

IPMK and β -catenin take part in PLC- β 1-dependent signaling pathway during myogenic differentiation

Giulia Ramazzotti¹, Irene Faenza¹, Anna Maria Billi¹, Stefano Ratti¹, Lucia Manzoli¹

¹Laboratorio di trasduzione del segnale, Dipartimento di Scienze Biomediche e Neuromotorie, Università di Bologna.

Phospholipase C (PLC)- β 1 catalytic activity plays an essential role in the initiation of myogenic differentiation but the effectors involved in its signaling pathway are not well defined[1,2]. Here, we show that the overexpression of the Inositol Polyphosphate Multikinase (IPMK) promotes myogenic differentiation, and that IPMK targets the same cyclin D3 promoter region activated by PLC- β 1. Moreover, cyclin D3 promoter activation relies upon c-jun binding to the promoter, both in response to PLC- β 1 and to IPMK overexpression. Furthermore, both IPMK and PLC- β 1 overexpression determines an increase in β -catenin translocation and accumulation to the nuclei of differentiating myoblasts resulting in higher MyoD activation. Therefore, our data show that PLC- β 1, IPMK and β -catenin are mediators of the same signaling pathway that regulates cyclin D3 and myosin heavy chain (MYH) induction during myogenic differentiation.

References

1. Faenza I, Ramazzotti G, Bavelloni A, Fiume R, Gaboardi GC, Follo MY, Gilmour RS, Martelli AM, Ravid K, Cocco L. Inositide-dependent phospholipase C signaling mimics insulin in skeletal muscle differentiation by affecting specific regions of the cyclin D3 promoter. *Endocrinology* [Internet]. 2007; 148: 1108–17. doi: 10.1210/en.2006-1003
2. Ramazzotti G, Faenza I, Gaboardi GC, Piazzini M, Bavelloni A, Fiume R, Manzoli L, Martelli AM, Cocco L. Catalytic activity of nuclear PLC-beta(1) is required for its signalling function during C2C12 differentiation. *Cell Signal* [Internet]. 2008; 20: 2013–21. doi: 10.1016/j.cellsig.2008.07.009

Keywords

Myogenic differentiation, phospholipase C- β 1, IPMK, β -catenin, inositol phosphates