

FACTA UNIVERSITATIS (NIŠ)
SER. MATH. INFORM. Vol. 33, No 1 (2018), 63–77
<https://doi.org/10.22190/FUMI1801063K>

AN ALTERNATIVE DECOMPOSITION OF CATALAN NUMBER *

Predrag V. Krtolica, Predrag S. Stanimirović and Igor Stojanović

Abstract. A particular integer sequence derived by the convex polygon triangulation is introduced and investigated. After some underlying results are presented, the forbidden (or improper) integer values relative to the triangulation are concerned. It is understood that the forbidden sequences do not correspond to any triangulation. Some of their properties are presented. These properties are used to count the forbidden values, which is, finally, exploited in stating another decomposition of the Catalan number.

Keywords: Catalan number, Convex polygon triangulation, Binomial coefficient.

1. Introduction and Preliminaries

The decomposition of two-dimensional objects into simpler components has a variety of applications in computer graphics as well as in different areas of engineering (robotics, pattern recognition and others mentioned in [3]).

The problem of the convex polygon triangulation assumes dividing a given polygon area by its internal diagonals into the triangles without any gaps and overlaps. This is a classical problem solved so far in several ways.

The number of triangulations of a convex n -gon is equal to the $(n-2)$ -th Catalan number, i.e.

$$(1.1) \quad C_{n-2} = \frac{1}{n-1} \binom{2n-4}{n-2}.$$

In [9] we have proposed a recursive convex polygon triangulation algorithm, partially based on the problem solution proposed in [7] where the context-free grammar is used.

Received March 21, 2017; accepted March 13, 2018

2010 *Mathematics Subject Classification.* Primary 11Y55; Secondary 68R05, 68U05

*The authors were supported in part by Research Project 174013 of the Serbian Ministry of Science and Project "Applying direct methods for digital image restoring" of the Goce Delčev University

Being inspired by the method described in [7], we have used the following rules generating the arithmetic expression in the reverse Polish notation.

$$(1.2) \quad S \rightarrow SS +$$

$$(1.3) \quad S \rightarrow b$$

Using the grammar rules (1.2) and (1.3) and the reverse Polish notation (RPN) method (cf. [8, 9, 10]) the algorithm for a convex polygon triangulation is suggested in [9]. Also, it is proved that this algorithm works in linear time. Moreover, the correlation between the arithmetic expressions in RPN with triangulations is investigated.

In [9], for every $n \geq 3$, we can consider the mapping $F_n : R_n \mapsto T_n$, whose domain is the set of expressions made by the $n - 2$ applications of the rule (1.2) and $n - 1$ applications of the rule (1.3), and the range is the set of an n -gon triangulations. It is not difficult to verify that some elements from R_n could be derived in multiple ways. The unique characterization for each element in R_n is given in [9].

Following the usual notations, by $\{+, b\}_{p,q}^*$ we denote the subset of the closure set $\{+, b\}^*$ consisting of p appearances of the character $+$ and q appearances of the character b . In the boundary case, we have $\{+, b\}_{0,0}^* = \{\epsilon\}$.

Lemma 1.1. [9] *An arbitrary element $r_n \in R_n$, corresponding to a particular triangulation of an n -sided polygon, is uniquely determined by the following two conditions:*

(C1) *It possesses the form*

$$r_n = bbA+, \quad A \in \{+, b\}_{n-3, n-3}^*$$

(C2) *Each initial part of the expression r_n (the substring of consecutive characters which starts from the first character) must be of the form*

$$\{+, b\}_{p,q}^* \quad p < q, \quad p = 1, \dots, n - 2, \quad q = 1, \dots, n - 1.$$

By P_n we denote the set of expressions generated by the grammar rules (1.2), (1.3) which satisfy conditions (C1) and (C2).

Lemma 1.2. [9] *The mapping $F_n : P_n \mapsto T_n$ is well defined, one-to-one and onto for any $n \geq 3$.*

In this paper, we assign the unique integer to every arithmetic expression in RPN, derived by rules (1.2) and (1.3), which corresponds to some n -gone triangulation. In other words, we have made a bijection between triangulations T_n and a subset of integers. We consider these numbers and give some of their properties.

As a consequence, we get an ordering in the set T_n , different than e.g. the one derived in [5]. In [5] the authors propose the infinite tree of all triangulations, where every tree node is a particular triangulation of an n -gon ($n \geq 3$) with a number

of descendants equal to the degree of the vertex n in range $2, \dots, n-1$. The descendant triangulations (which are the $(n+1)$ -gon triangulations) are derived by "splits" of diagonals $\delta_{i,n}$, $i \in \{1, \dots, n-1\}$, incident to the vertex n . If these splits are performed in the increasing order of i , the unique order among all triangulations (and unique tree of all triangulations) is derived.

In the following section, we define the forbidden combinations (which do not correspond to any element of T_n) and count their number.

There is a variety of recursive definitions of Catalan numbers in literature. Many of them are derived using their combinatorial essence. For example, a well known Touchard's formula (see e.g. [6]) is derived using the generating function for Catalan numbers. Also, Hurtado in [4] gives another recursive definition counting the triangulations by the ears. Stanimirović et al. in [13] introduce Catalan matrix and derive several combinatorial identities with Catalan numbers.

Catalan numbers appear not only in counting the polygon triangulations, but in lattice paths, ballot problem and several tents of other situations which can be found summarized in [14].

They have even a broader application concerning many problems in physics and engineering (see e.g. [2]). Integer sequences themselves are also applicable in variety of areas as queueing theory and probability theory (see [1]) and many others. An interesting application of Catalan numbers sequence in cryptography can be found in [11].

Actually, the present literature on Catalan numbers is far more exhaustive, containing many different formulas for Catalan numbers derived in the variety of approaches and contexts.

Finally, using the combinatorial argument and definition of the forbidden combinations, we give another recursive definition of Catalan numbers.

2. Arithmetic Expressions and Integer Sequences

It is obvious that by changing b 's with 1's and $+$'s with 0's we get a unique binary number for every triangulation of the convex polygon.

Denote the integer value corresponding to the i th triangulation from T_n by d_n^i . It is of our interest to consider sets

$$D_n = \{d_n^i \mid i = 1, \dots, C_{n-2}\}, \quad D = \bigcup_{n \geq 3} D_n.$$

The condition (C1) of Lemma 1.1 imposes that each of those numbers starts with '11' and ends with '0'. If so, we could consider only the central part of each d_n^i . Denote this binary number by $[x_n^i]$ and its value by x_n^i , where $i \in \{1, \dots, C_{n-2}\}$

corresponds to a particular triangulation. Similarly, we will use the sets

$$X_n = \{x_n^i \mid i = 1, \dots, C_{n-2}\}, \quad X = \bigcup_{n \geq 3} X_n.$$

The expressions P_n and corresponding numbers for smaller values of n are presented in Table 2.1.

Table 2.1: Sets P_n , D_n and X_n .

n	P_n	$[p_n^i]$	D_n	X_n
3	$bb+$	110	6	0
4	$bbb++$	11100	28	2
	$bb+b+$	11010	26	1
5	$bbbb+++$	1111000	120	12
	$bbb+b++$	1110100	116	10
	$bbb++b+$	1110010	114	9
	$bb+bb++$	1101100	108	6
	$bb+b+b+$	1101010	106	5
6	$bbbbbb++++$	111110000	496	56
	$bbbb+b+++$	111101000	488	52
	$bbbb++b++$	111100100	484	50
	$bbbb+++b+$	111100010	482	49
	$bbb+bb+++$	111011000	472	44
	$bbb+b+b++$	111010100	468	42
	$bbb+b++b+$	111010010	466	41
	$bbb++bb++$	111001100	460	38
	$bbb++b+b+$	111001010	458	37
	$bb+bbbb+++$	110111000	440	28
	$bb+bb+b++$	110110100	436	26
	$bb+bb++b+$	110110010	434	25
	$bb+b+bb++$	110101100	428	22
	$bb+b+b+b+$	110101010	426	21

It is interesting that the sequence X appears in [12] noted as A085183, derived from the terms of sequence A014486 without their most significant bit (1) and the least significant bit (0). The sequence A014486 represents a list of totally balanced sequences of $2n$ binary digits written in base 10. A binary expansion of each term from A014486 contains n 0s and n 1s and reading from left to right (the most significant to the least significant bit), the number of 0s never exceeds the number of 1s [12].

Also, note that among famous Stanley's 66 combinatorial interpretations of Catalan numbers ([14]) there is no example identical to the sequence X_n . Of course,

being counted by Catalan numbers, a bijection can be established between all of them.

Now, we investigate some basic properties of the sets D and X .

Proposition 2.1. *The value of the number corresponding to the postfix expression characteristic for particular triangulation of the convex n -gon is*

$$d_n^i = 2(2^{2n-5} + 2^{2n-6} + x_n^i),$$

where x_n^i is the changeable part or a shorter binary (decimal) equivalent.

Vice versa

$$x_n^i = \frac{d_n^i - 2^{2n-4} - 2^{2n-5}}{2}.$$

Proof. It follows immediately from Lemma 1.1. \square

In order to clarify Table 2.1, in Figure 2.1 we present the relation between the arithmetic expression $bbbb+++$ from P_5 , derived by productions (1.2) and (1.3), and producing the pentagon triangulation (for the algorithm see [9]). The rest of arithmetic expressions and the corresponding triangulations for $n = 5$ is presented in Figure 2.2.





FIG. 2.1: Generating the triangulation corresponding to $bbbb+++$.

It is well known that there is a bijection between the triangulations and binary trees. It is interesting to note that triangulation ordering imposed by sequence X_n does not match the usual ordering between the trees corresponding to the particular triangulation (see e.g. [6, Chapter 8]). This can be easily seen in Figure 2.3. In the first column, corresponding arithmetic expression in RPN is presented, then there is the corresponding parsing tree, followed by triangulation and the binary tree for this triangulation. In five rows in Figure 2.3 we have five possible triangulations of a pentagon corresponding to integers $x_5^i = 5, 6, 9, 10, 12$, $i = 1, 2, 3, 4, 5$ respectively.

Let us give some observations about $[x_n^i]$. According to Lemma 1.1, it is obvious that the length of each $[x_n^i]$ is equal to $|[x_n^i]| = 2n - 6$ and $[x_n^i] \in \{0, 1\}_{n-3, n-3}^*$. Moreover, $[x_n^i]$ can not start with more than one '0' and can not end with more than one '1'. So, it must have one of the forms $[y_1(n)], \dots, [y_9(n)]$ presented in Table 2.2.

FIG. 2.2: Illustration of the rest from T_5 and P_5 .

In Table 2.2

$$\begin{aligned} [\alpha] &\in \{0, 1\}_{n-6, n-4}^* \\ [\beta] &\in \{0, 1\}_{n-5, n-5}^* \\ [\gamma] &\in \{0, 1\}_{n-4, n-6}^*. \end{aligned}$$

Note that some of $[\alpha], [\beta], [\gamma]$ type strings are proper (correspond to particular polygon triangulations) while some of them are not. These strings are proper under conditions described below.

The $[\alpha]$ type string can be found only within the contexts

$$1101[\alpha]000, 1110[\alpha]000$$

and, according to the conditions of Lemma 1.1, start of proper $[\alpha]$ should satisfy

$$(2.1) \quad [\alpha]_{ps} \in \{0, 1\}_{p, q}^*, \quad p \leq q + 1, \quad p \in \{0, \dots, n - 6\}; \quad q \in \{0, \dots, n - 4\}.$$

The substring of $[\beta]$ type has to be considered in two subtypes: $[\beta']$ and $[\beta'']$. The subtype $[\beta']$ can be found in the contexts

$$1101[\beta']010, 1101[\beta']100, 1110[\beta']010, 1110[\beta']100,$$

and, according to Lemma 1.1, every start of proper $[\beta']$ should satisfy

$$(2.2) \quad [\beta']_{ps} \in \{0, 1\}_{p, q}^*, \quad p \leq q + 1, \quad p \in \{0, \dots, n - 5\}; \quad q \in \{0, \dots, n - 5\}.$$

Table 2.2: String types.

weights	$2n - 7$	$2n - 8$...	1	0
$[y_1(n)]$	0	1	$[\alpha]$	0	0
$[y_2(n)]$	0	1	$[\beta]$	0	1
$[y_3(n)]$	0	1	$[\beta]$	1	0
$[y_4(n)]$	1	0	$[\alpha]$	0	0
$[y_5(n)]$	1	0	$[\beta]$	0	1
$[y_6(n)]$	1	0	$[\beta]$	1	0
$[y_7(n)]$	1	1	$[\beta]$	0	0
$[y_8(n)]$	1	1	$[\gamma]$	0	1
$[y_9(n)]$	1	1	$[\gamma]$	1	0

The subtype $[\beta'']$ should be found within context

$$1111[\beta'']000$$

and every start of proper $[\beta'']$ should satisfy

$$(2.3) \quad [\beta'']_{ps} \in \{0, 1\}_{p,q}^*, \quad p \leq q + 3, \quad p \in \{0, \dots, n - 5\}; \quad q \in \{0, \dots, n - 5\}.$$

Because of the fact that $[\gamma]$ type strings can be found only within the contexts

$$1111[\gamma]010, \quad 1111[\gamma]100$$

and the conditions of Lemma 1.1, it is clear that every start of proper $[\gamma]$ should satisfy

$$(2.4) \quad [\gamma]_{ps} \in \{0, 1\}_{p,q}^*, \quad p \leq q + 3, \quad p \in \{0, \dots, n - 4\}; \quad q \in \{0, \dots, n - 6\}.$$

The number of the bits in $[\alpha]$, $[\beta]$ and $[\gamma]$ is

$$|[\alpha]| = |[\beta]| = |[\gamma]| = 2n - 10$$

and values of the above binary numbers $[y_i(n)]$ are correspondingly:

$$(2.5) \quad y_1(n) = 2^{2n-8} + 4\alpha$$

$$(2.6) \quad y_2(n) = 2^{2n-8} + 4\beta + 1$$

$$(2.7) \quad y_3(n) = 2^{2n-8} + 4\beta + 2$$

$$(2.8) \quad y_4(n) = 2^{2n-7} + 4\alpha$$

$$(2.9) \quad y_5(n) = 2^{2n-7} + 4\beta + 1$$

$$(2.10) \quad y_6(n) = 2^{2n-7} + 4\beta + 2$$

$$(2.11) \quad y_7(n) = 2^{2n-7} + 2^{2n-8} + 4\beta$$

$$(2.12) \quad y_8(n) = 2^{2n-7} + 2^{2n-8} + 4\gamma + 1$$

$$(2.13) \quad y_9(n) = 2^{2n-7} + 2^{2n-8} + 4\gamma + 2$$

Proposition 2.2. *The minimal value among the values (2.5)-(2.13) is*

$$y_{\min}(n) = \frac{4^{n-3} - 1}{3}, \quad n \geq 4.$$

Proof. It is not difficult to observe that the minimal value $y_{\min}(n)$ must have the form $01[\beta_{\min}]01$, where $[\beta_{\min}]$ corresponds to the minimal value in the context $01[\beta]01$. Note that β_{\min} is not the minimal β in general.

As a consequence of Lema 1.1 and its condition (C2), every beginning of β , $\beta_s \in \{0, 1\}_{p,q}^*$, satisfies the condition $p \leq q + 1$. The β_{\min} for an n -gon (denoted as $\beta_{\min}(n)$) is of the following form:

$$\begin{aligned} \beta_{\min}(n) &= \epsilon, & 3 \leq n \leq 5 \\ \beta_{\min}(6) &= 01 = 1_{10}, \\ \beta_{\min}(7) &= 0101 = 5_{10}, \\ \beta_{\min}(n > 7) &= \overbrace{01}^{n-5} = \sum_{k=1}^{n-5} 2^{2(k-1)}. \end{aligned}$$

Therefore

$$\begin{aligned} y_{\min}(n) &= 2^{2n-8} + 4 \sum_{k=1}^{n-5} 2^{2(k-1)} + 1 \\ &= 2^{2n-8} + 4 \frac{2^{2n-10} - 1}{3} + 1 \\ &= 4^{n-4} + 4 \frac{4^{n-5} - 1}{3} + 1 \\ &= \frac{3 \cdot 4^{n-4} + 4 \cdot 4^{n-5} - 4 + 3}{3} \\ &= \frac{4^{n-3} - 1}{3}. \end{aligned}$$

□

Proposition 2.3. *The maximal value among the values (2.5)-(2.13) is*

$$y_{\max}(n) = 2^{n-3} (2^{n-3} - 1).$$

Proof. It is straightforward to conclude that the maximal value must have the form $11[\beta_{\max}]00$, where $[\beta_{\max}]$ corresponds to the maximal value in the context $11[\beta]00$, since it starts with two 1's and β has more 1's than γ .

The maximal β (which is allowed by Lemma 1.1) is

$$[\beta_{\max}] = \underbrace{1 \dots 1}_{n-5} \underbrace{0 \dots 0}_{n-5} = 1^{n-5} 0^{n-5}.$$

The corresponding value is

$$\beta_{\max} = 2^{n-5} \sum_{k=0}^{n-6} 2^k = 2^{n-5}(2^{n-5} - 1)$$

Therefore, using

$$y_{\max}(n) = 2^{2n-7} + 2^{2n-8} + 4\beta_{\max}$$

and the elementary algebraic operations the proof can be completed. \square

Proposition 2.4. *The following inequality is satisfied for $n \geq 3$:*

$$y_{\min}(n+1) > y_{\max}(n).$$

Proof. According to Propositions 2.2 and 2.3

$$\begin{aligned} y_{\min}(n+1) - y_{\max}(n) &= \frac{4^{n-2} - 1}{3} - 2^{n-3}(2^{n-3} - 1) \\ &= \frac{a^2 + 3a - 1}{3}, \end{aligned}$$

where $a = 2^{n-3}$. As $n \geq 4$ and $a \geq 2$, it is not difficult to verify that the quadratic inequality

$$a^2 + 3a - 1 > 0$$

is satisfied. \square

As we have already seen, some combinations are forbidden (do not correspond to any n -gon triangulation). Later, we count these forbidden combinations. In accordance to the previous results we state the next definition.

Definition 2.1. The forbidden starts are defined in the following way:

(a) By the forbidden start of $[\alpha]$ we consider

$$[\alpha]_{fs} \in \{0, 1\}_{p,q}^*, \quad p > q + 1,$$

where all its starts with length less than $p + q$ are proper.

(b) By the forbidden start of $[\beta']$ we consider

$$[\beta']_{fs} \in \{0, 1\}_{p,q}^*, \quad p > q + 1,$$

where all its starts with length less than $p + q$ are proper.

(c) The forbidden start of $[\beta'']$ is given by

$$[\beta'']_{fs} \in \{0, 1\}_{p,q}^*, \quad p > q + 3,$$

where all its starts with length less than $p + q$ are proper.

(d) By the forbidden start of $[\gamma]$ we consider

$$[\gamma]_{fs} \in \{0, 1\}_{p,q}^*, p > q + 3,$$

where all its starts with length less than $p + q$ are proper.

Proposition 2.5. *Any combination having the forbidden start is also forbidden.*

Remark 2.1. *The forbidden combination is not a proper string $\{0, 1\}_{n-3, n-3}^*$ which corresponds to some n -gon triangulation. The purpose of the forbidden start notion is to count the non-proper combinations which do not correspond to any triangulation. If we recognize a forbidden start, then any pattern of bits following it makes a certain number of forbidden combinations. This will be used later in Theorems 2.1 and 2.2.*

The following lemmas hold.

Lemma 2.1. *If there exists a forbidden start in $[\alpha]$, $[\beta]$ or $[\gamma]$ it must contain an even number of characters.*

Proof. According to (2.1) and (2.2), the shortest forbidden start for $[\alpha]$ and $[\beta']$ is '00', where $p + q = 2$, an even integer. Similarly, according to (2.3) and (2.4), the shortest forbidden start for $[\beta'']$ and $[\gamma]$ is '0000', in which case holds $p + q = 4$.

The case $p + q = 2q + 1$ is the limit case for allowed start in (2.1) and (2.2). The forbidden start can grow from it by adding one zero at the end, making the length of this forbidden start an even number $p + 1 + q = 2q + 2$.

For all allowed starts of $[\beta'']$ and $[\gamma]$ holds $p \leq q + 3$. The forbidden start can be made by appending one zero only for the case $p = q + 3$. Therefore, $p + q = 2q + 3$ must be satisfied again. In this case, the forbidden start has the even length $p + 1 + q = 2q + 4$. \square

Lemma 2.2. *The following holds:*

(a) *If we have a forbidden start $2k$ characters long in $[\alpha]$ or $[\beta']$, then it contains exactly $k - 1$ 1s.*

(b) *If we have a forbidden start $2k$ characters long in $[\beta'']$ or $[\gamma]$, then it contains exactly $k - 2$ 1s.*

Proof. (a) Let us consider firstly the $[\alpha]$ type case. As $p + q = 2k$ and (2.) are satisfied, we have $p = q + 2$ which implies $q + 2 + q = 2k$ and the proof is completed.

The rest of the proof could be done in a similar way. \square

Remark 2.2. It is easy to observe that the shortest forbidden start '00' for $[\alpha]$ occurs at $n = 8$. Regarding $[\alpha] \in \{0, 1\}_{n-6, n-4}^*$, this forbidden start can be found in the combinations 1101001111000 or 1110001111000.

Also, for $[\beta']$ the shortest forbidden start '00' occurs at $n = 7$ in 11010011010, 11010011100, 11100011010 or 11100011100.

In the case of $[\beta'']$ the shortest forbidden start is '0000', at $n = 9$ in combination 111100001111000.

The shortest forbidden start '0000' for $[\gamma]$ occurs at $n = 8$ in

$$1111 \underline{000011} 010$$

or

$$1111 \underline{000011} 100.$$

Theorem 2.1. The number of forbidden combinations for $[\alpha]$ is

$$n_{\alpha_{fs}}(n) = \binom{2n-10}{n-8}, \quad n \geq 8.$$

The number of forbidden combinations for $[\beta']$ is

$$n_{\beta'_{fs}}(n) = \binom{2n-10}{n-7}, \quad n \geq 7.$$

The number of forbidden combinations for $[\beta'']$ and $[\gamma]$ is

$$n_{\beta''_{fs}}(n) = \binom{2n-10}{n-9}, \quad n \geq 9,$$

and

$$n_{\gamma_{fs}}(n) = \binom{2n-10}{n-8}, \quad n \geq 8,$$

respectively.

Proof. As $[\alpha]_{fs} \in \{0, 1\}_{p,q}^*$, $p > q + 1$, and $[\alpha] \in \{0, 1\}_{n-6, n-4}^*$, we actually want to count all paths from $(0, 0)$ to $(n-6, n-4)$ touching the line $p = q + 2$. Using the reflection principle, this is the same number as the number of paths from $(2, -2)$ to $(n-6, n-4)$. This yields

$$n_{\alpha_{fs}}(n) = \binom{n-6+n-4}{n-6-2} = \binom{2n-10}{n-8}.$$

In a similar way, we can complete the rest of the proof. Note that for $[\beta']_{fs} \in \{0, 1\}_{p,q}^*$, $p > q + 1$ and $[\beta] \in \{0, 1\}_{n-5, n-5}^*$ we need to observe count of paths from

$(2, -2)$ to $(n - 5, n - 5)$, while for $[\beta'']_{fs} \in \{0, 1\}_{p,q}^*$, $p > q + 3$, we observe line $p = q + 4$ and paths from $(4, -4)$ to $(n - 5, n - 5)$.

For $[\gamma]_{fs} \in \{0, 1\}_{p,q}^*$, $p > q + 3$, and $[\gamma] \in \{0, 1\}_{n-4, n-6}^*$ we need to observe line $p = q + 4$ and paths from $(4, -4)$ to $(n - 4, n - 6)$. \square

In the next theorem, we state another way to express Catalan number.

Theorem 2.2.

$$(2.14) \quad C_{n-2} = 4 \binom{2n-10}{n-6} + 5 \binom{2n-10}{n-5} - 4 \binom{2n-10}{n-8} - 4 \binom{2n-10}{n-7} - \binom{2n-10}{n-9}, \quad n \geq 9.$$

Proof. The proof could be derived in the combinatorial way. The overall number of possible combinations in the case of $[\alpha]$ and $[\gamma]$ type strings from Table 2.2 is $4 \binom{2n-10}{n-6}$, while the number of possible combinations in the case of $[\beta]$ is $5 \binom{2n-10}{n-5}$.

Theorem 2.1 counts all forbidden combinations. The overall number of possible combinations diminished by the number of the forbidden combinations results in the number of the allowed combinations, which is equal to C_{n-2} . \square

Corollary 1. *The following holds*

$$\begin{aligned} C_{n-2} &= 8 \frac{8n^3 - 84n^2 + 286n - 315}{(n-1)(n-2)(n-3)(n-4)} \binom{2n-10}{n-5} \\ &= 8 \frac{8n^3 - 84n^2 + 286n - 315}{(n-1)(n-2)(n-3)} C_{n-5}. \end{aligned}$$

Proof. Using Cardan's rule of succession

$$\binom{n}{r} = \frac{n-r+1}{r} \binom{n}{r-1}$$

each binomial coefficient from (2.14) can be reduced on the same type, e.g. $\binom{2n-10}{n-5}$. After some algebraic transformations we get the result. \square




FIG. 2.3: Illustrations of arithmetic expressions, expression trees, triangulations and binary trees.

3. Conclusion

We have made the presentation of the convex polygon triangulations in the form of integers. This imposes one of the triangulation ordering different from e.g. the one derived in [5].

Moreover, the boundaries for these integers are found in terms of polygon vertices number n . The forbidden patterns in the binary presentation of these integers are counted. Another way to express Catalan number via binomial coefficients is presented.

Acknowledgments

The authors wish to express their gratitude to prof. Dragan Stevanović who read the draft of this paper and gave a number of useful comments and suggestions. His effort provides a considerable contribution to the quality of this paper.

REFERENCES

1. J. Abate, W. Whitt, *Integer sequences from queueing theory*, Journal of Integer Sequences, **13** (2010), 21p.
2. J. Abate, W. Whitt, *Brownian motion and the generalized Catalan numbers*, Journal of Integer Sequences, **14** (2011), 15p.
3. B. Chazelle, L. Palios, *Decomposition algorithms in geometry*, Algebraic Geometry and its Application (ed. Ch. L. Bajaj) **27**, (1994), 419–447.
4. F. Hurtado, M. Noy, *Ears of triangulations and Catalan numbers*, Discrete Mathematics, **149** (1996), 319–324.
5. F. Hurtado, M. Noy, *Graph of triangulations of a convex polygon and tree of triangulations*, Comput. Geom., **13** (1999), 179–188.
6. T. Koshy, *Catalan numbers with applications*, Oxford University Press, Oxford, New York, 2009.
7. M. Kross, A. Lentin, *Notions sur les grammaires formelles*, Gauthier-Villars, 1967.
8. P. V. Krtolica, P. S. Stanimirović, *On Some properties of Reverse Polish Notation*, FILOMAT, **13** (1999), 157–172.
9. P. V. Krtolica, P. S. Stanimirović, R. Stanojević, *Reverse Polish notation in constructing the algorithm for polygon triangulation*, FILOMAT, **15** (2001), 25–33.
10. P. V. Krtolica, P. S. Stanimirović, *Reverse Polish notation method*, Int. J. Comput. Math., **81**(3) (2004), 273–284.
11. D. Sravana Kumar, CH. Suneetha, A. Chandrasekhar, *Novel encryption schemes based on Catalan numbers*, International Journal of Engineering Research and Applications (IJERA), **2**(2) (2012), 161–166.
12. *The On-Line Encyclopedia of Integer Sequences (OEIS)*. <http://oeis.org/Seis.html>.

13. S. Stanimirović, P. Stanimirović, M. Miladinović, A. Ilić, *Catalan matrix and related combinatorial identities*, Appl. Math. Comput., **215** (2009), 796-805.
14. R. P. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge University Press, Cambridge, 1999.

Predrag Krtolica
University of Niš, Faculty of Science and Mathematics
Department of Computer Science
Višegradaska 33, 18000 Niš, Serbia
`krca@pmf.ni.ac.rs`

Predrag Stanimirović
University of Niš, Faculty of Science and Mathematics
Department of Computer Science
Višegradaska 33, 18000 Niš, Serbia
`pecko@pmf.ni.ac.rs`

Igor Stojanović
Goce Delčev University, Faculty of Computer Science
Goce Delčev 89, 2000 Štip, Macedonia
`igor.stojanovik@ugd.edu.mk`