

FACTA UNIVERSITATIS (NIŠ)  
SER. MATH. INFORM. Vol. 30, No 5 (2015), 741–752

## CURVATURE TENSORS AND PSEUDOTENSORS IN A GENERALIZED FINSLER SPACE\*

Milan Lj. Zlatanović, Svetislav M. Minčić, Miloš Z. Petrović

**Abstract.** We examine relations between the curvature tensor of the associated symmetric connection and curvature tensors, curvature pseudotensors and derived curvature tensors of non-symmetric affine connection in Rund's sense.

**Keywords:** curvature tensor; curvature pseudotensor; derived curvature tensor; generalized Finsler space; non-symmetric connection.

### 1. Introduction and preliminaries

Eisenhart [3] gave a generalization of Riemannian spaces by using (in general) the non-symmetric basic tensor. In the same way Shamihoke [11]–[14] gave a generalization of Finsler spaces. He introduced two kinds of generalized Finsler spaces. The first kind was a generalization of Finsler spaces in Rund's [10] sense and the second kind was a generalization of Cartan's [1] Finsler spaces. In the present paper, we follow Shamihoke's definition of generalized Finsler spaces of the first kind. We continue the results derived in several previously published papers [2, 7, 8, 9, 15].

A generalized Finsler space  $GF_N$  is a differentiable manifold equipped with a basic tensor  $g_{ij}(x^1, \dots, x^N, \dot{x}^1, \dots, \dot{x}^N) \equiv g_{ij}(x, \dot{x})$ , where

$$(1.1) \quad g_{ij}(x, \dot{x}) \neq g_{ji}(x, \dot{x}), \text{ in generally, } \quad (g = \det(g_{ij}) \neq 0, \dot{x} = dx/dt).$$

Based on (1.1), one can define the symmetric and the anti-symmetric part of  $g_{ij}$ , respectively, by

$$g_{\underline{ij}} = \frac{1}{2}(g_{ij} + g_{ji}), \quad g_{\underset{\vee}{ij}} = \frac{1}{2}(g_{ij} - g_{ji}),$$

---

Received July 10, 2015; Accepted August 14, 2015

2010 *Mathematics Subject Classification.* Primary 53A45; Secondary 53B05, 53B40

\*The authors gratefully acknowledge support from the research project 174012 of the Serbian Ministry of Science.

where, following [13], it is

$$(1.2) \quad a) g_{ij}(x, \dot{x}) = \frac{1}{2} \frac{\partial^2 F^2(x, \dot{x})}{\partial \dot{x}^i \partial \dot{x}^j}, \quad b) \frac{\partial g_{ij}}{\partial \dot{x}^k} = 0.$$

The function  $F(x, \dot{x})$  is a metric function in  $GF_N$ , having the properties known from the theory of usual Finsler spaces ( $F_N$ ). The following conditions are valid:

1.  $F(x, \dot{x})$  is continuously differentiable at least four times in its  $2N$  arguments.
2.  $F(x, \dot{x}) > 0$  providing all  $dx^i$  are not 0.
3.  $F(x, \dot{x})$  is positively homogeneous of the 1<sup>st</sup> degree in  $\dot{x}$ , i.e.

$$F(x, k\dot{x}) = kF(x, \dot{x}), \quad k > 0.$$

4.  $\frac{\partial^2 F^2(x, \dot{x})}{\partial \dot{x}^i \partial \dot{x}^j} \xi^i \xi^j > 0$  for any given  $\dot{x}$ , and  $\sum_i (\xi^i)^2 > 0$ ,  $\xi^i \in R$ .

The lowering and the raising of indices are defined by the tensors  $g_{ij}$  and  $h^{ij}$ , respectively, where  $h^{ij}$  is defined as follows

$$g_{ij} h^{jk} = \delta_i^k, \quad (g = \det(g_{ij}) \neq 0).$$

Generalized Cristoffel symbols of the 1<sup>st</sup> and the 2<sup>nd</sup> kind are defined by

$$\begin{aligned} \gamma_{i,jk} &= \frac{1}{2} (g_{ji,k} - g_{jk,i} + g_{ik,j}) \neq \gamma_{i,kj}, \\ \gamma_{jk}^i &= h^{ip} \gamma_{p,jk} = \frac{1}{2} h^{ip} (g_{jp,k} - g_{jk,p} + g_{pk,j}) \neq \gamma_{kj}^i, \end{aligned}$$

where,  $g_{ji,k} = \partial g_{ji} / \partial x^k$ .

Then we have

$$\gamma_{jk}^p g_{ip} = \gamma_{s,jk} h^{ps} g_{ip} = \gamma_{s,jk} \delta_i^s = \gamma_{i,jk}.$$

Introducing the tensor  $C_{ijk}$  analogously as in  $F_N$ , we have

$$(1.3) \quad C_{ijk}(x, \dot{x}) \stackrel{def}{=} \frac{1}{2} g_{ij,\dot{x}^k} \stackrel{(1.2b)}{=} \frac{1}{2} g_{ij,\dot{x}^k} \stackrel{(1.2a)}{=} \frac{1}{4} F_{\dot{x}^i \dot{x}^j \dot{x}^k}^2,$$

where " $\stackrel{(1.2b)}{=}$ " signifies "equal based on (1.2b)".

From (1.3) we can conclude that  $C_{ijk}$  is symmetric with respect to each pair of indices. Also, we have

$$C_{jk}^i \stackrel{def}{=} h^{ip} C_{pjk} = h^{ip} C_{jpk} = h^{ip} C_{jkp}.$$

With the help of coefficients  $P^i_{jk'}$  given by

$$P^i_{jk} = \gamma^i_{jk} - C^i_{jp} \gamma^p_{sk} \dot{x}^s \neq P^i_{kj'}$$

one obtains coefficients of non-symmetric affine connection in Rund's sense [14]:

$$P^*_{jk} = \gamma^i_{jk} - h^{iq}(C_{jqp}P^p_{ks} + C_{kqp}P^p_{js} - C_{jkp}P^p_{qs})\dot{x}^s \neq P^*_{kj'}$$

$$P^*_{i,jk} = P^*_{jk}g_{i\bar{r}} = \gamma^{i,jk} - (C_{ijp}P^p_{ks} + C_{ikp}P^p_{js} - C_{jkp}P^p_{is})\dot{x}^s \neq P^*_{i.kj}$$

In  $GF_N$  we denote the double anti-symmetric and the double symmetric part of connection  $P^*_{jk}$ , respectively, by:

$$a) T^*_{jk}(x, \dot{x}) = P^*_{[jk]} = P^*_{jk} - P^*_{kj'}, \quad b) P^*_{(jk)} = P^*_{jk} + P^*_{kj'}$$

where  $T^*_{jk}$  is the *torsion tensor*.

## 2. Curvature tensors and pseudotensors of non-symmetric connection in Rund's sense

In the generalized Finsler space  $GF_N$  based on the non-symmetry of connection coefficients  $P^*_{jk}$  there exist four kinds of covariant derivative of a tensor (see, e.g.[7]–[9]). For a tensor  $a^i_j(x, \dot{x})$  it is

$$(2.1) \quad a^i_{j|m}(x, \dot{x}) = \delta_m a^i_j + P^*_{pm} a^p_j - P^*_{jm} a^i_p,$$

$\begin{matrix} 1 \\ 2 \\ 3 \\ 4 \end{matrix}$

$\begin{matrix} mp \\ pm \\ mp \end{matrix}$

$\begin{matrix} mj \\ mj \\ jm \end{matrix}$

where

$$\delta_m = \frac{\partial}{\partial x^m} + \frac{\partial \dot{x}^p}{\partial x^m} \frac{\partial}{\partial \dot{x}^p}$$

By using the first and the second kind of covariant derivative, given in (2.1), we examine ten Ricci type identities [8]

$$\begin{aligned}
a_{j|_1mn}^i - a_{j|_1nm}^i &= R_{1\ pmn}^{*i} a_j^p - R_{1\ jmn}^{*p} a_p^i - T_{nm}^{*p} a_{j|_1p}^i, \\
a_{j|_2mn}^i - a_{j|_2nm}^i &= R_{2\ pmn}^{*i} a_j^p - R_{2\ jmn}^{*p} a_p^i + T_{nm}^{*p} a_{j|_2p}^i, \\
a_{j|_1\ 2}^i - a_{j|_1\ 2}^i &= A_{1\ pmn}^i a_j^p - A_{2\ jmn}^p a_p^i + a_{j<[mn]>}^i + a_{j\leq[mm]\geq}^i + T_{nm}^{*p} a_{j|_1p}^i, \\
a_{j|_2\ 1}^i - a_{j|_2\ 1}^i &= A_{3\ pmn}^i a_j^p - A_{4\ jmn}^p a_p^i - a_{j<[mn]>}^i - a_{j\leq[mm]\geq}^i - T_{nm}^{*p} a_{j|_2p}^i, \\
a_{j|_1mn}^i - a_{j|_2nm}^i &= A_{5\ pmn}^i a_j^p - A_{6\ jmn}^p a_p^i + a_{j<(mn)>}^i + a_{j\leq(mn)\geq}^i - P_{mn}^{*p} (a_{j|_1p}^i - a_{j|_2p}^i), \\
a_{j|_1mn}^i - a_{j|_2nm}^i &= A_{7\ pmn}^i a_j^p - A_{8\ jmn}^p a_p^i + a_{j<mn>}^i + a_{j\leq mn\geq}^i, \\
a_{j|_1mn}^i - a_{j|_2nm}^i &= A_{9\ pmn}^i a_j^p - A_{10\ jmn}^p a_p^i + a_{j<nm>}^i + a_{j\leq nm\geq}^i - (P_{mn}^{*p} a_{j|_1p}^i - P_{nm}^{*p} a_{j|_2p}^i), \\
a_{j|_2mn}^i - a_{j|_2nm}^i &= A_{11\ pmn}^i a_j^p - A_{12\ jmn}^p a_p^i - a_{j<nm>}^i + a_{j\leq mn\geq}^i + P_{mn}^{*p} a_{j|_1p}^i - P_{nm}^{*p} a_{j|_2p}^i, \\
a_{j|_2mn}^i - a_{j|_2nm}^i &= A_{13\ pmn}^i a_j^p - A_{14\ jmn}^p a_p^i - a_{j<nm>}^i + a_{j\leq nm\geq}^i, \\
a_{j|_1\ 2}^i - a_{j|_2\ 1}^i &= A_{15\ pmn}^i a_j^p - A_{15\ jmn}^p a_p^i - P_{nm}^{*p} (a_{j|_1p}^i - a_{j|_2p}^i), \quad \text{i.e.} \\
a_{j|_1\ 2}^i - a_{j|_2\ 1}^i &= R_{3\ pmn}^{*i} a_j^p - R_{3\ jmn}^{*p} a_p^i,
\end{aligned}$$

where

$$\begin{aligned}
a_{j<mn>}^i &= T_{pm}^{*i} \delta_n a_j^p - T_{jm}^{*p} \delta_n a_p^i, \\
a_{j\leq mn\geq}^i &= (P_{np}^{*i} P_{jn}^{*s} - P_{pm}^{*i} P_{nj}^{*s}) a_s^p, \\
a_{j\leq mn\geq}^i &= (P_{np}^{*i} P_{nj}^{*s} - P_{pm}^{*i} P_{jn}^{*s}) a_s^p, \\
a_{j\leq mn\geq}^i &= (T_{np}^{*i} P_{jn}^{*s} - P_{pn}^{*i} T_{mj}^{*s}) a_s^p, \\
a_{j\leq mn\geq}^i &= (P_{np}^{*i} T_{jn}^{*s} - T_{pn}^{*i} P_{mj}^{*s}) a_s^p,
\end{aligned}$$

the magnitudes  $R_{t\ jmn}^{*i}$ ,  $t = 1, 2, 3$ , are curvature tensors and the magnitudes  $A_{t\ jmn}^{*i}$ ,  $t = 1, \dots, 15$ , are not tensors (we will verify this fact in what follows), we call them curvature pseudotensors of the first, the second, ..., and the fifteenth kind, respectively.

Using the third and the fourth kind of the covariant derivative given in (2.1), one gets ten new Ricci type identities (see, [7]). In these identities appear the same quantities  $R_{t\ jmn}^{*i}$ ,  $t = 1, 2, 3$ ,  $A_{s\ jmn}^{*i}$ ,  $s = 1, \dots, 15$ , but in different distribution. Only in the last case appears a new curvature tensor, the curvature tensor of the fourth kind  $R_{4\ jmn}^{*i}$  [7]:

$$a_{j|_3\ 4}^i - a_{j|_4\ 3}^i = R_{4\ pmn}^{*i} a_j^p + R_{3\ jmn}^{*p} a_p^i.$$

The curvature tensors  $R^*{}^i{}_{jmn}$ ,  $t = 1, \dots, 4$ , are given by [7]–[9]

$$(2.2) \quad \begin{aligned} R^*{}^i{}_{1jmn} &= \delta_n P^*{}^i{}_{jm} - \delta_m P^*{}^i{}_{jn} + P^*{}^{pp} P^*{}^i{}_{pn} - P^*{}^{pp} P^*{}^i{}_{pm} \\ &= (\delta_n P^*{}^i{}_{jm} + P^*{}^{pp} P^*{}^i{}_{pn})_{(mn)}, \end{aligned}$$

$$(2.3) \quad R^*{}^i{}_{2jmn} = (\delta_n P^*{}^i{}_{mj} + P^*{}^{pp} P^*{}^i{}_{np})_{(mn)},$$

$$(2.4) \quad R^*{}^i{}_{3jmn} = \delta_n P^*{}^i{}_{jm} - \delta_m P^*{}^i{}_{nj} + P^*{}^{pp} P^*{}^i{}_{np} - P^*{}^{pp} P^*{}^i{}_{pm} + P^*{}^{pp} P^*{}^i{}_{[pj]},$$

$$(2.5) \quad R^*{}^i{}_{4jmn} = \delta_n P^*{}^i{}_{jm} - \delta_m P^*{}^i{}_{nj} + P^*{}^{pp} P^*{}^i{}_{np} - P^*{}^{pp} P^*{}^i{}_{pm} + P^*{}^{pp} P^*{}^i{}_{[pj]}.$$

In [15] are given some properties of the mentioned curvature tensors (the antisymmetry with respect to two indices, the cyclic symmetry, the symmetry with respect to pairs of indices).

The curvature pseudotensors  $A^*{}^i{}_{tjmn}$ ,  $t = 1, \dots, 15$ , are given by [8]

$$(2.6) \quad A^*{}^i{}_{1jmn} = (\delta_n P^*{}^i{}_{jm} + P^*{}^{pp} P^*{}^i{}_{np})_{(mn)},$$

$$(2.7) \quad A^*{}^i{}_{2jmn} = (\delta_n P^*{}^i{}_{jm} + P^*{}^{pp} P^*{}^i{}_{pn})_{(mn)},$$

$$(2.8) \quad A^*{}^i{}_{3jmn} = (\delta_n P^*{}^i{}_{mj} + P^*{}^{pp} P^*{}^i{}_{pn})_{(mn)},$$

$$(2.9) \quad A^*{}^i{}_{4jmn} = (\delta_n P^*{}^i{}_{mj} + P^*{}^{pp} P^*{}^i{}_{np})_{(mn)},$$

$$(2.10) \quad A^*{}^i{}_{5jmn} = \delta_n P^*{}^i{}_{jm} - \delta_m P^*{}^i{}_{nj} + P^*{}^{pp} P^*{}^i{}_{pn} - P^*{}^{pp} P^*{}^i{}_{mp},$$

$$(2.11) \quad A^*{}^i{}_{6jmn} = \delta_n P^*{}^i{}_{jm} - \delta_m P^*{}^i{}_{nj} + P^*{}^{pp} P^*{}^i{}_{np} - P^*{}^{pp} P^*{}^i{}_{pm},$$

$$(2.12) \quad A^*{}^i{}_{7jmn} = (\delta_n P^*{}^i{}_{jm})_{(mn)} + P^*{}^{pp} P^*{}^i{}_{pn} - P^*{}^{pp} P^*{}^i{}_{mp},$$

$$(2.13) \quad A^*{}^i{}_{8jmn} = (\delta_n P^*{}^i{}_{jm})_{(mn)} + P^*{}^{pp} P^*{}^i{}_{pn} - P^*{}^{pp} P^*{}^i{}_{pm},$$

$$(2.14) \quad A^*{}^i{}_{9jmn} = \delta_n P^*{}^i{}_{jm} - \delta_m P^*{}^i{}_{nj} + P^*{}^{pp} P^*{}^i{}_{pn} - P^*{}^{pp} P^*{}^i{}_{pm},$$

$$(2.15) \quad A^*{}^i{}_{10jmn} = \delta_n P^*{}^i{}_{jm} - \delta_m P^*{}^i{}_{nj} + P^*{}^{pp} P^*{}^i{}_{np} - P^*{}^{pp} P^*{}^i{}_{pm},$$

$$(2.16) \quad A^*{}^i{}_{11jmn} = \delta_n P^*{}^i{}_{mj} - \delta_m P^*{}^i{}_{jn} + P^*{}^{pp} P^*{}^i{}_{np} - P^*{}^{pp} P^*{}^i{}_{mp},$$

$$(2.17) \quad A^*{}^i{}_{12jmn} = \delta_n P^*{}^i{}_{mj} - \delta_m P^*{}^i{}_{jn} + P^*{}^{pp} P^*{}^i{}_{pn} - P^*{}^{pp} P^*{}^i{}_{mp},$$

$$(2.18) \quad A^*{}^i{}_{13jmn} = \delta_n P^*{}^i{}_{mj} - \delta_m P^*{}^i{}_{nj} + P^*{}^{pp} P^*{}^i{}_{np} - P^*{}^{pp} P^*{}^i{}_{pm},$$

$$(2.19) \quad A^*{}^i{}_{14jmn} = \delta_n P^*{}^i{}_{mj} - \delta_m P^*{}^i{}_{nj} + P^*{}^{pp} P^*{}^i{}_{np} - P^*{}^{pp} P^*{}^i{}_{mp},$$

$$(2.20) \quad A^*{}^i{}_{15jmn} = \delta_n P^*{}^i{}_{jm} - \delta_m P^*{}^i{}_{nj} + P^*{}^{pp} P^*{}^i{}_{np} - P^*{}^{pp} P^*{}^i{}_{pm}.$$

The signature  $( )_{(mn)}$  denotes an antisymmetrization without division with respect to  $m, n$ , of the expression in the bracket. Analogously, the signature  $( )_{[mn]}$  signifies a symmetrization without division with respect to  $m, n$ .

We should mention that Ricci type identities, curvature tensors and pseudotensors, given in [7, 8] are analogous to Ricci type identities, curvature tensors and pseudotensors in the space of non-symmetric affine connection, given in [4, 6].

The non-symmetric connection  $P_{jk}^{*i}$  can be represented as follows:

$$(2.21) \quad P_{jk}^{*i} = \frac{1}{2}P_{(jk)}^{*i} + \frac{1}{2}T_{jk}^{*i}.$$

Let us denote by  $R_{jmn}^{*i}$  the curvature tensor of symmetric connection  $\frac{1}{2}P_{(jk)}^{*i}$  i.e.

$$(2.22) \quad R_{jmn}^{*i}(x, \dot{x}) = \frac{1}{2}(\delta_n P_{(jm)}^{*i} + \frac{1}{2}P_{(jm)}^{*p} P_{(pn)}^{*i})_{[mnn]}.$$

According to (2.21) the curvature tensor of the first kind can be represented in the following manner:

$$(2.23) \quad \begin{aligned} R_{1jmn}^{*i} &= \frac{1}{2} \left\{ \delta_n P_{jm}^{*i} + \delta_n T_{jm}^{*i} + \frac{1}{2} (P_{(jm)}^{*p} P_{(pn)}^{*i} + T_{jm}^{*p} P_{(pn)}^{*i} \right. \\ &\quad \left. + P_{(jm)}^{*p} T_{pn}^{*i} + T_{jm}^{*p} T_{pn}^{*i}) \right\}_{[mnn]}. \end{aligned}$$

If one denotes by semicolon (;) the covariant derivative with respect to symmetric connection  $\frac{1}{2}P_{(jk)}^{*i}$ , equation (2.23) becomes

$$(2.24) \quad R_{1jmn}^{*i} = R_{jmn}^{*i} + \frac{1}{2} (T_{jm;n}^{*i} + \frac{1}{2} T_{jm}^{*p} T_{pn}^{*i})_{[mnn]},$$

that is, we obtained the relation between  $R_{1jmn}^{*i}$ ,  $R_{jmn}^{*i}$  and the torsion tensor  $T_{jk}^{*i}$ .

Following this procedure for the curvature tensors  $R_{tjmn}^{*i}$ ,  $t = 2, \dots, 4$ , in  $GF_N$ , we get:

$$(2.25) \quad R_{2jmn}^{*i} = R_{jmn}^{*i} + \frac{1}{2} (T_{mj;n}^{*i} + \frac{1}{2} T_{mj}^{*p} T_{np}^{*i})_{[mnn]},$$

$$(2.26) \quad R_{3jmn}^{*i} = R_{jmn}^{*i} + \frac{1}{2} ((T_{jm;n}^{*i})_{(nm)} + \frac{1}{2} (T_{jm}^{*p} T_{np}^{*i})_{[mnn]} + T_{mn}^p T_{jp}^i),$$

$$(2.27) \quad R_{4jmn}^{*i} = R_{jmn}^{*i} + \frac{1}{2} ((T_{jm;n}^{*i})_{(nm)} + \frac{1}{2} (T_{jm}^{*p} T_{np}^{*i})_{[mnn]} - T_{mn}^{*p} T_{jp}^{*i}).$$

Based on the discussion given above, we can formulate the next theorem.

**Theorem 2.1.** *The curvature tensors  $R_{\theta jmn}^{*i}$  ( $\theta = 1, \dots, 4$ ), corresponding to the non-symmetric connection  $P_{jk}^{*i}$  given by (2.2)–(2.5), and the curvature tensor  $R_{jmn}^{*i}$ , corresponding to the symmetric connection  $\frac{1}{2}P_{(jk)}^{*i} = \frac{1}{2}(P_{jk}^{*i} + P_{kj}^{*i})$ , given by (2.22), satisfy relations (2.24)–(2.27), where  $T_{jk}^{*i}$  is the torsion tensor of connection  $P_{jk}^{*i}$ .*

In a similar manner, we can prove the following theorem.

**Theorem 2.2.** *The curvature pseudotensors  $A_{t\ jmn}^{*i}$ ,  $t = 1, \dots, 15$ , given by equations (2.6)–(2.20) and the curvature tensor  $R_{jmn}^{*i}$ , given by Eq. (2.22), satisfy the following relations*

$$\begin{aligned} A_{1\ jmn}^{*i} &= R_{jmn}^{*i} + \left( \frac{1}{2} \delta_n T_{jm}^{*i} + \frac{1}{4} T_{jm}^{*p} P_{(np)}^{*i} + \frac{1}{4} T_{jm}^{*p} T_{np}^{*i} + \frac{1}{4} P_{(jm)}^{*p} T_{np}^{*i} \right)_{[mn]}, \\ A_{2\ jmn}^{*i} &= R_{jmn}^{*i} + \left( \frac{1}{2} \delta_n T_{jm}^{*i} + \frac{1}{4} T_{mj}^{*p} P_{(pn)}^{*i} + \frac{1}{4} T_{mj}^{*p} T_{pn}^{*i} + \frac{1}{4} P_{(mj)}^{*p} T_{pn}^{*i} \right)_{[mn]}, \\ A_{3\ jmn}^{*i} &= R_{jmn}^{*i} + \left( \frac{1}{2} \delta_n T_{mj}^{*i} + \frac{1}{4} T_{mj}^{*p} P_{(pn)}^{*i} + \frac{1}{4} T_{mj}^{*p} T_{pn}^{*i} + \frac{1}{4} P_{(mj)}^{*p} T_{pn}^{*i} \right)_{[mn]}, \\ A_{4\ jmn}^{*i} &= R_{jmn}^{*i} + \left( \frac{1}{2} \delta_n T_{mj}^{*i} + \frac{1}{4} T_{jm}^{*p} P_{(np)}^{*i} + \frac{1}{4} T_{jm}^{*p} T_{np}^{*i} + \frac{1}{4} P_{(jm)}^{*p} T_{np}^{*i} \right)_{[mn]}, \end{aligned}$$

$$\begin{aligned} A_{5\ jmn}^{*i} &= R_{jmn}^{*i} + \frac{1}{2} \delta_n T_{jm}^{*i} - \frac{1}{2} \delta_m T_{nj}^{*i} + \frac{1}{4} P_{(jm)}^{*p} T_{pn}^{*i} + \frac{1}{4} T_{jm}^{*p} P_{(pn)}^{*i} \\ &\quad + \frac{1}{4} T_{jm}^{*p} T_{pn}^{*i} - \frac{1}{4} P_{(nj)}^{*p} T_{mp}^{*i} - \frac{1}{4} T_{nj}^{*p} P_{(mp)}^{*i} - \frac{1}{4} T_{nj}^{*p} T_{mp}^{*i}, \\ A_{6\ jmn}^{*i} &= R_{jmn}^{*i} + \frac{1}{2} \delta_n T_{jm}^{*i} - \frac{1}{2} \delta_m T_{nj}^{*i} + \frac{1}{4} P_{(mj)}^{*p} T_{np}^{*i} + \frac{1}{4} T_{mj}^{*p} P_{(np)}^{*i} \\ &\quad + \frac{1}{4} T_{mj}^{*p} T_{np}^{*i} - \frac{1}{4} P_{(jn)}^{*p} T_{pm}^{*i} - \frac{1}{4} T_{jn}^{*p} P_{(pm)}^{*i} - \frac{1}{4} T_{jn}^{*p} T_{pm}^{*i}, \end{aligned}$$

$$\begin{aligned}
A_{7\ jmn}^{*i} &= R_{jmn}^{*i} + \frac{1}{2}\delta_n T_{jm}^{*i} - \frac{1}{2}\delta_m T_{jn}^{*i} + \frac{1}{4}P_{(jm)}^{*p} T_{pn}^{*i} + \frac{1}{4}T_{jm}^{*p} P_{(pn)}^{*i} \\
&\quad + \frac{1}{4}T_{jm}^{*p} T_{pn}^{*i} - \frac{1}{4}P_{(jn)}^{*p} T_{mp}^{*i} - \frac{1}{4}T_{jn}^{*p} P_{(mp)}^{*i} - \frac{1}{4}T_{jn}^{*p} T_{mp}^{*i}, \\
A_{8\ jmn}^{*i} &= R_{jmn}^{*i} + \frac{1}{2}\delta_n T_{jm}^{*i} - \frac{1}{2}\delta_m T_{jn}^{*i} + \frac{1}{4}P_{(mj)}^{*p} T_{pn}^{*i} + \frac{1}{4}T_{mj}^{*p} P_{(pn)}^{*i} \\
&\quad + \frac{1}{4}T_{mj}^{*p} T_{pn}^{*i} - \frac{1}{4}P_{(jn)}^{*p} T_{pm}^{*i} - \frac{1}{4}T_{jn}^{*p} P_{(pm)}^{*i} - \frac{1}{4}T_{jn}^{*p} T_{pm}^{*i}, \\
A_{9\ jmn}^{*i} &= R_{jmn}^{*i} + \frac{1}{2}\delta_n T_{jm}^{*i} - \frac{1}{2}\delta_m T_{nj}^{*i} + \frac{1}{4}P_{(jm)}^{*p} T_{pn}^{*i} + \frac{1}{4}T_{jm}^{*p} P_{(pn)}^{*i} \\
&\quad + \frac{1}{4}T_{jm}^{*p} T_{pn}^{*i} - \frac{1}{4}P_{(nj)}^{*p} T_{pm}^{*i} - \frac{1}{4}T_{nj}^{*p} P_{(pm)}^{*i} - \frac{1}{4}T_{nj}^{*p} T_{pm}^{*i}, \\
A_{10\ jmn}^{*i} &= R_{jmn}^{*i} + \frac{1}{2}\delta_n T_{jm}^{*i} - \frac{1}{2}\delta_m T_{nj}^{*i} + \frac{1}{4}P_{(jm)}^{*p} T_{np}^{*i} + \frac{1}{4}T_{jm}^{*p} P_{(np)}^{*i} \\
&\quad + \frac{1}{4}T_{jm}^{*p} T_{np}^{*i} - \frac{1}{4}P_{(jn)}^{*p} T_{pm}^{*i} - \frac{1}{4}T_{jn}^{*p} P_{(pm)}^{*i} - \frac{1}{4}T_{jn}^{*p} T_{pm}^{*i}, \\
A_{11\ jmn}^{*i} &= R_{jmn}^{*i} + \frac{1}{2}\delta_n T_{mj}^{*i} - \frac{1}{2}\delta_m T_{jn}^{*i} + \frac{1}{4}P_{(mj)}^{*p} T_{np}^{*i} + \frac{1}{4}T_{mj}^{*p} P_{(np)}^{*i} \\
&\quad + \frac{1}{4}T_{mj}^{*p} T_{np}^{*i} - \frac{1}{4}P_{(jn)}^{*p} T_{mp}^{*i} - \frac{1}{4}T_{jn}^{*p} P_{(mp)}^{*i} - \frac{1}{4}T_{jn}^{*p} T_{mp}^{*i}, \\
A_{12\ jmn}^{*i} &= R_{jmn}^{*i} + \frac{1}{2}\delta_n T_{mj}^{*i} - \frac{1}{2}\delta_m T_{jn}^{*i} + \frac{1}{4}P_{(mj)}^{*p} T_{pn}^{*i} + \frac{1}{4}T_{mj}^{*p} P_{(pn)}^{*i} \\
&\quad + \frac{1}{4}T_{mj}^{*p} T_{pn}^{*i} - \frac{1}{4}P_{(nj)}^{*p} T_{mp}^{*i} - \frac{1}{4}T_{nj}^{*p} P_{(mp)}^{*i} - \frac{1}{4}T_{nj}^{*p} T_{mp}^{*i}, \\
A_{13\ jmn}^{*i} &= R_{jmn}^{*i} + \frac{1}{2}\delta_n T_{mj}^{*i} - \frac{1}{2}\delta_m T_{nj}^{*i} + \frac{1}{4}P_{(mj)}^{*p} T_{np}^{*i} + \frac{1}{4}T_{mj}^{*p} P_{(np)}^{*i} \\
&\quad + \frac{1}{4}T_{mj}^{*p} T_{np}^{*i} - \frac{1}{4}P_{(nj)}^{*p} T_{pm}^{*i} - \frac{1}{4}T_{nj}^{*p} P_{(pm)}^{*i} - \frac{1}{4}T_{nj}^{*p} T_{pm}^{*i}, \\
A_{14\ jmn}^{*i} &= R_{jmn}^{*i} + \frac{1}{2}\delta_n T_{mj}^{*i} - \frac{1}{2}\delta_m T_{nj}^{*i} + \frac{1}{4}P_{(jm)}^{*p} T_{np}^{*i} + \frac{1}{4}T_{jm}^{*p} P_{(np)}^{*i} \\
&\quad + \frac{1}{4}T_{jm}^{*p} T_{np}^{*i} - \frac{1}{4}P_{(nj)}^{*p} T_{mp}^{*i} - \frac{1}{4}T_{nj}^{*p} P_{(mp)}^{*i} - \frac{1}{4}T_{nj}^{*p} T_{mp}^{*i}, \\
A_{15\ jmn}^{*i} &= R_{jmn}^{*i} + \frac{1}{2}\delta_n T_{jm}^{*i} - \frac{1}{2}\delta_m T_{nj}^{*i} + \frac{1}{4}P_{(jm)}^{*p} T_{np}^{*i} + \frac{1}{4}T_{jm}^{*p} P_{(np)}^{*i} \\
&\quad + \frac{1}{4}T_{jm}^{*p} T_{np}^{*i} - \frac{1}{4}P_{(nj)}^{*p} T_{pm}^{*i} - \frac{1}{4}T_{nj}^{*p} P_{(pm)}^{*i} - \frac{1}{4}T_{nj}^{*p} T_{pm}^{*i},
\end{aligned}$$

where  $T_{jk}^{*i}$  is the torsion tensor of connection  $P_{jk}^{*i}$ .

From the obtained expressions of magnitudes  $A_{t\ jmn}^{*i}$ ,  $t = 1, \dots, 15$ , given in Theorem 2.2, we see that these magnitudes are not tensors because  $T_{jk}^{*i}$  is a torsion tensor and  $P_{(jk)}^{*i}$  is not a tensor. Also, we can obtain the transformation laws for curvature pseudotensors. For example, for  $A_{7\ jmn}^{*i}$ , we see that the last addend is not



a tensor, in coordinates  $x^{i'}$  it is

$$(2.28) \quad P^{*p'}_{(j'n')} T^{*i'}_{m'p'} = (P^{*p}_{(jn)} x^{p'}_j x^j_{n'} + x^p_{j'n'} x^{p'}_p) T^{*i} x^i_{m'} x^m_{p'} x^q_{p'} = P^{*p}_{(jn)} T^{*i}_{mp} x^i_{j'} x^j_{m'} x^n_{n'} + x^p_{j'n'} T^{*i}_{mp} x^i_{j'} x^m_{m'}$$

because

$$T^{*i}_{mq} x^p_{j'} x^q_{p'} = T^{*i}_{mq} \delta^q_p = T^{*i}_{mp}$$

Now, from (2.28) we obtain

$$A^{*i'}_{7j'n'n'} = A^{*i}_{7jmn} x^i_{j'} x^j_{m'} x^n_{n'} - 2x^p_{j'n'} T^{*i}_{mp} x^i_{j'} x^m_{m'}$$

**Remark 2.1.** If  $g_{ij} = 0$  i.e.  $g_{ij}(x, \dot{x}) = g_{ij}(x, \dot{x})$ , then the generalized Finsler space reduces to the usual Finsler space. In this case the curvature tensors  $R^{*i}_{1jmn}, \dots, R^{*i}_{4jmn}$  and the curvature pseudotensors  $A^{*i}_{1jmn}, \dots, A^{*i}_{15jmn}$  reduce to the curvature tensor  $R^{*i}_{jmn}$  of usual Finsler space.

### 3. Derived curvature tensors of non-symmetric connection in Rund's sense

In the paper [9] we examined combined Ricci type identities and derived eight new curvature tensors  $\widetilde{R}^{*i}_{tjmn}$ ,  $t = 1, \dots, 8$ , in the generalized Finsler space, it is analogous to related results in the non-symmetric affine connection space [5]. Derived curvature tensors  $\widetilde{R}^{*i}_{tjmn}$ ,  $t = 1, \dots, 8$ , are given by

$$(3.1) \quad \widetilde{R}^{*i}_{1jmn} = \frac{1}{2}(A^{*i}_{1jmn} + A^{*i}_{3jmn}) = \frac{1}{2}(A^{*i}_{2jmn} + A^{*i}_{4jmn}) = \frac{1}{2}A^{*i}_{j[lmn]}$$

$$(3.2) \quad \widetilde{R}^{*i}_{2jmn} = \frac{1}{2}(A^{*i}_{7jmn} + A^{*i}_{13jmn}) = \frac{1}{2}(A^{*i}_{9jmn} + A^{*i}_{11jmn}),$$

$$(3.3) \quad \widetilde{R}^{*i}_{3jmn} = \frac{1}{2}(A^{*i}_{8jmn} + A^{*i}_{14jmn}) = \frac{1}{2}(A^{*i}_{10jmn} + A^{*i}_{12jmn}),$$

$$(3.4) \quad \begin{aligned} \widetilde{R}^{*i}_{4jmn} &= \frac{1}{6}(R^*_{3jmn} + A^*_{11jmn} + A^*_{13jmn}) - A^{*i}_{13jmn} \\ &= \frac{1}{6}(R^*_{3jmn} + A^*_{12jmn} + A^*_{14jmn}) - A^{*i}_{13jmn} \end{aligned}$$

$$(3.5) \quad \widetilde{R}^{*i}_{5jmn} = (A^*_{1jmn} - A^*_{7jmn}) - A^{*i}_{13jmn} = -(A^*_{11jmn} + A^*_{15jmn}) - A^{*i}_{7jmn}$$

$$(3.6) \quad \widetilde{R}^{*i}_{6jmn} = (A^*_{2jmn} - A^*_{8jmn}) - A^{*i}_{14jmn} = -(A^*_{12jmn} + A^*_{15jmn}) - A^{*i}_{8jmn}$$

$$(3.7) \quad \widetilde{R}^{*i}_{7jmn} = (A^*_{3jmn} + A^*_{7jmn}) + A^{*i}_{13jmn} = (A^*_{13jmn} - A^*_{15jmn}) + A^{*i}_{9jmn}$$

$$(3.8) \quad \widetilde{R}^{*i}_{8jmn} = (A^*_{4jmn} + A^*_{8jmn}) + A^{*i}_{14jmn} = (A^*_{14jmn} - A^*_{15jmn}) + A^{*i}_{10jmn}$$

where  $R_3^*$  is the third curvature tensor and  $A_t^*$ ,  $t = 1, \dots, 15$ , are the curvature pseudotensors of non-symmetric connection in Rund's sense.

As an easy corollary of Theorem 2.1 and Theorem 2.2 we obtain relations between derived curvature tensors  $\widetilde{R}_t^{*i}$ ,  $t = 1, \dots, 8$ , and the curvature tensor  $R_{jmn}^{*i}$  of associated symmetric connection.

**Corollary 3.1.** *Derived curvature tensors  $\widetilde{R}_\theta^*$  ( $\theta = 1, \dots, 8$ ), corresponding to the non-symmetric connection  $P^*$ , given by (3.1)–(3.8) and the curvature tensor  $R_{jmn}^{*i}$ , corresponding to the symmetric connection  $\frac{1}{2}P_{(jk)}^{*i} = \frac{1}{2}(P_{jk}^{*i} + P_{kj}^{*i})$ , satisfy the following relations*

$$\begin{aligned}\widetilde{R}_1^{*i}{}_{jmn} &= R_{jmn}^{*i} - \frac{1}{4}T_{jm}^{*p}T_{pn}^{*i} + \frac{1}{4}T_{jn}^{*p}T_{pm}^{*i}, \\ \widetilde{R}_2^{*i}{}_{jmn} &= R_{jmn}^{*i} + \frac{1}{4}T_{jm}^{*p}T_{pn}^{*i} + \frac{1}{4}T_{jn}^{*p}T_{pm}^{*i}, \\ \widetilde{R}_3^{*i}{}_{jmn} &= R_{jmn}^{*i} - \frac{1}{4}T_{jm}^{*p}T_{pn}^{*i} - \frac{1}{4}T_{jn}^{*p}T_{pm}^{*i}, \\ \widetilde{R}_4^{*i}{}_{jmn} &= R_{jmn}^{*i} - \frac{1}{6}\left((T_{jm;n}^{*i} + \frac{1}{4}T_{jm}^{*p}T_{pn}^{*i})_{[mn]} + T_{mn}^{*p}T_{pj}^{*i}\right), \\ \widetilde{R}_5^{*i}{}_{jmn} &= R_{jmn}^{*i} - (T_{jm;n}^{*i})_{[mn]} - \frac{3}{4}T_{jm}^{*p}T_{pn}^{*i} - \frac{1}{4}T_{jn}^{*p}T_{pm}^{*i}, \\ \widetilde{R}_6^{*i}{}_{jmn} &= R_{jmn}^{*i} - (T_{jm;n}^{*i})_{[mn]} + \frac{1}{4}T_{jm}^{*p}T_{pn}^{*i} + \frac{3}{4}T_{jn}^{*p}T_{pm}^{*i}, \\ \widetilde{R}_7^{*i}{}_{jmn} &= R_{jmn}^{*i} + (T_{jm;n}^{*i})_{[mn]} + \frac{1}{4}T_{jm}^{*p}T_{pn}^{*i} + \frac{3}{4}T_{jn}^{*p}T_{pm}^{*i}, \\ \widetilde{R}_8^{*i}{}_{jmn} &= R_{jmn}^{*i} + (T_{jm;n}^{*i})_{[mn]} - \frac{3}{4}T_{jm}^{*p}T_{pn}^{*i} - \frac{1}{4}T_{jn}^{*p}T_{pm}^{*i}\end{aligned}$$

where  $T_{jk}^{*i}$  is the torsion tensor of connection  $P_{jk}^{*i}$ .

**Remark 3.1.** *If  $T_{jk}^{*i} = 0$ , then derived curvature tensors  $\widetilde{R}_t^{*i}$ ,  $t = 1, \dots, 8$ , reduce to the curvature tensor  $R_{jmn}^{*i}$ . This fact evidently follows from Corollary 3.1.*

#### 4. Concluding remarks

The presented results are generalizations of related results in the generalized (non-symmetric) Riemannian spaces.

If  $g_{ij}(x, \dot{x}) = g_{ij}(x) \neq g_{ji}(x)$ , in general,  $GF_N$  reduces to  $GR_N$  (generalized Riemannian space), if  $g_{ij}(x, \dot{x}) = g_{ij}(x) = g_{ji}(x)$  we obtain the usual Riemannian space  $R_N$ .

## Acknowledgement

The authors are grateful to Professor Irena Čomić for her valuable remarks and suggestions.

## REFERENCES

1. E. CARTAN: *Les Espaces de Finsler*. Paris, 1934.
2. M.D. CVETKOVIĆ, M.LJ. ZLATANOVIĆ: *New Cartan's tensors and pseudotensors in a Generalized Finsler Space*. *Filomat* **28**(2014), 107–117.
3. L.P. EISENHART: *Generalized Riemannian Spaces*. *Proc. Nat. Acad. Sci. USA* **37**(1951), 311–315.
4. S.M. MINIČIĆ: *Ricci identities in the space of non-symmetric affine connexion*. *Mat. Vesnik* **25**(1973), 161–172
5. S.M. MINIČIĆ: *Curvature tensors of the space of non-symmetric affine connexion, obtained from the curvature pseudotensors*. *Mat. Vesnik* **28**(1976), 421–435.
6. S.M. MINIČIĆ: *New commutation formulas in the non-symmetric affine connexion space*. *Publ. Inst. Math. (Beograd) (N.S)* **22**(1977), 189–199.
7. S.M. MINIČIĆ, M.LJ. ZLATANOVIĆ: *New commutation formulas for  $\delta$ -differentiation in a generalized Finsler space*. *Differential Geometry-Dynamical Systems* **12**(2010), 145–157.
8. S.M. MINIČIĆ, M.LJ. ZLATANOVIĆ: *Commutation formulas for  $\delta$ -differentiation in a generalized Finsler space*. *Kragujevac Journal of Mathematics*, **35**(2011), 277–289.
9. S.M. MINIČIĆ, M.LJ. ZLATANOVIĆ: *Derived curvature tensors in generalized Finsler space*. *Differential Geometry-Dynamical Systems* **13**(2011), 179–190.
10. H. RUND: *The Differential Geometry of Finsler Space*. Springer-Verlag, Berlin-Göttingen-Heidelberg, 1959.
11. A.C. SHAMIHOKE: *Some properties of a curvature tensors in a generalised Finsler space*. *Tensor N.S.* **12**(1962), 97–109.
12. A.C. SHAMIHOKE: *Hypersurfaces of a generalised Finsler space*. *Tensor N.S.* **13** (1963), 129–144.
13. A.C. SHAMIHOKE: *A note on a curvature tensor in a generalized Finsler space*. *Tensor N.S.* **15**(1964), 20–22.
14. A.C. SHAMIHOKE: *Parallelism and covariant differentiation in a generalized Finsler space of  $n$  dimensions*. *Riv. Mat. Univ. Parma II. Ser.* **5**(1964), 189–200.
15. M.LJ. ZLATANOVIĆ, S.M. MINIČIĆ: *Identities for curvature tensors in generalized Finsler space*. *Filomat* **23**(2009), 34–42.

Milan Lj. Zlatanović  
Faculty of Sciences and Mathematics  
Department of Mathematics

Višegradska 33, 18000 Niš, Serbia  
zlatmilan@yahoo.com

Svetislav M. Minčić  
Faculty of Sciences and Mathematics  
Department of Mathematics  
Višegradska 33, 18000 Niš, Serbia  
svetislavmincic@yahoo.com

Miloš Z. Petrović  
Faculty of Sciences and Mathematics  
Department of Mathematics  
Višegradska 33, 18000 Niš, Serbia  
somipetrovic89@gmail.com