

FACTA UNIVERSITATIS (NIS)  
SER. MATH. INFORM. Vol. 31, No 3 (2016), 595–608

## PERTURBED COMPANION OF OSTROWSKI TYPE INEQUALITY FOR TWICE DIFFERENTIABLE FUNCTIONS

Mehmet Zeki Sarıkaya, Hüseyin Budak, Tuba Tunç, Samet Erden  
and Hatice Yıldız

**Abstract.** In this paper, some perturbed companion of Ostrowski type integral inequalities for functions whose second derivatives are either bounded or of bounded variation are established.

**Keywords:** Perturbed Ostrowski type inequality, function of bounded variation.

### 1. Introduction

In 1938, Ostrowski [29] established a following useful inequality:

**Theorem 1.1.** *Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  whose derivative  $f' : (a, b) \rightarrow \mathbb{R}$  is bounded on  $(a, b)$ , i.e.  $\|f'\|_\infty := \sup_{t \in (a, b)} |f'(t)| < \infty$ .*

*Then, we have the inequality*

$$(1.1) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[ \frac{1}{4} + \frac{\left(x - \frac{a+b}{2}\right)^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty,$$

for all  $x \in [a, b]$ .

The constant  $\frac{1}{4}$  is the best possible.

**Definition 1.1.** Let  $P : a = x_0 < x_1 < \dots < x_n = b$  be any partition of  $[a, b]$  and let  $\Delta f(x_i) = f(x_{i+1}) - f(x_i)$ , then  $f$  is said to be of bounded variation if the sum

$$\sum_{i=1}^m |\Delta f(x_i)|$$

is bounded for all such partitions.

Received April 11, 2016; accepted June 18, 2016

2010 *Mathematics Subject Classification.* Primary 26D15; Secondary 26A45, 26D10.

**Definition 1.2.** Let  $f$  be of bounded variation on  $[a, b]$ , and  $\sum \Delta f(P)$  denotes the sum  $\sum_{i=1}^n |\Delta f(x_i)|$  corresponding to the partition  $P$  of  $[a, b]$ . The number

$$\bigvee_a^b(f) := \sup \left\{ \sum \Delta f(P) : P \in P([a, b]) \right\},$$

is called the total variation of  $f$  on  $[a, b]$ . Here  $P([a, b])$  denotes the family of partitions of  $[a, b]$ .

In [16], Dragomir proved the following Ostrowski type inequalities for functions of bounded variation:

**Theorem 1.2.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be a mapping of bounded variation on  $[a, b]$ . Then

$$(1.2) \quad \left| \int_a^b f(t) dt - (b-a)f(x) \right| \leq \left[ \frac{1}{2}(b-a) + \left| x - \frac{a+b}{2} \right| \right] \bigvee_a^b(f)$$

holds for all  $x \in [a, b]$ . The constant  $\frac{1}{2}$  is the best possible.

In [13], the authors obtained the following Ostrowski type inequalities for functions whose second derivatives are bounded:

**Theorem 1.3.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be continuous on  $[a, b]$  and twice differentiable on  $(a, b)$ , whose second derivative  $f'' : (a, b) \rightarrow \mathbb{R}$  is bounded on  $(a, b)$ . Then we have the inequality

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt - \frac{f(b) - f(a)}{b-a} \left( x - \frac{a+b}{2} \right) \right| \\ & \leq \frac{1}{2} \left\{ \left[ \frac{\left( x - \frac{a+b}{2} \right)^2}{(b-a)^2} + \frac{1}{4} \right]^2 + \frac{1}{12} \right\} (b-a)^2 \|f''\|_\infty \\ & \leq \frac{\|f''\|_\infty}{6} (b-a)^2 \end{aligned}$$

for all  $x \in [a, b]$ .

Ostrowski inequality has potential applications in mathematical sciences. In the past, many authors have worked on Ostrowski type inequalities for functions (bounded, of bounded variation, etc.) see for example ([1]-[10], [13]-[19], [27],[28],[30]-[37]). Furthermore, several works were devoted to study of perturbed Ostrowski type inequalities for bounded functions and functions of bounded variation, please refer to ([11],[12], [20]-[26]). In this study, we establish some perturbed companion of Ostrowski type inequalities for twice differentiable functions whose second derivatives are either bounded or of bounded variation.

2. Some Identities

Before we start our main results, we state and prove the following lemma:

**Lemma 2.1.** *Let  $f : [a, b] \rightarrow \mathbb{C}$  be a twice differentiable function on  $(a, b)$ . Then for any  $\lambda_i(x)$ ,  $i = 1, 2, 3$  complex number and all  $x \in [a, \frac{a+b}{2}]$  the following identity holds*

$$\begin{aligned}
 (2.1) \quad & \left(x - \frac{3a+b}{4}\right) \frac{f'(x) - f'(a+b-x)}{2} - \frac{f(x) + f(a+b-x)}{2} + \frac{1}{b-a} \int_a^b f(t) dt \\
 & - \frac{1}{6(b-a)} \left[ (x-a)^3 (\lambda_1(x) + \lambda_3(x)) + 2 \left(\frac{a+b}{2} - x\right)^3 \lambda_2(x) \right] \\
 = & \frac{1}{2(b-a)} \left[ \int_a^x (t-a)^2 [f''(t) - \lambda_1(x)] dt + \int_x^{a+b-x} \left(t - \frac{a+b}{2}\right)^2 [f''(t) - \lambda_2(x)] dt \right. \\
 & \left. + \int_{a+b-x}^b (t-b)^2 [f''(t) - \lambda_3(x)] dt \right]
 \end{aligned}$$

where the integrals in the right hand side are taken in the Lebesgue sense.

*Proof.* Using the integration by parts, we have

$$\begin{aligned}
 (2.2) \quad & \int_a^x (t-a)^2 [f''(t) - \lambda_1(x)] dt \\
 = & \int_a^x (t-a)^2 f''(t) dt - \lambda_1(x) \int_a^x (t-a)^2 dt \\
 = & (t-a)^2 f'(t) \Big|_a^x - 2 \int_a^x (t-a) f'(t) dt - \frac{\lambda_1(x)}{3} (t-a)^3 \Big|_a^x \\
 = & (x-a)^2 f'(x) - 2 \left[ (t-a) f(t) \Big|_a^x - \int_a^x f(t) dt \right] - \frac{\lambda_1(x)}{3} (x-a)^3 \\
 = & (x-a)^2 f'(x) - 2(x-a) f(x) + 2 \int_a^x f(t) dt - \frac{\lambda_1(x)}{3} (x-a)^3,
 \end{aligned}$$

$$\begin{aligned}
(2.3) \quad & \int_x^{a+b-x} \left(t - \frac{a+b}{2}\right)^2 [f''(t) - \lambda_2(x)] dt \\
&= \int_x^{a+b-x} \left(t - \frac{a+b}{2}\right)^2 f''(t) dt - \lambda_2(x) \int_x^{a+b-x} \left(t - \frac{a+b}{2}\right)^2 dt \\
&= \left(t - \frac{a+b}{2}\right)^2 f'(t) \Big|_x^{a+b-x} - 2 \left(t - \frac{a+b}{2}\right) f(t) \Big|_x^{a+b-x} \\
&\quad + 2 \int_x^{a+b-x} f(t) dt - \frac{\lambda_2(x)}{3} \left(t - \frac{a+b}{2}\right)^3 \Big|_x^{a+b-x} \\
&= \left(\frac{a+b}{2} - x\right)^2 [f'(a+b-x) - f'(x)] \\
&\quad - 2 \left(\frac{a+b}{2} - x\right) [f(a+b-x) + f(x)] + 2 \int_x^{a+b-x} f(t) dt \\
&\quad - \frac{2}{3} \lambda_2(x) \left(\frac{a+b}{2} - x\right)^3
\end{aligned}$$

and

$$\begin{aligned}
(2.4) \quad & \int_{a+b-x}^b (t-b)^2 [f''(t) - \lambda_3(x)] dt \\
&= \int_{a+b-x}^b (t-b)^2 f''(t) dt - \lambda_3(x) \int_{a+b-x}^b (t-b)^2 dt \\
&= (t-b)^2 f'(t) \Big|_{a+b-x}^b - 2(t-b) f(t) \Big|_{a+b-x}^b \\
&\quad + 2 \int_{a+b-x}^b f(t) dt - \frac{\lambda_3(x)}{3} (t-b)^3 \Big|_{a+b-x}^b \\
&= -(x-a)^2 f'(a+b-x) - 2(x-a) f(a+b-x) \\
&\quad + 2 \int_{a+b-x}^b f(t) dt - \frac{\lambda_3(x)}{3} (x-a)^3.
\end{aligned}$$

If we add the equality (2.2)-(2.4) and divide by  $2(b-a)$ , we obtain required identity.  $\square$

**Corollary 2.1.** *Under assumption of Lemma 2.1 with  $\lambda_i(x) = \lambda_i, i = 1, 2, 3$*

*i) if we choose  $x = a$ , we have*

$$\begin{aligned}
 (2.5) \quad & \frac{b-a}{8} [f'(b) - f'(a)] - \frac{f(a) + f(b)}{2} + \frac{1}{b-a} \int_a^b f(t) dt - \frac{(b-a)^2}{24} \lambda_2 \\
 & = \frac{1}{2(b-a)} \int_a^b \left(t - \frac{a+b}{2}\right)^2 [f''(t) - \lambda_2] dt,
 \end{aligned}$$

ii) if we choose  $x = \frac{a+b}{2}$ , we have

$$\begin{aligned}
 (2.6) \quad & \frac{1}{b-a} \int_a^b f(t) dt - f\left(\frac{a+b}{2}\right) - \frac{(b-a)^2}{48} (\lambda_1 + \lambda_3) \\
 & = \frac{1}{2(b-a)} \left[ \int_a^{\frac{a+b}{2}} (t-a)^2 [f''(t) - \lambda_1] dt + \int_{\frac{a+b}{2}}^b (t-b)^2 [f''(t) - \lambda_3] dt \right],
 \end{aligned}$$

iii) if we choose  $x = \frac{3a+b}{4}$ , we have

$$\begin{aligned}
 (2.7) \quad & \frac{1}{b-a} \int_a^b f(t) dt - \frac{1}{2} \left[ f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) \right] - \frac{(b-a)^2}{384} (\lambda_1 + 2\lambda_2 + \lambda_3) \\
 & = \frac{1}{2(b-a)} \left[ \int_a^{\frac{3a+b}{4}} (t-a)^2 [f''(t) - \lambda_1] dt + \int_{\frac{3a+b}{4}}^{\frac{a+3b}{4}} \left(t - \frac{a+b}{2}\right)^2 [f''(t) - \lambda_2] dt \right. \\
 & \quad \left. + \int_{\frac{a+3b}{4}}^b (t-b)^2 [f''(t) - \lambda_3] dt \right].
 \end{aligned}$$

**Corollary 2.2.** If we take  $\lambda_1 = -\lambda_3$  in (2.6), then we get

$$\begin{aligned}
 & \frac{1}{b-a} \int_a^b f(t) dt - f\left(\frac{a+b}{2}\right) \\
 & = \frac{1}{2(b-a)} \left[ \int_a^{\frac{a+b}{2}} (t-a)^2 [f''(t) - \lambda_1] dt + \int_{\frac{a+b}{2}}^b (t-b)^2 [f''(t) + \lambda_1] dt \right],
 \end{aligned}$$

and choosing  $\lambda_1 = \lambda_3 = -\lambda_2$  in (2.7), we have the inequality

$$\begin{aligned}
 & \frac{1}{b-a} \int_a^b f(t) dt - \frac{1}{2} \left[ f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) \right] \\
 & = \frac{1}{2(b-a)} \left[ \int_a^{\frac{3a+b}{4}} (t-a)^2 [f''(t) - \lambda_1] dt + \int_{\frac{3a+b}{4}}^{\frac{a+3b}{4}} \left(t - \frac{a+b}{2}\right)^2 [f''(t) + \lambda_1] dt \right. \\
 & \quad \left. + \int_{\frac{a+3b}{4}}^b (t-b)^2 [f''(t) - \lambda_1] dt \right].
 \end{aligned}$$

### 3. Inequalities for Functions Whose Second Derivatives are Bounded

Recall the sets of complex-valued functions:

$$\overline{U}_{[a,b]}(\gamma, \Gamma) := \left\{ f : [a, b] \rightarrow \mathbb{C} \mid \operatorname{Re} \left[ (\Gamma - f(t)) \overline{(f(t) - \gamma)} \right] \geq 0 \text{ for almost every } t \in [a, b] \right\}$$

and

$$\overline{\Delta}_{[a,b]}(\gamma, \Gamma) := \left\{ f : [a, b] \rightarrow \mathbb{C} \mid \left| f(t) - \frac{\gamma + \Gamma}{2} \right| \leq \frac{1}{2} |\Gamma - \gamma| \text{ for a.e. } t \in [a, b] \right\}.$$

**Proposition 3.1.** [25] For any  $\gamma, \Gamma \in \mathbb{C}$ ,  $\gamma \neq \Gamma$ , we have that  $\overline{U}_{[a,b]}(\gamma, \Gamma)$  and  $\overline{\Delta}_{[a,b]}(\gamma, \Gamma)$  are nonempty and closed sets and

$$\overline{U}_{[a,b]}(\gamma, \Gamma) = \overline{\Delta}_{[a,b]}(\gamma, \Gamma).$$

**Theorem 3.1.** Let  $f : [a, b] \rightarrow \mathbb{C}$  be a twice differentiable function on  $(a, b)$  and  $x \in (a, b)$ . Suppose that  $\gamma_i(x), \Gamma_i(x) \in \mathbb{C}$ ,  $\gamma_i(x) \neq \Gamma_i(x)$ ,  $i = 1, 2, 3$  and  $f'' \in \overline{U}_{[a,x]}(\gamma_1, \Gamma_1) \cap \overline{U}_{[x, a+b-x]}(\gamma_2, \Gamma_2) \cap \overline{U}_{[a+b-x, b]}(\gamma_3, \Gamma_3)$ , then we have the inequality

$$\begin{aligned} & \left| \left( x - \frac{3a+b}{4} \right) \frac{f'(x) - f'(a+b-x)}{2} - \frac{f(x) + f(a+b-x)}{2} + \frac{1}{b-a} \int_a^b f(t) dt \right. \\ & \quad \left. - \frac{1}{12(b-a)} [(x-a)^3 [\gamma_1(x) + \Gamma_1(x) + \gamma_3(x) + \Gamma_3(x)] \right. \\ & \quad \left. + 2 \left( \frac{a+b}{2} - x \right)^3 [\gamma_2(x) + \Gamma_2(x)] \right] \\ & \leq \frac{1}{12(b-a)} [(x-a)^3 (|\Gamma_1(x) - \gamma_1(x)| + |\Gamma_3(x) - \gamma_3(x)|) \\ & \quad + 2 \left( \frac{a+b}{2} - x \right)^3 |\Gamma_2(x) - \gamma_2(x)|] \end{aligned}$$

for all  $x \in [a, \frac{a+b}{2}]$ .

*Proof.* Taking the modulus identity (2.1) for  $\lambda_1(x) = \frac{\gamma_1(x) + \Gamma_1(x)}{2}$ ,  $\lambda_2(x) = \frac{\gamma_2(x) + \Gamma_2(x)}{2}$  and  $\lambda_3(x) = \frac{\gamma_3(x) + \Gamma_3(x)}{2}$ , since  $f'' \in \overline{U}_{[a,x]}(\gamma_1, \Gamma_1) \cap \overline{U}_{[x, a+b-x]}(\gamma_2, \Gamma_2) \cap \overline{U}_{[a+b-x, b]}(\gamma_3, \Gamma_3)$ , we have

$$\begin{aligned}
 & \left| \left( x - \frac{3a+b}{4} \right) \frac{f'(x) - f'(a+b-x)}{2} - \frac{f(x) + f(a+b-x)}{2} + \frac{1}{b-a} \int_a^b f(t) dt \right. \\
 & \left. - \frac{1}{12(b-a)} [(x-a)^3 [\gamma_1(x) + \Gamma_1(x) + \gamma_3(x) + \Gamma_3(x)] \right. \\
 & \left. + 2 \left( \frac{a+b}{2} - x \right)^3 [\gamma_2(x) + \Gamma_2(x)] \right] \Bigg| \\
 & \leq \frac{1}{2(b-a)} \left[ \int_a^x (t-a)^2 \left| f''(t) - \frac{\gamma_1(x) + \Gamma_1(x)}{2} \right| dt \right. \\
 & \left. + \int_x^{a+b-x} \left( t - \frac{a+b}{2} \right)^2 \left| f''(t) - \frac{\gamma_2(x) + \Gamma_2(x)}{2} \right| dt \right. \\
 & \left. + \int_{a+b-x}^b (t-b)^2 \left| f''(t) - \frac{\gamma_3(x) + \Gamma_3(x)}{2} \right| dt \right] \\
 & \leq \frac{1}{4(b-a)} \left[ |\Gamma_1(x) - \gamma_1(x)| \int_a^x (t-a)^2 dt + |\Gamma_2(x) - \gamma_2(x)| \int_x^{a+b-x} \left( t - \frac{a+b}{2} \right)^2 dt \right. \\
 & \left. + |\Gamma_3(x) - \gamma_3(x)| \int_{a+b-x}^b (t-b)^2 dt \right] \\
 & = \frac{1}{12(b-a)} \left[ (x-a)^3 |\Gamma_1(x) - \gamma_1(x)| + 2 \left( \frac{a+b}{2} - x \right)^3 |\Gamma_2(x) - \gamma_2(x)| \right. \\
 & \left. + (x-a)^3 |\Gamma_3(x) - \gamma_3(x)| \right]
 \end{aligned}$$

which completes the proof.  $\square$

**Corollary 3.1.** *Under the assumption of Theorem 3.1,*

*i) if we choose  $x = a$ , then we have*

$$\begin{aligned}
 (3.1) \quad & \left| \frac{b-a}{8} [f'(b) - f'(a)] - \frac{f(a) + f(b)}{2} + \frac{1}{b-a} \int_a^b f(t) dt - \frac{(b-a)^2}{48} [\gamma_2(x) + \Gamma_2(x)] \right| \\
 & \leq \frac{(b-a)^2}{48} |\Gamma_2(x) - \gamma_2(x)|,
 \end{aligned}$$

*ii) if we choose  $x = \frac{a+b}{2}$ , then we have*

$$\begin{aligned}
 (3.2) \quad & \left| \frac{1}{b-a} \int_a^b f(t) dt - f \left( \frac{a+b}{2} \right) - \frac{(b-a)^2}{96} [\gamma_1(x) + \Gamma_1(x) + \gamma_3(x) + \Gamma_3(x)] \right| \\
 & \leq \frac{(b-a)^2}{96} [|\Gamma_1(x) - \gamma_1(x)| + |\Gamma_3(x) - \gamma_3(x)|],
 \end{aligned}$$

iii) if we choose  $x = \frac{3a+b}{4}$ , then we have

$$(3.3) \quad \left| \frac{1}{b-a} \int_a^b f(t) dt - \frac{1}{2} \left[ f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) \right] - \frac{(b-a)^2}{768} (\gamma_1(x) + \Gamma_1(x) + 2\gamma_2(x) + 2\Gamma_2(x) + \gamma_3(x) + \Gamma_3(x)) \right| \\ \leq \frac{(b-a)^2}{768} [|\Gamma_1(x) - \gamma_1(x)| + 2|\Gamma_2(x) - \gamma_2(x)| + |\Gamma_3(x) - \gamma_3(x)|].$$

**Corollary 3.2.** If we choose  $\Gamma_1(x) = -\Gamma_3(x)$  and  $\gamma_1(x) = -\gamma_3(x)$  in (3.2), then

$$\left| \frac{1}{b-a} \int_a^b f(t) dt - f\left(\frac{a+b}{2}\right) \right| \leq \frac{(b-a)^2}{48} [|\Gamma_1(x) - \gamma_1(x)|],$$

and choosing  $\Gamma_1(x) = \Gamma_3(x) = -\Gamma_2(x)$  and  $\gamma_1(x) = \gamma_3(x) = -\gamma_2(x)$  in (3.3), we have

$$\left| \frac{1}{b-a} \int_a^b f(t) dt - \frac{1}{2} \left[ f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) \right] \right| \\ \leq \frac{(b-a)^2}{192} [|\Gamma_1(x) - \gamma_1(x)|].$$

#### 4. Inequalities for Functions Whose Second Derivatives are of Bounded Variation

Assume that  $f : [a, b] \rightarrow \mathbb{C}$  be a twice differentiable function on  $I^\circ$  (the interior of  $I$ ) and  $[a, b] \subset I^\circ$ . Then, from (2.1), we have for  $\lambda_1(x) = f''(a)$ ,  $\lambda_2(x) = \frac{f''(x) + f''(a+b-x)}{2}$  and  $\lambda_3(x) = f''(b)$

$$(4.1) \quad \left( x - \frac{3a+b}{4} \right) \frac{f'(x) - f'(a+b-x)}{2} - \frac{f(x) + f(a+b-x)}{2} + \frac{1}{b-a} \int_a^b f(t) dt \\ - \frac{1}{6(b-a)} \left[ (x-a)^3 (f''(a) + f''(b)) + \left( \frac{a+b}{2} - x \right)^3 [f''(x) + f''(a+b-x)] \right] \\ = \frac{1}{2(b-a)} \left[ \int_a^x (t-a)^2 [f''(t) - f''(a)] dt + \int_x^{a+b-x} \left( t - \frac{a+b}{2} \right)^2 \left[ f''(t) - \frac{f''(x) + f''(a+b-x)}{2} \right] dt \right. \\ \left. + \int_{a+b-x}^b (t-b)^2 [f''(t) - f''(b)] dt \right]$$

for any  $x \in [a, \frac{a+b}{2}]$ .



**Theorem 4.1.** Let  $f : [a, b] \rightarrow \mathbb{C}$  be a twice differentiable function on  $I^\circ$  (the interior of  $I$ ) and  $[a, b] \subset I^\circ$ . If the second derivative  $f''$  is of bounded variation on  $[a, b]$ , then

$$\begin{aligned}
 & \left( \left| \left( x - \frac{3a+b}{4} \right) \frac{f'(x) - f'(a+b-x)}{2} - \frac{f(x) + f(a+b-x)}{2} \right. \right. \\
 & \quad \left. \left. + \frac{1}{b-a} \int_a^b f(t) dt - \frac{1}{6(b-a)} [(x-a)^3 (f''(a) + f''(b)) \right. \right. \\
 & \quad \left. \left. + \left( \frac{a+b}{2} - x \right)^3 [f''(x) + f''(a+b-x)] \right] \right| \\
 & \leq \frac{1}{6(b-a)} \left[ (x-a)^3 \bigvee_a^x (f'') + \left( \frac{a+b}{2} - x \right)^3 \bigvee_x^{a+b-x} (f'') + (x-a)^3 \bigvee_{a+b-x}^b (f'') \right] \\
 & \leq \frac{1}{6(b-a)} \left\{ \begin{aligned} & \max \left\{ (x-a)^3, \left( \frac{a+b}{2} - x \right)^3 \right\} \bigvee_a^b (f''), \\ & \left[ 2(x-a)^3 + \left( \frac{a+b}{2} - x \right)^3 \right] \max \left\{ \bigvee_a^x (f''), \bigvee_x^{a+b-x} (f''), \bigvee_{a+b-x}^b (f'') \right\} \end{aligned} \right\}
 \end{aligned}$$

for any  $x \in [a, b]$ .

*Proof.* Taking modulus (4.1), we get

$$\begin{aligned}
 (4.3) \quad & \left| \left( x - \frac{3a+b}{4} \right) \frac{f'(x) - f'(a+b-x)}{2} - \frac{f(x) + f(a+b-x)}{2} \right. \\
 & \quad \left. + \frac{1}{b-a} \int_a^b f(t) dt - \frac{1}{6(b-a)} [(x-a)^3 (f''(a) + f''(b)) \right. \\
 & \quad \left. + \left( \frac{a+b}{2} - x \right)^3 [f''(x) + f''(a+b-x)] \right] \Big| \\
 & \leq \frac{1}{2(b-a)} \left[ \int_a^x (t-a)^2 |f''(t) - f''(a)| dt \right. \\
 & \quad \left. + \int_x^{a+b-x} \left( t - \frac{a+b}{2} \right)^2 \left| f''(t) - \frac{f''(x) + f''(a+b-x)}{2} \right| dt \right. \\
 & \quad \left. + \int_{a+b-x}^b (t-b)^2 |f''(t) - f''(b)| dt \right] := T.
 \end{aligned}$$

Since  $f''$  is of bounded variation on  $[a, b]$ , we get

$$(4.4) \quad |f''(t) - f''(a)| \leq \bigvee_a^t (f'') \text{ for } t \in [a, x],$$

$$\begin{aligned}
(4.5) \quad & \left| f''(t) - \frac{f''(x) + f''(a+b-x)}{2} \right| \\
& \leq \frac{1}{2} [|f''(t) - f''(x)| + |f''(a+b-x) - f''(t)|] \\
& \leq \frac{1}{2} \bigvee_x^{a+b-x} (f'') \text{ for } t \in [x, a+b-x]
\end{aligned}$$

and

$$(4.6) \quad |f''(t) - f''(b)| \leq \bigvee_t^b (f'') \text{ for } t \in [a+b-x, b].$$

If we put (4.4)-(4.6) in (4.3), we have

$$\begin{aligned}
T & \leq \frac{1}{2(b-a)} \left[ \int_a^x (t-a)^2 \left( \bigvee_a^t (f'') \right) dt + \int_x^{a+b-x} \left( t - \frac{a+b}{2} \right)^2 \left( \frac{1}{2} \bigvee_x^{a+b-x} (f'') \right) dt \right. \\
& \quad \left. + \int_{a+b-x}^b (t-b)^2 \left( \bigvee_t^b (f'') \right) dt \right] \\
& \leq \frac{1}{2(b-a)} \left[ \bigvee_a^x (f'') \int_a^x (t-a)^2 dt + \frac{1}{2} \bigvee_x^{a+b-x} (f'') \int_x^{a+b-x} \left( t - \frac{a+b}{2} \right)^2 dt \right. \\
& \quad \left. + \bigvee_{a+b-x}^b (f'') \int_{a+b-x}^b (t-b)^2 dt \right] \\
& = \frac{1}{6(b-a)} \left[ (x-a)^3 \bigvee_a^x (f'') + \left( \frac{a+b}{2} - x \right)^3 \bigvee_x^{a+b-x} (f'') + (x-a)^3 \bigvee_{a+b-x}^b (f'') \right]
\end{aligned}$$

which completes the proof of the first inequality in (4.2).

The proof of the second inequality in (4.2) is obvious from properties of maximum.  $\square$

**Corollary 4.1.** *Under assumptions of Theorem 4.1,*

*i) if we take  $x = a$ , we have the inequality*

$$\begin{aligned}
& \left| \frac{b-a}{8} [f'(b) - f'(a)] - \frac{f(a) + f(b)}{2} + \frac{1}{b-a} \int_a^b f(t) dt - \frac{(b-a)^2}{48} [f''(a) + f''(b)] \right| \\
& \leq \frac{(b-a)^2}{48} \bigvee_a^b (f'')
\end{aligned}$$

ii) if we take  $x = \frac{a+b}{2}$ , we have

$$\left| \frac{1}{b-a} \int_a^b f(t)dt - f\left(\frac{a+b}{2}\right) - \frac{(b-a)^2}{48} [f''(a) + f''(b)] \right| \leq \frac{(b-a)^2}{48} \mathcal{V}_a^b(f'')$$

iii) if we take  $x = \frac{3a+b}{4}$ , we get

$$\begin{aligned} & \left| \frac{1}{b-a} \int_a^b f(t)dt - \frac{1}{2} \left[ f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) \right] \right. \\ & \quad \left. - \frac{(b-a)^2}{384} \left[ f''(a) + f''(b) + f''\left(\frac{3a+b}{4}\right) + f''\left(\frac{a+3b}{4}\right) \right] \right| \\ & \leq \frac{(b-a)^2}{384} \mathcal{V}_a^b(f''). \end{aligned}$$

#### REFERENCES

1. M. W. Alomari, *A Generalization of weighted companion of Ostrowski integral inequality for mappings of bounded variation*, RGMIA Research Report Collection, 14(2011), Article 87, 11 pp.
2. M. W. Alomari and M.A. Latif, *Weighted Companion for the Ostrowski and the Generalized Trapezoid Inequalities for Mappings of Bounded Variation*, RGMIA Research Report Collection, 14(2011), Article 92, 10 pp.
3. M.W. Alomari and S.S. Dragomir, *Mercer–Trapezoid rule for the Riemann–Stieltjes integral with applications*, Journal of Advances in Mathematics, 2 (2)(2013), 67–85.
4. H. Budak and M.Z. Sarikaya, *On generalization of Dragomir’s inequalities*, RGMIA Research Report Collection, 17(2014), Article 155, 10 pp.
5. H. Budak and M.Z. Sarikaya, *New weighted Ostrowski type inequalities for mappings with first derivatives of bounded variation*, RGMIA Research Report Collection, 18(2015), Article 43, 8 pp.
6. H. Budak and M.Z. Sarikaya, *A new generalization of Ostrowski type inequality for mappings of bounded variation*, RGMIA Research Report Collection, 18(2015), Article 47, 9 pp.
7. H. Budak and M.Z. Sarikaya, *On generalization of weighted Ostrowski type inequalities for functions of bounded variation*, RGMIA Research Report Collection, 18(2015), Article
8. H. Budak and M. Z. Sarikaya, *A new Ostrowski type inequality for functions whose first derivatives are of bounded variation*, Moroccan J. Pure Appl. Anal., 2(1)(2016), 1–11.

9. H. Budak and M.Z. Sarikaya, *A companion of Ostrowski type inequalities for mappings of bounded variation and some applications*, RGMIA Research Report Collection, 19(2016), Article 24, 10 pp.
10. H. Budak, M.Z. Sarikaya and A. Qayyum, *Improvement in companion of Ostrowski type inequalities for mappings whose first derivatives are of bounded variation and application*, RGMIA Research Report Collection, 19(2016), Article 25, 11 pp.
11. H. Budak, M.Z. Sarikaya and S.S. Dragomir, *Some perturbed Ostrowski type inequality for twice differentiable functions*, RGMIA Research Report Collection, 19(2016), Article 47, 14 pp.
12. H. Budak and M. Z. Sarikaya, *Some perturbed Ostrowski type inequality for functions whose first derivatives are of bounded variation*, Submitted
13. S. S. Dragomir and N.S. Barnett, *An Ostrowski type inequality for mappings whose second derivatives are bounded and applications*, RGMIA Research Report Collection, 1(2)(1998) .
14. S. S. Dragomir, *The Ostrowski integral inequality for mappings of bounded variation*, Bulletin of the Australian Mathematical Society, 60(1) (1999), 495-508.
15. S. S. Dragomir, and A. Sofo, *An integral inequality for twice differentiable mappings and application*, Tamkang J. Math., 31(4) 2000.
16. S. S. Dragomir, *On the Ostrowski's integral inequality for mappings with bounded variation and applications*, Mathematical Inequalities & Applications, 4 (2001), no. 1, 59–66.
17. S. S. Dragomir, *A companion of Ostrowski's inequality for functions of bounded variation and applications*, International Journal of Nonlinear Analysis and Applications, 5 (2014) No. 1, 89-97 pp.
18. S. S. Dragomir, *Some perturbed Ostrowski type inequalities for functions of bounded variation*, Preprint RGMIA Research Report Collection, 16 (2013), Art. 93.
19. S. S. Dragomir, *Approximating real functions which possess  $n$ th derivatives of bounded variation and applications*, Computers and Mathematics with Applications 56 (2008) 2268–2278.
20. S. S. Dragomir, *Some perturbed Ostrowski type inequalities for functions of bounded variation*, Asian-European Journal of Mathematics, 8(4)(2015, ),14 pages. DOI:10.1142/S1793557115500692
21. S. S. Dragomir, *Some Perturbed Ostrowski Type Inequalities for Absolutely Continuous Functions (I)*, Acta Universitatis Matthiae Belii, series Mathematics 23(2015), 71–86.
22. S. S. Dragomir, *Some Perturbed Ostrowski Type Inequalities for Absolutely Continuous Functions (II)*, RGMIA Research Report Collection, 16(2013), Article 93, 16 pp.
23. S. S. Dragomir, *Some Perturbed Ostrowski Type Inequalities for Absolutely Continuous Functions (III)*, TJMM, 7(1)(2015),31-43.
24. S. S. Dragomir, *Perturbed Companions of Ostrowski's Inequality for Functions of Bounded Variation*, RGMIA Research Report Collection, 17(2014), Article 1, 16 pp.

25. S. S. Dragomir, *Perturbed Companions of Ostrowski's Inequality for Absolutely Continuous Functions (I)*, RGMIA Research Report Collection, 17(2014), Article 7, 15 pp.
26. S. S. Dragomir, *Perturbed Companions of Ostrowski's Inequality for Absolutely Continuous Functions (II)*, GMIA Research Report Collection, 17(2014), Article 19, 11 pp.
27. W. Liu and Y. Sun, *A Refinement of the Companion of Ostrowski inequality for functions of bounded variation and Applications*, arXiv:1207.3861v1, (2012).
28. Z. Liu, *Some Ostrowski type inequalities*, Mathematical and Computer Modelling 48 (2008) 949–960.
29. A. M. Ostrowski, *Über die absolutabweichung einer differentiebaren funktion von ihrem integralmittelwert*, Comment. Math. Helv. 10(1938), 226-227.
30. A. Rafiq, N.A. Mir and F. Zafar, *A generalized Ostrowski-Grüss Type inequality for twice differentiable mappings and application*, JIPAM, 7(4)(2006), article 124.
31. M. Z. Sarikaya, *On the Ostrowski type integral inequality*, Acta Mathematica Universitatis Comenianae, Vol. LXXIX, 1(2010),129-134.
32. M. Z. Sarikaya and E. Set, *On New Ostrowski Type Integral Inequalities*, Thai Journal of Mathematics, 12(1)(2014) 145-154.
33. E. Set and M. Z. Sarikaya, *On a new Ostroski-type inequality and related results*, Kyungpook Mathematical Journal, 54(2014), 545-554.
34. J. Park, *Some Companions of an Ostrowski-like Type Inequality for Twice Differentiable Functions*, Applied Mathematical Sciences, Vol. 8, 2014, no. 47, 2339 - 2351.
35. M. Liu, Y. Zhu and J. Park, *Some companions of perturbed Ostrowski-type inequalities based on the quadratic kernel function with three sections and applications*, J. of Ineq. and Applications, 2013 2013:226.
36. A. Qayyum, M. Shoaib and I. Faye, *Companion of Ostrowski-type inequality based on 5-step quadratic kernel and applications*, Journal of Nonlinear Science and Applications, 9 (2016), 537–552.
37. A. Qayyum, M. Shoaib and I. Faye, *On new refinements and applications of efficient quadrature rules using n-times differentiable mappings*, RGMIA Research Report Collection, 19(2016), Article 9, 22 pp.

Mehmet Zeki Sarikaya  
Faculty of Science and Arts  
Department of Mathematics  
Düzce University  
Düzce, TURKEY  
sarikayamz@gmail.com

Hüseyin Budak  
Faculty of Science and Arts  
Department of Mathematics

Düzce University  
Düzce, TURKEY  
hsyn.budak@gmail.com

Tuba Tunç  
Faculty of Science and Arts  
Department of Mathematics  
Düzce University  
Düzce, TURKEY  
tubatunc03@gmail.com

Samet Erden  
Faculty of Science  
Department of Mathematics  
Bartın University  
Bartın, TURKEY  
erdem1627@gmail.com

Hatice Yıldız  
Faculty of Science and Arts  
Department of Mathematics  
Düzce University  
Düzce, TURKEY  
yaldizhatice@gmail.com