

## Three-dimensional facial morphometry in patients rehabilitated with implant-supported prostheses

Davide G. Tommasi, Luca Pisoni, Giuseppe C. Lanza Attisano, Alberto Rossetti, Claudia Dolci and Virgilio F. Ferrario

Dept. Biomedical Sciences for Health, Università degli Studi di Milano, Milano, Italy

The aim of the present study was to assess a low-cost, non-invasive facial morphometric digitizer to assist the practitioner in three-dimensional soft-tissue changes before and after oral rehabilitation. The method should provide quantitative data to support an objective assessment of the facial esthetic outcome [1].

Twenty-two patients aged 45-82 years, all with edentulous maxilla and mandible, were assessed both before and after receiving their definitive complete implant-supported prostheses (each received 4-11 implants in each dental arch; full-arch fixed prostheses were made). The three-dimensional coordinates of 50 soft-tissue facial landmarks were collected with a non-invasive digitizer; labial and facial areas, volumes, angles and distances were compared without/ with the prostheses [2].

Dental prostheses induced significant reductions in the nasolabial, mentolabial and interlabial angles, with increased labial prominence ( $p < 0.05$ , Wilcoxon test). Lip vermilion area and volume significantly increased; significant increments were found in the vertical and anteroposterior labial dimensions. The presence of the dental prostheses significantly ( $p < 0.001$ ) modified the three-dimensional positions of several soft-tissue facial landmarks.

The current approach enabled quantitative evaluation of the final soft-tissue results of oral rehabilitation with implant-supported prostheses, without submitting the patients to invasive procedures. The method could assess the three-dimensional appearance of the facial soft tissues of the patient while planning the provisional prosthetic restoration, providing quantitative information to prepare the best definitive prosthesis.

Dote ricerca: FSE, Regione Lombardia

### References

- [1] Bidra (2011) Three-dimensional esthetic analysis in treatment planning for implant-supported fixed prosthesis in the edentulous maxilla: review of the esthetics literature. *J Esthet Restor Dent* 23: 219-237.
- [2] Sforza et al. (2010) Age- and sex-related changes in three-dimensional lip morphology. *Forensic Sci Int* 200:182.e1-182.e7.

Keywords: Computerized anthropometry, face, oral rehabilitation, implant-supported prostheses.