

The anastomotic network around the anterior superior alveolar nerve: an anatomical and radiological study

Marco Ferrari¹ - Alberto Schreiber² - Davide Lancini¹ - Stefania Castrezzi¹ - Lena Hirtler³ - Elena Mercandelli¹ - Roberto Maroldi⁴ - Piero Nicolai² - [Luigi F. Rodella](#)¹

¹ Section of Anatomy and Physiopathology, Department of Clinical and Experimental Sciences, University of Brescia, Brescia, Italia – ²Unit of Otorhinolaryngology-Head and Neck Surgery, Department of Medical and Surgical Specialties, Radiological Sciences, and Public Health, University of Brescia, Italia – ³Center for Anatomy and Cell Biology, Department for Systematic Anatomy, Medical University Vienna, Vienna, Austria – ⁴Unit of Radiology, Department of Medical and Surgical Specialties, Radiological Sciences and Public Health, University of Brescia, Brescia, Italia

Innervation of superior teeth is supplied by the posterior (PSAN), anterior (ASAN) and sometimes by middle superior alveolar nerve (MSAN). PSAN arises from the maxillary nerve and passes through the posterolateral maxillary wall towards the posterior teeth. ASAN arises from the anterior portion of the infraorbital nerve and courses within the infraorbital canal passing nearby the piriform aperture and premaxilla. When present, MSAN arises from the posterior portion of the infraorbital nerve and runs along the lateral maxillary wall. However, an additional nasopalatine or sublabial injection is frequently required to obtain a complete anesthesia of the maxillary teeth due to rich anastomotic network (1-2). With the aim to better describe the complexity of the superior alveolar nerve network, fifty-seven high-definition sinonasal cone-beam CT (CBCT) were analyzed. PSAN, ASAN and MSAN were detected by specific bony landmarks/canals and nervous anastomoses were accurately evaluated. In addition, medial anastomotic branches from the palatal and/or nasal nervous plexi were also considered. PSAN and ASAN were identified in 100% of cases whereas MSAN in 19.6% of cases. Anastomotic branch versus ASAN was identified in all cases from MSAN and in 60.3% from PSAN. Medial anastomotic branch was detected in 62.0% of cases from the nasal plexus and in 6.2% from the palatal plexus: the former passed through a bony defect in the floor of the piriform aperture or at the base of the nasal septum; the latter passed through a tiny canal in the interface between maxilla and premaxilla. These data confirm that maxillary teeth innervation, especially for incisor teeth, could be provided not only by alveolar nerves but also from palatal and nasal plexi via small branches running within maxillary bony canals. These results support the need of additional anesthetic injection to obtain adequate anesthesia of the maxillary teeth; moreover, the role of CBCT in the identification of the nervous pattern was underlined.

References

- [1] Rodella et al. (2007) A review of the mandibular and maxillary nerve supplies and their clinical relevance. *Arch Oral Biol* 57: 323; doi: 10.1016/j.archoralbio.2011.09.007.
- [2] Mayer et al. (2007) Effectiveness of nasopalatine nerve block for anesthesia of maxillary central incisors after failure of the anterior superior alveolar nerve block technique. *Braz Dent J* 18: 69; doi: 10.1590/S0103-64402007000100015.

Keywords

Superior alveolar nerve; anastomosis; CBCT.