

Morphometric analysis of Huguier's canal by Cone Beam CT

Barbara Buffoli¹, Massimo Galli², Andrea Papini³, Marco Ferrari¹, Francesco Belotti¹, Mauro Labanca¹, Manfred Tschabitscher¹ and Luigi Fabrizio Rodella¹

¹ Section of Anatomy and Physiopathology, Department of Clinical and Experimental Sciences, University of Brescia, Viale Europa 11, 25123 Brescia, Italy

² Private Oral Surgeon, Castagnola Foundation, Italy

³ Private Dentist Gnathologist, Castagnola Foundation, Italy

The middle ear and the stomatognathic system are closely anatomically and functionally related. The anterior chordal canal of Huguier connects the temporomandibular joint (TMJ) and the middle ear. This canal is formed by the inferior process of tegment tympani and the sphenoid bone and it is located at the medial end of the petrotympanic fissure. To date, few studies aimed to describe Huguier's canal morphology and its related structures (Toth et al., 2006; Sato et al., 2008; Aristeguieta et al., 2009). The aim of this study is to describe the radiological anatomy of the Huguier's canal using cone beam CT (CBCT, Scanora 3D, Soredex). We measured 438 Huguier's canals from 219 human skulls (Section of Anthropology and Ethnology, Museum of Natural History, Florence, Italy). The measurements were made at three levels: 1) near the TMJ (lateral-glenoidal side) that was 1.961 ± 0.472 mm; 2) the narrowest point of the middle area that was 0.494 ± 0.24 mm; 3) near the middle ear (medial, acoustic meatus side) that was 1.085 ± 0.354 mm. 21 on 439 Huguier's canal (4.79%) were ossificated: 1 only in the medial side, 11 only in the middle area and 9 in all the three levels. Considering the high number of measurements, the values obtained were comparable, suggesting that CBCT can be useful to detect these anatomical details.

References

- [1] Toth et al. (2006) Development of the anterior chordal canal. *Ann Anat* 188: 7-11.
- [2] Sato et al. (2008) Classifications of tunnel-like structure of human petrotympanic fissure by cone beam CT. *Surg Radiol Anat* 30: 323-326.
- [3] Aristeguieta et al. (2009) A direct anatomical study of the morphology and functionality of dis-co-malleolar and anterior malleolar ligaments. *Int J Morphol* 27: 367-379.

Key words

Huguier's canal, middle ear, morphometric analysis.