

Cytotoxic activity of a plant extract on cancer cells

Enrico Catalano¹, Paolo Maestri², Andrea Cochis¹, Barbara Azzimonti¹, Elena M. Varoni^{1,3,4}, Franco Fraschini², Lia Rimondini¹, Saverio Megna² and Marcello Iriti⁵

¹ Department of Health Sciences, University of Piemonte Orientale 'Amedeo Avogadro', via Solaroli 17, 28100 Novara, (Italy)

² Department of Medical Biotechnology and Translational Medicine, Milan State University

³ Department of Biomedical, Surgical and Dental Sciences, Milan State University

⁴ Department of Mining and Materials Engineering, McGill University, Montreal, Canada

⁵ Department of Agricultural and Environmental Sciences, Milan State University, via G. Celoria 2, 20133 Milan, (Italy), marcello.iriti@unimi.it

Chemoprevention by natural products may be considered a promising approach to cancer control and management [1]. Many studies have demonstrated antiproliferative, cytostatic and cytotoxic activities of phytochemicals against cancer cells [2]. In this study, a plant extract from *Arctium lappa*, *Berberis vulgaris* and *Eschscholtia californica* was tested as potential anticancer agent. The antitumoral activity of this plant extract was tested on four human cancer cell lines: MCF-7 (breast carcinoma cells), Huh-7 (hepatic carcinoma cells), HTB-43 (oropharyngeal carcinoma cells) and ECV-304 (urinary bladder carcinoma cells). The efficacy of the extract was compared to the common chemotherapeutic agent cyclophosphamide. Three plant extract concentrations were tested: 800, 650 and 450 ng/ml; for cyclophosphamide, three concentrations were assayed, according to literature data: 1300, 1000 and 850 ng/ml [3]. In addition, plant extract and cyclophosphamide were tested on two primary cell lines as controls, human gingival fibroblasts and human mammary fibroblasts. Cell viability was evaluated by the MTT [(3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, Sigma] colorimetric assay and the new xCELLigence system (Roche) for real-time monitoring of cell viability. All concentrations of plant extract exhibited a high level of cytotoxicity on MCF-7, Huh-7, HTB-43 and ECV-304 cancer cells, similar to cyclophosphamide, though they slightly reduced viability of human gingival and mammary fibroblasts. Conversely, the conventional chemotherapeutic drug showed a marked cytotoxicity on control cells. The potential of the plant extract has been demonstrated *in vitro* on various types of cancers, suggesting a possible use of this natural product as a promising anticancer agent. Further studies are needed to ascertain its efficacy *in vivo* and to elucidate its mechanism(s) of action at molecular and biochemical levels.

References

- [1] Karikas et al. (2010) Anticancer and chemopreventing natural products: some biochemical and therapeutic aspects. *J Buon* 15: 627-638.
- [2] Gordaliza et al. (2007) Natural products as leads to anticancer drugs. *Clin Transl Oncol* 9: 767-776.
- [3] Vetvicka et al. (2012) Procathepsin D involvement in chemoresistance of cancer cells. *N Am J Med Sci* 4: 174-179.

Key words

Antitumor activity, anticancer agents, xCELLigence System, phytotherapy, cancer chemoprevention, natural products.