

The role of caveolin-1 in the regulation of angiogenesis

Alessio D'Alessio and Gigliola Sica

Istituto di Istologia ed Embriologia - Facoltà di Medicina e Chirurgia "A. Gemelli", Roma, Italy

Caveolin-1 (cav-1) is the principal structural component of caveolae which functions as scaffolding protein for the integration of a variety of signaling pathways. In this study, we show that siRNA-induced cav-1 down regulation in human endothelial cells (EC) increased cell size and provoked cell cycle arrest at G1/S phase transition. In addition, silencing of cav-1 reduced matrix metalloproteinases (MMPs) activity which, in turn, affected cell migration and VEGF-induced tube formation of EC in vitro. These data indicate that proper expression of cav-1 is required for maintaining typical functions of EC such as proliferation and the formation of new blood vessels. In addition, we observed a marked increase of cell size, after cav-1 silencing, which might indicate the involvement of this scaffolding protein in the way by which cells perceive changes in their microenvironment. In conclusion, this study proposes cav-1 as an interesting target molecule for studying cellular mechanisms which occur in physiological as well as pathological conditions such as senescence and tumorigenesis.

References

- D'Alessio, A., et al. (2010) Targeting of tumor necrosis factor receptor 1 to low density plasma membrane domains in human endothelial cells. *J Biol Chem.* 285(31): p. 23868-79.
- D'Alessio, A., et al. (2005) Caveolae Participate in Tumor Necrosis Factor Receptor 1 Signaling and Internalization in a Human Endothelial Cell Line. *Am J Pathol.* 166(4): p. 1273-1282.

Keywords: Caveolin, endothelial cells, angiogenesis.