

Sella turcica dimensions between 7 and 13 years: a novel radiographic method for age estimation

Daniele Gibelli¹, Elisa Cerutti¹, Laura Spagnoli¹, Matteo Zago¹, Cristina Cattaneo², Chiarella Sforza¹

¹ Dipartimento di Scienze Biomediche per la Salute, Università degli Studi di Milano, Milan, Italy - ² LABANOF, Laboratorio di Antropologia e Odontologia Forense, Dipartimento di Scienze Biomediche per la Salute, Università degli Studi di Milano, Milan, Italy

Age estimation of the living is becoming a relevant field of application of forensic anthropology, where the development of novel methods is urging (1). Several articles highlight the relation with age of different metrical parameters of the sella turcica (2,3), although no study so far has produced regression formulae which may be applicable to the forensic context. The aim of this study was to evaluate the metrical standards of three different measurements of sella turcica (length, depth, diameter) in 177 lateral cephalometric radiograms belonging to male and female individuals aged between 7 and 13 years in order to find standards useful for age estimation. Results were evaluated by two-way ANOVA. Sella diameter was significantly larger in older children ($p < 0.01$), while sella depth was larger in males than in females ($p < 0.01$); there was no significant gender \times age interaction in any variable. Principal component analysis (PCA) was conducted to highlight the sources of variability in data. The first principal component accounted for 76% of the overall variance and it was closely correlated with length and diameter ($r = 0.93$ and $r = 0.92$, respectively, $p < 0.01$). The linear regression model fitted on age and diameter measures yielded the following equation: $\text{age (years)} = 3.81 * \text{diameter (cm)} + 6.12$. Slope and intercept 95% CI were respectively 4.64 to 7.61 years/cm and 2.34 to 5.28 years. The related coefficient of determination was $R^2 = 0.123$, while the root mean square error was 1.74 years. The present results provide a novel method useful for age estimation in the living minors: further studies are needed in order to test its applicability to the forensic scenario.

References

- [1] Cunha et al. (2009) The problem of aging human remains and living individuals: a review. *Forensic Sci Int* 193:1-13 10.1016/j.forsciint.2009.09.008
- [2] Axelsson et al. (2004) Post-natal size and morphology of the sella turcica. Longitudinal cephalometric standards for Norwegians between 6 and 21 years of age. *Eur J Orthod* 26: 613-21
- [3] Silverman (1957) Roentgen standards for size of the pituitary fossa from infancy through adolescence. *Am J Roentgenol Radium Ther Nucl Med* 78: 451-60

Keywords

Anatomy; age estimation; subadults; sella turcica; X-ray examination.