



ISSN 1678-3921

Journal homepage: www.embrapa.br/pabFor manuscript submission and journal contents, access: www.scielo.br/pab

Pomology/ Scientific Notes

Physicochemical attributes and functional properties of flowers of Brazilian feijoa genotypes


Abstract – The objective of this work was to characterize entire flowers of the feijoa (*Acca sellowiana*) genotypes 'Alcântara', 'Helena', 'Mattos', 'Nonante', and the accession 2316 for dry matter, total antioxidant activity (TAA), and the contents of soluble solids, ascorbic acid, total anthocyanins, total flavonoids, and total phenolic compounds (TPC), as well as for the petal surface areas and colors. The TAA was quantified by the ABTS and DPPH methods. Flowers of the accession 2316 and of the cultivars Alcântara and Helena showed the highest contents of anthocyanins and flavonoids, as well as the highest values of TPC and TAA. The feijoa genotypes cultivated in Southern Brazil show high contents of soluble solids and represent a natural source of antioxidants.

Index terms: *Acca sellowiana*, anthocyanins, ascorbic acid, flavonoids, soluble solids, total antioxidant activity, total phenolic compounds.

Atributos físico-químicos e propriedades funcionais de flores de genótipos brasileiros de goiabeira-serrana

Resumo – O objetivo deste trabalho foi caracterizar flores inteiras dos genótipos de goiabeira-serrana (*Acca sellowiana*) 'Alcântara', 'Helena', 'Mattos', 'Nonante' e o acesso 2316 quanto à matéria seca, à atividade antioxidante total (AAT), e aos teores de sólidos solúveis, ácido ascórbico, antocianinas totais, flavonoides totais e compostos fenólicos totais (CFT), assim como as áreas e as cores das pétalas. A AAT foi quantificada por meio dos métodos ABTS e DPPH. As flores do acesso 2316 e das cultivares Alcântara e Helena apresentaram os maiores teores de antocianinas e flavonoides, bem como os maiores valores de CFT e AAT. Os genótipos de goiabeira-serrana cultivados no Sul do Brasil apresentam alto teor de sólidos solúveis e representam uma fonte natural de antioxidantes.

Termos para indexação: *Acca sellowiana*, antocianinas, ácido ascórbico, flavonoides, sólidos solúveis, atividade antioxidante total, compostos fenólicos totais.

Cassandro Vidal Talamini do Amarante⁽¹⁾ ,
Alexandra Goede de Souza⁽¹⁾,
Thalita Dal Toé Benincá⁽¹⁾,
Cristiano André Steffens⁽¹⁾ and
Marlise Nara Ciotta⁽²⁾

⁽¹⁾ Universidade do Estado de Santa Catarina, Centro de Ciências Agroveterinárias, Avenida Luiz de Camões, nº 2.090, CEP 88520-000 Lages, SC, Brazil. E-mail: cassandro.amarante@udesc.br, alexandra.goede@gmail.com, thalitabeninca@hotmail.com, cristiano.steffens@udesc.br

⁽²⁾ Empresa de Pesquisa Agropecuária e Extensão Rural de Santa Catarina, Estação Experimental de São Joaquim, Rua João Araújo Lima, nº 102, Jardim Caiçara, CEP 88600-000 São Joaquim, SC, Brazil. E-mail: marlise@epagri.sc.gov.br

✉ Corresponding author

Received
February 25, 2019

Accepted
March 15, 2019

How to cite
AMARANTE, C.V.T. do; SOUZA, A.G. de; BENINCÁ, T.D.T.; STEFFENS, C.A.; CIOTTA, M.N. Physicochemical attributes and functional properties of flowers of Brazilian feijoa genotypes. *Pesquisa Agropecuária Brasileira*, v.54, e00445, 2019. DOI: <https://doi.org/10.1590/S1678-3921.pab2019.v54.00445>.

Feijoa [*Acca sellowiana* (O. Berg.) Burret, Myrtaceae] is native to Southern Brazil and to the Northeastern of Uruguay (Ducroquet & Hickel, 1991). The fruit has a singular sweet and sour taste and a very aromatic flavor (Amarante et al., 2017a), and its flowers are edible (Souza et al., 2016). Edible flowers, besides enhancing the appearance of prepared foods, are also an important source of bioactive compounds that help maintain and improve human health (Bazylko et al., 2014). These include phenols that have been attracting particular attention



for their antioxidant activity and ability to scavenge different free radicals, leading to the protection of biological molecules against oxidation (Li et al., 2014). However, as far as we know, there are no reports concerning the characterization of phytochemical composition of feijoa flowers and the beneficial effects of its consumption.

The objective of this work was to characterize entire flowers of the feijoa genotypes 'Alcântara', 'Helena', 'Mattos', 'Nonante', and the accession 2316 for dry matter, total antioxidant activity, and the contents of soluble solids, ascorbic acid, total anthocyanins, total flavonoids, and total phenolic compounds, as well as for the petal surface areas and colors.

Feijoa flowers were harvested early in the morning (between 8:00 and 10:00 h) from the orchard of Empresa de Pesquisa Agropecuária e Extensão Rural de Santa Catarina (Epagri), in the municipality of São Joaquim, in the state of Santa Catarina, Brazil (28°16'40"S, 49°56'09"W, at 1,400 m altitude), in 2012, 2013, and 2014. Newly opened flowers (at the phenological stage F1, as described by Ducroquet & Hickel, 1991) were collected from the cultivars Alcântara, Helena, Mattos and Nonante, and from the accession 2316; the samples were then placed in trays, and immediately transported, under refrigeration (~10°C), to the laboratory of postharvest physiology and technology, in the Universidade do Estado de Santa Catarina (Udesc), for analysis.

Flower petals were assessed for their colors and surface areas, while the entire flowers were assessed for dry matter (DM), as well as for their contents of soluble solids (SSC), ascorbic acid, anthocyanins, flavonoids, and total phenolic compounds (TPC), and their total antioxidant activity (TAA).

The colors of the petals were determined on the adaxial (colorful) surface for lightness (L), chroma (C), and hue angle (h°), using a Minolta CR 400 colorimeter (Konica Minolta, Tokyo, Japan). Petal area (cm²) was assessed using a LI-COR leaf area integrator (Model LI-3050A, Lincoln, NE, USA). DM (%) was quantified after leaving the material in an oven at 70°C for 72 hours, until constant mass was obtained. SSC (°Brix) was assessed in the extract (undiluted, total volume) collected from the entire flowers (obtained by grinding the tissue with pestle and mortar), using a digital refractometer (Atago, Model PR201α, Tokyo, Japan). The ascorbic acid content [mg 100 g⁻¹ fresh

matter (FW)] was determined by a colorimetric method using 2,4-dinitrophenyl hydrazine, as described by Amarante et al. (2017a).

Anthocyanins and flavonoids were quantified (mg 100 g⁻¹ FW) according to the method described by Lees & Francis (1972).

The procedure to obtain the hydroalcoholic extract for quantification of TPC and TAA was described by Amarante et al. (2017b). TPC [mg gallic acid equivalents (GAE) 100 g⁻¹ FM] was quantified using the Folin-Ciocalteu method, and TAA was determined based on the ability of the extract to scavenge the radicals ABTS [2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid)] and DPPH (1,1-diphenyl-2-picrylhydrazyl), as described by Amarante et al. (2017b). TAA was expressed as Trolox equivalent antioxidant capacity (TEAC, μmol Trolox g⁻¹ FW) for ABTS method, and as EC₅₀ (substrate concentration which leads to 50% reduction of DPPH, determined as mg FW g⁻¹ DPPH) for DPPH method.

The experiment was carried out in a completely randomized design with five genotypes and four replicates (with 30 flowers and 120 petals by replicate). Since data obtained for all attributes were consistent during the three-year period, only the average values for the three harvests were subjected to the analysis of variance by the SAS software (SAS Institute Inc., Cary, NC, USA), and the means of the treatments were compared by Tukey's test, at 5% probability.

The petals of feijoa flowers showed an intense color and variations of tonality among the genotypes (Table 1). The adaxial (colorful) surface of the accession 2316 showed an intense dark-bluish violet color, characterized by lower h° and L values and higher C values than the other genotypes. 'Helena' showed an intense reddish violet (or pink) color, characterized by higher h° and C values and intermediate L values, compared with other genotypes. 'Mattos', 'Nonante', and 'Alcântara' showed an intermediate coloring between dark-bluish violet and reddish violet. The vivid and vibrant color of feijoa petals is very attractive for human consumption, and this attribute is highly desirable for edible flowers used to decorate dishes, salads, and desserts (Lee et al., 2011; Li et al., 2014).

The area of four petals (number of petals within each flower) ranged from 1.59 cm² in 'Nonante' to 3.91 cm² in 'Mattos', and DM of entire flowers varied from 10.5 g in 'Mattos' to 13.3 g in the accession 2316 (Table

1). The SSC of entire flowers ranged from 10.10 °Brix in 'Mattos' to 14.17 °Brix in 'Helena' (Table 1), which characterizes the flowers of these genotypes as sweet and attractive for human consumption.

Ascorbic acid content of the entire flowers ranged from 18.6 mg 100 g⁻¹ FW ('Helena') to 32.7 mg 100 g⁻¹ FW ('Alcântara' and 'Nonante') (Table 1). These values are lower than those reported for flowers of *Malva sylvestris* (111 mg 100 g⁻¹ FW) (Barros et al., 2010), and *Tropaeolum majus* (1,810 mg 100 g⁻¹ FW) (Bazylo et al., 2014). However, they are higher than the ascorbic acid content reported for fruits, such as passion fruit (12.2 mg 100 g⁻¹ FW), pineapple (11 mg 100 g⁻¹ FW), starfruit (2.73 mg 100 g⁻¹ FW), and avocado (8.53 mg per 100 g FW) (Valente et al., 2011).

Anthocyanin contents ranged from 3.79 mg 100 g⁻¹ FW in the genotype 'Mattos' to 10.15 mg 100 g⁻¹ FW in the accession 2316 (Table 2). Flavonoid contents ranged from 7.99 mg 100 g⁻¹ FW in 'Mattos' to 12.18 mg 100 g⁻¹

FW in the genotype 'Helena' (Table 2). The TPC ranged from 50.8 mg GAE 100 g⁻¹ FW ('Mattos') to 70.1 mg GAE 100 g⁻¹ FW (accession 2316) (Table 2). There was a linear relationship between anthocyanin contents and TPC ($y = 2.4127x + 41.632$, $R^2 = 0.8819$) and between flavonoid contents and TPC ($y = 3.3055x + 26.267$, $R^2 = 0.7509$). Anthocyanin contents and TPC were highly correlated in *Sophora viciifolia* flowers (Tai et al., 2011) and in petals of *Rosa hybrida* cv. Noblered (Lee et al., 2011).

TAA was significantly different between genotypes, as 'Mattos' showed the lowest TAA values, and the accession 2316 the highest ones, regardless of the assay method used (DPPH or ABTS) (Table 2). There was a linear relationship between TPC and TAA, assessed by ABTS and DPPH methods (Figure 1). Therefore, phenolic compounds are the major contributors to the TAA in feijoa flowers, as reported for other flowers (Li et al., 2014).

Table 1. Color attributes (lightness = L; chroma = C; and hue angle = h°) on the adaxial (colorful) surface of petals, petal area, and dry matter, and contents of soluble solids (SSC) and ascorbic acid of the entire flower of Brazilian feijoa (*Acca sellowiana*) genotypes⁽¹⁾.

Genotype	Petal				Entire flower		
	Color			Area ⁽²⁾ (cm ²)	Dry matter (g)	SSC (°Brix)	Ascorbic acid (mg 100 g ⁻¹ FW)
	L	C	h°				
Alcântara	44.26c	27.89a	353.96bc	2.87c	12.321ab	11.03c	32.724a
Mattos	52.62a	21.45b	348.48c	3.91a	10.487c	10.10d	22.333b
Helena	41.62cb	29.31a	359.17a	3.13bc	11.823b	14.17a	18.578b
Nonante	48.98b	17.91b	352.46bc	1.59d	11.414bc	12.42b	32.714a
Accession 2316	39.48d	28.06a	335.13d	3.49ab	13.311a	11.15c	19.912b
Mean	45.39	24.91	349.84	2.99	11.871	11.75	25.252
CV (%)	18.1	36.9	48.7	7.18	4.59	1.85	12.68

⁽¹⁾Means followed by equal letters within the columns do not differ by Tukey's test, at 5% probability. ⁽²⁾Area of four petals (number of petals within each flower). The means refer to averages from 2012, 2013, and 2014.

Table 2. Anthocyanins, flavonoids, total phenolic compounds, and total antioxidant activity (by ABTS and DPPH methods) of entire flowers of Brazilian feijoa (*Acca sellowiana*) genotypes⁽¹⁾.

Genotype	Anthocyanins (mg 100 g ⁻¹ FW)	Flavonoids (mg 100 g ⁻¹ FW)	Total phenolic compounds (mg GAE 100 g ⁻¹ FW) ⁽²⁾	Total antioxidant activity	
				ABTS (µmol Trolox g ⁻¹ FW)	DPPH (EC ₅₀ ⁽³⁾ ; mg FW g ⁻¹ DPPH)
Alcântara	8.77b	11.22a	61.223c	22.858b	4.717bc
Mattos	3.79d	7.99c	50.817d	14.057d	9.014a
Helena	10.12a	12.18a	63.186b	20.512c	5.147bc
Nonante	5.27c	8.19b	54.752d	14.397d	7.093b
Accession 2316	10.15a	11.47a	70.098a	23.744a	4.071c
Mean	7.70	10.30	60.015	19.114	6.008
CV (%)	35.9	19.1	11.5	22.2	44.0

⁽¹⁾Means followed by equal letters within the columns do not differ by Tukey's test, at 5% probability. The means refer to averages from 2012, 2013, and 2014. ⁽²⁾GAE, gallic acid equivalent. ⁽³⁾EC₅₀, substrate concentration which leads to 50% reduction of the DPPH.

The values of TPC (from 50.8 to 70.10 mg GAE 100 g⁻¹ FW) and TAA assessed by the ABTS method (from 14.1 to 23.7 μmol Trolox g⁻¹ FW) (Table 2) are within those reported for other edible flowers. Li et al. (2014) evaluated 51 edible and wild flowers, and found TPC values from 13.0 to 1,148 mg GAE 100 g⁻¹ FW, and TAA (ABTS method) from 0.23 to 175.39 μmol Trolox g⁻¹ FW.

Physicochemical attributes and functional properties of feijoa flowers differed between the evaluated cultivars. Genotypes with high contents of anthocyanins and flavonoids, and high TPC, showed also a high TAA in the entire flower (accession 2316, 'Alcântara' and 'Helena'). Feijoa flowers show high SSC contents, and represent an important source of phenolic compounds with antioxidant activity.




Figure 1. Relationship between total phenolic content and total antioxidant activity assessed by ABTS (A) and DPPH (B) methods of entire flowers of five Brazilian genotypes of feijoa (*Acca sellowiana*). GAE, gallic acid equivalent. EC₅₀, substrate concentration which leads to 50% reduction of the DPPH. The means refer to averages from 2012, 2013, and 2014.

Acknowledgments

To Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Process number 301595/2013-5), for financial support; and to Empresa de Pesquisa Agropecuária e Extensão Rural de Santa Catarina (Epagri), Experimental Station of São Joaquim, for supplying feijoa flowers.

References

- AMARANTE, C.V.T. do; SOUZA, A.G. de; BENINCÁ, T.D.T.; STEFFENS, C.A. Fruit quality of Brazilian genotypes of feijoa at harvest and after storage. *Pesquisa Agropecuária Brasileira*, v.52, p.734-742, 2017a. DOI: <https://doi.org/10.1590/S0100-204X2017000900005>.
- AMARANTE, C.V.T. do; SOUZA, A.G. de; BENINCÁ, T.D.T.; STEFFENS, C.A. Phenolic content and antioxidant activity of fruit of Brazilian genotypes of feijoa. *Pesquisa Agropecuária Brasileira*, v.52, p.1223-1230, 2017b. DOI: <https://doi.org/10.1590/S0100-204X2017001200011>.
- BARROS, L.; CARVALHO, A.M.; FERREIRA, I.C.F.R. Leaves, flowers, immature fruits and leafy flowered stems of *Malva sylvestris*: a comparative study of the nutraceutical potential and composition. *Food and Chemical Toxicology*, v.48, p.1466-1472, 2010. DOI: <https://doi.org/10.1016/j.fct.2010.03.012>.
- BAZYLKO, A.; PARZONKO, A.; JEZ, W.; OSINSKA, E.; KISS, A.K. Inhibition of ROS production, photoprotection, and total phenolic, flavonoids and ascorbic acid content of fresh herb juice and extracts from the leaves and flowers of *Tropeaeolum majus*. *Industrial Crops and Products*, v.55, p.19-24, 2014. DOI: <https://doi.org/10.1016/j.indcrop.2014.01.056>.
- DUCROQUET, J.-P.H.J.; HICKEL, E.R. Fenologia da goiabeira serrana (*Feijoa sellowiana* Berg) no Alto Vale do Rio do Peixe, Santa Catarina. *Revista Brasileira de Fruticultura*, v.13, p.313-320, 1991.
- LEE, J.H.; LEE, H.-J.; CHOUNG, M.-G. Anthocyanin compositions and biological activities from the red petals of Korean edible rose (*Rosa hybrida* cv. Noblered). *Food Chemistry*, v.129, p.272-278, 2011. DOI: <https://doi.org/10.1016/j.foodchem.2011.04.040>.
- LEES, D.H.; FRANCIS, F.J. Standardization of pigment analyses in cranberries. *HortScience*, v.7, p.83-84, 1972.
- LI, A.-N.; LI, S.; LI, H.-B.; XU, D.-P.; XU, X.-R.; CHEN, F. Total phenolic contents and antioxidant capacities of 51 edible and wild flowers. *Journal of Functional Foods*, v.6, p.319-330, 2014. DOI: <https://doi.org/10.1016/j.jff.2013.10.022>.
- SOUZA, A.G. de; AMARANTE, C.V.T. do; STEFFENS, C.A.; BENINCÁ, T.D.T.; PADILHA, M. Postharvest quality of feijoa flowers treated with different preservative solutions and 1-methylcyclopropene. *Revista Brasileira de Fruticultura*, v.38, e-759, 2016. DOI: <https://doi.org/10.1590/0100-29452016759>.

TAI, Z.; CAI, L.; DAI, L.; DONG, L.; WANG, M.; YANG, Y.; CAO, Q.; DING, Z. Antioxidant activity and chemical constituents of edible flower of *Sophora viciifolia*. **Food Chemistry**, v.126, p.1648-1654, 2011. DOI: <https://doi.org/10.1016/j.foodchem.2010.12.048>.

VALENTE, A.; ALBUQUERQUE, T.G.; SANCHES-SILVA, A.; COSTA, H.S. Ascorbic acid content in exotic fruits: a contribution to produce quality data for food composition databases. **Food Research International**, v.44, p.2237-2242, 2011. DOI: <https://doi.org/10.1016/j.foodres.2011.02.012>.
