

ACUTE GASTROINTESTINAL ILLNESS IN TWO INUIT COMMUNITIES: BURDEN OF ILLNESS IN RIGOLET AND IQALUIT, CANADA

**S.L. HARPER¹, V.L. EDGE^{1,2}, J. FORD³, M.K. THOMAS^{1,4}, D.L. PEARL¹, IHACC⁵,
RICG⁶, S.A. MCEWEN¹**

¹ Department of Population Medicine, University of Guelph, Guelph, Ontario, Canada

² Office of Public Health Practice, Public Health Agency of Canada, Guelph, Ontario, Canada

³ Department of Geography, McGill University, Montreal, Quebec, Canada

⁴ Centre for Food-borne, Environmental & Zoonotic Infectious Diseases, Public Health Agency of Canada, Guelph, Ontario, Canada

⁵ Indigenous Health Adaptation to Climate Change Research Group: Lea Berrang-Ford, Cesar Carcamo, Alejandro Llanos, Shuaib Lwasa, Didacus Bambahi Namanya

⁶ Rigolet Inuit Community Government, Rigolet, Nunatsiavut, Labrador, Canada

Correspondence: Sherilee Harper, Department of Population Medicine, University of Guelph, Guelph, Ontario, Canada, N1G 2W1, Email: harpers@uoguelph.ca, Tel: 519-824-4120 ext. 54595

Running Title: Inuit Burden of Acute Gastrointestinal Illness

Key Words: Aboriginal health, acute gastrointestinal illness, burden of illness, foodborne disease, Inuit, waterborne disease

Competing financial interests declaration: None

Abbreviations: Acute Gastrointestinal Illness (AGI)

DOI: <https://doi.org/10.1017/S0950268814003744>

SUMMARY

Food- and waterborne disease is thought to be high in some Canadian Indigenous communities; however, the burden of acute gastrointestinal illness (AGI) is not well understood due to limited availability and quality of surveillance data. This study estimated the burden of community-level self-reported AGI in the Inuit communities of Rigolet, Nunatsiavut, and Iqaluit, Nunavut, Canada. Cross-sectional retrospective surveys captured information on AGI and potential environmental risk factors. Multivariable logistic regression models identified potential AGI risk factors. The annual incidence of AGI ranged from 2.9-3.9 cases/person/year in Rigolet and Iqaluit. In Rigolet, increased spending on obtaining country foods, a homeless person in the house, not visiting a cabin recently, exposure to puppies, and alternative sources of drinking water were associated with increased odds of AGI. In Iqaluit, eating country fish often, exposure to cats, employment status of the person responsible for food preparation, not washing the countertop with soap after preparing meat, a homeless person in the house, and over-crowding were associated with increased odds of AGI. The results highlight the need for systematic data collection to better understand and support previously anecdotal indications of high AGI incidence, as well as insights into unique AGI environmental risk factors in Indigenous populations.

Key Words: Aboriginal health, acute gastrointestinal illness, burden of illness, foodborne disease, Inuit, waterborne disease

INTRODUCTION

Diarrheal disease continues to be a global public health priority [1], especially in the context of climate change, global environmental change, and the globalization of travel and trade [2]. Endemic levels and outbreaks of foodborne, waterborne, and person-to-person transmission of acute gastrointestinal illness (AGI) contribute to considerable morbidity, mortality, and economic costs in developed countries [3–7], and could be particularly important to understand among high risk populations [3–5]. Therefore, environmental and public health practitioners continue to prioritize the surveillance of AGI to monitor the burden of illness, detect and control outbreaks, and evaluate control measures. Under-reporting of AGI is a limitation of surveillance data, especially when used to estimate the burden of illness [8]. Considering this high level of under-reporting, burden of illness studies that estimate rates and identify risk factors for AGI at the community-level are employed internationally [4, 6, 7, 9–12].

While research on the burden of AGI is burgeoning, there is still a limited understanding of enteric illness in sub-sets of these populations, including populations who could be the most vulnerable to enteric illness [13–16]. For instance, although high quality data are not currently available to accurately estimate the burden of AGI in Indigenous communities, research has documented environmental conditions that could increase the risk of AGI (e.g., over-crowding, limited accessibility and availability of safe drinking water) and some researchers have hypothesized that waterborne and foodborne disease is likely disproportionately higher in many Indigenous communities [17–20]. Indeed, some Indigenous populations in Canada, USA, and Australia live in substandard living conditions with more limited access to the quality and quantity of services and resources than other non-Indigenous citizens [19]. These conditions

contribute to disparities in several health outcomes between Indigenous people and non-Indigenous people living in the same country [19, 21]. In addition, effective and timely public health surveillance for Indigenous populations can be challenging, particularly in rural or remote communities due to limited accessibility of healthcare services [19, 22]. This lack of access to these services can result in reduced care-seeking behaviour, which can compromise the quality and quantity of available surveillance data [19, 22–24].

Considering the possible increased vulnerability to AGI and the limitations of surveillance data in some Indigenous communities, there is a clear need for burden of AGI studies to estimate the incidence of, and identify potential risk factors for, AGI at the community-level in these populations. In this context, the goal of this study was to investigate the burden of AGI in the Inuit communities of Rigolet, Nunatsiavut and Iqaluit, Nunavut, Canada. The specific objectives were to estimate the incidence and prevalence, and identify potential socio-economic and environmental risk factors of AGI in each community in the fall (September) and spring (May) 2012-13. The results of this study are important to identify local risk factors that can point to possible intervention strategies, inform decision-making, and better understand the burden of AGI in the North.

METHODS

Study Locations

The term Indigenous is an all-encompassing term that refers to Indigenous inhabitants of Canada, including Inuit, First Nations, and Métis. Many Inuit live in Northern Canada in one of four Inuit regions: Nunatsiavut, Nunavut, Nunavik and Inuvialuit. Inuit have developed a culture and

lifestyle that is dependent upon cold climatic conditions, with the extensive ice and snow coverage providing transportation opportunities between communities, to hunting grounds, and to important cultural activities that are essential for health and wellbeing [25–27]. Canadian Inuit also face challenges regarding access to health services, low socio-economic status compared to the national average, crowded and poor-quality housing, and concerns regarding basic services such as drinking water quality and sanitation [19].

This study took place in two Inuit communities: Iqaluit, Nunavut and Rigolet, Nunatsiavut, Canada. Nunavut (“Our Land”) is Canada’s newest Territory spanning nearly two million square kilometres, with 26 communities and a population of over 31,000 residents (Figure 1). Iqaluit is the capital of Nunavut with a population size of 6,699 people, 62% of whom identify as Aboriginal [28]. Nunatsiavut (“Our Beautiful Land”) is Canada’s most eastern Inuit region, with five remote communities and a growing population of 2,617 [28]. The community of Rigolet is the most southern of five communities in Nunatsiavut (Figure 1), with a population size of 306, 94% of whom identify as Aboriginal [28].

Study Design and Data Collection Tool

The National Studies on Acute Gastrointestinal Illness (NSAGI) study methodology used in Canada [29] was adapted to reflect Inuit culture and the Northern context. Two retrospective cross-sectional surveys were conducted in person in the fall and spring season. The fall survey took place from September 17 – October 3, 2012 in Rigolet, and from September 15 – October 5, 2012 in Iqaluit. The spring survey took place from May 22 – June 19, 2013 in Rigolet, and from May 18 – June 2, 2013 in Iqaluit. Questionnaires previously used in the Canadian burden of

AGI surveys [3, 5, 30], were modified for use in Rigolet and Iqaluit (questionnaires available from the corresponding author upon request). The primary outcome measure was self-reported AGI, which was captured by asking participants if they had experienced a new case of diarrhea and/or vomiting in the past 14 and 28 days. The case definition used for AGI was self-reported vomiting and/or diarrhea (any loose stool) in the past 14 days (September and May) as well as the past 28 days (May), excluding cases who reported vomiting or diarrhea due to pregnancy, medication use, alcohol/drug use, or diagnosed chronic conditions (e.g., colitis, diverticulitis, Crohn's disease, irritable bowel syndrome, *H. pylori*, or other diagnosed chronic conditions) [30]. In an attempt to capture incident cases, if the date of reported AGI symptom onset was prior to the recall period, the case was excluded. Data on potential risk factors were also gathered (Figure 2) [3, 5]. The questionnaire was extensively pre-tested for content and context by academics, health practitioners, and Inuit community members.

Sample Size, Sampling Framework, and Survey Administration

Rigolet, Nunatsiavut:

Given the small population size of Rigolet (n=306), a census sample was attempted; every individual in every household who was in the community during the sampling period was invited to participate. The private and confidential questionnaire was administered by trained community personnel using iPads[®] in the language preferred by the participant, including English and Inuttitut; however, all participants chose English, which is their first language [28].

Iqaluit, Nunavut:

A target sample size of 498 participants was calculated using EpiInfo (Centers for Disease Control and Prevention, Atlanta, Georgia, USA, 2000), to detect an expected 14-day period prevalence of 6% [3, 5]. with a 2% allowable error and a 95% confidence level in a population of 6,184. A two-stage random sample approach was employed. The first stage involved proportionally, randomly selecting blocks of households in Iqaluit and a census of the block was attempted. The second stage involved randomly selecting a household member to participate in the survey using the last birthday-method [31], and all ages were eligible to participate. Participants had the option of conducting the interview in-person or scheduling a telephone interview. Contact with each house was attempted twice during different times of the day. The private and confidential questionnaire was administered by trained community personnel using iPads[®] in the language preferred by the participant, including English, French, and Inuktitut.

Consent and Ethics

Each participant was invited to watch an informational video and written (Rigolet) or oral (Iqaluit) informed consent was obtained. Participants under the age of 18 required parental permission. For participants under the age of 12, the parent could act as a proxy respondent for the child. Compensation for the participant's time and information was provided following the recommendations of local project partners [32, 33]. The study protocol was approved by the Research Ethics Board at the University of Guelph and McGill University, Health Canada Research Ethics Board, and the Nunatsiavut Government Research Advisory Committee. A research license was obtained from Nunavut Research Institute under the Nunavut Scientist Act.

Data Analysis

Data were exported from the iPad[®] iSurveySoft[®] software into Excel[®] (version 14.2.5) and all analyses were conducted using Stata IC (version 11.2), using a significance level of $\alpha= 0.05$. Data from participants responding “unsure” or “refused to answer” were excluded from the analysis of that question. To examine the representativeness of the data, the sample population demographics were compared to census demographics [28] using Pearson χ^2 tests. The annual incidence rate and annual incidence proportion were estimated using equations 1 and 2, respectively [11].

eq. 1: Annual incidence rate

$$= \frac{\text{cases}}{\frac{1}{2}[(\text{total at risk})+(\text{total at risk} - \text{cases})]} * \frac{365}{14 \text{ day recall period}}$$

eq. 2: Annual incidence proportion

$$= 1 - (1 - x)^{(365/14 \text{ day recall period})}, \text{ where } x = (\text{cases})/(\text{total at risk})$$

Multivariable logistic models were built to examine potential risk factors for AGI, with a null hypothesis of no association between AGI (based on 14-day period prevalence) and potential risk factor variables of interest. There were four models; one for September and one for May in each community. For each model, a causal diagram was created to explore potential risk factors of interest based on peer-reviewed literature and biological plausibility. The assumption of linearity was assessed graphically by plotting the continuous variables against the log-odds of the outcome using lowess curves (i.e. locally weighted regression for smoothed scatterplots). To avoid collinearity issues, the correlation between predictors variables was assessed using Spearman rank correlation analysis, using a cut-point value of 70%. If the correlation was above 70%, the most biologically plausible variable was used in the model building process.

Since a census was attempted in Rigolet, a mixed multivariable logistic regression model with a random intercept to control for clustering by household was used for these data. To build the Rigolet models, a series of univariable (i.e., one fixed effect) logistic regression models with a random intercept for household were conducted to explore potential unconditional associations between predictor variables and the outcome variable. Then, a manual backward stepwise approach was used to build a mixed multivariable logistic regression model, considering all predictor variables with $p < 0.20$ in univariable analyses, with a significance level of $\alpha = 0.05$ for risk factor variables to stay in the model. Full and reduced models were compared at each step using likelihood ratio tests; confounding was assessed at each step and if adding or removing a variable resulted in a 30% change in the model β -coefficients, the variable was considered a confounder and remained in the model regardless of statistical significance. All two-way interactions amongst predictors with $p < 0.10$ in the univariable analyses were assessed. Finally, the fit of the models were assessed via graphical examination of Pearson residuals, and assessing the assumptions of normality and homogeneity of variance for the best linear unbiased predictors (BLUPs).

For the Iqaluit data, multivariable exact logistic regression models were built since one individual per household was randomly selected and, thus, there was no household level clustering for these data. Specifically, a series of univariable exact logistic regressions using conditional scores tests were conducted to explore potential unconditional associations between predictor variables and the outcome variable. A manual backward stepwise approach was used to build the multivariable models, considering all predictor variables with $p < 0.20$ in univariable

analyses, with a significance level of $\alpha = 0.05$ for risk factor variables to stay in the model regardless of statistical significance. Confounding was assessed at each step and if adding or removing a variable resulted in a 30% change in the model β -coefficients, the variable was considered a confounder and remained in the model. Finally, all two-way interactions amongst predictors with $p < 0.10$ in the univariable analyses were assessed. For Iqaluit and Rigolet models, if sex and age were not significant in the final model, we forced these variables into the model to examine any changes in predictor variable coefficients and confidence intervals; if little change was observed (i.e. less than 30%), the age and/or sex variable was not included the final model.

RESULTS

In Rigolet, a total of 245 people were in the community during the September study period, and 226 questionnaires were completed (response rate = 92%); a total of 249 were in the community during the May study period and 236 questionnaires were completed (response rate = 95%). In Iqaluit, a total of 532 questionnaires were completed in September (response rate = 75%) and a total of 523 questionnaires were completed in May (response rate = 55%). Each questionnaire took an average of 14 minutes to complete.

The demographics of September and May survey participants compared to the 2011 census of Rigolet and Iqaluit are shown in Table 1. Compared to the census demographics of Iqaluit, females, older people, and Indigenous people were over-represented in the September survey ($p < 0.05$), and females and older people were over-represented in the May survey ($p < 0.05$). In

Rigolet, there were no significant differences in sex, age, and Indigenous identity between survey participants and the 2011 census ($p>0.05$).

AGI Prevalence, Incidence, and Risk Factors

In Rigolet, the estimated annual incidence rate was 3.8 and 3.9 episodes per person per year in September and May, respectively (Table 2). In Iqaluit, the estimated annual incidence rate was 3.8 and 2.8 episodes per person per year in September and May, respectively. The 14-day and 28-day prevalence, estimated annual incidence, and estimated annual incidence proportion for Rigolet and Iqaluit are presented in Table 2. In the September final multivariable model for Rigolet, not visiting a cabin in the past month, increased spending on obtaining country food (i.e. locally harvest food including berries, duck eggs, fish, caribou, marine mammals, ducks), and a homeless person staying in the home (as defined by the Inuit Health Survey [34]) increased the odds of AGI (Table 3). Final model results indicated no significant difference between models that did and did not account for clustering at the household-level (Table 3); however, a random intercept was forced in the model to control for household-level clustering due to the structure of the Rigolet data, as well as similar variable coefficients, variable confidence intervals, and model fit (based on Akaike information criterion) for models that did and did not control for clustering. In the May final multivariable model for Rigolet, age group (0-19, 20-55, over 55), exposure to a puppy in the past month, and consumption of tap water alternatives (e.g. not drinking tap water) in the past two weeks were significantly associated with increased odds of AGI (Table 4). Final model results indicated a significant difference between models that did and did not account for clustering at the household-level; as such, the final mixed multivariable model is presented in Table 4.

In the September final multivariable model for Iqaluit, monthly frequency of eating country fish (i.e. locally harvested fish including salmon, char, trout), washing kitchen counter-tops and cutting boards with soap after preparing meat, employment status of the person responsible for food preparation, and cat exposure in the past month significantly affected the odds of AGI (Table 5). Finally, in the May final multivariable model for Iqaluit, a homeless person staying in the home, as well as an interaction between pet ownership and over-crowding affected the odds of AGI (Table 6).

DISCUSSION

The estimated annual incidence rates of AGI in Rigolet and Iqaluit were higher than those reported in other studies using similar methodology in Canada [3–5, 29, 30], New Zealand [35], the Netherlands [36], Italy [10], Australia [37], Malta [38], Cuba [39], Poland [9], Denmark [40], Argentina [12], USA [7], and Hong Kong [41] (Table 2). It is important to note, however, that we sampled during potentially high-risk AGI seasons, which could result in over-estimates. Nonetheless, the estimated annual incidence in the two communities was higher than that observed in Chile during high risk seasons, using a similar design, case definition, and recall period [11].

Hunting, trapping, fishing, gathering, and sharing country food are still practiced in Rigolet and Iqaluit, and are important aspects of Inuit livelihoods, health, and wellbeing [27]. In Rigolet and Iqaluit, some factors related to country food (consumption of country fish and amount of money spent obtaining country foods) increased the odds of AGI in the statistical model. Other research

has suggested that some country foods and country food preparation in the North could increase the risk of AGI [42, 43]. The routes of contamination, as well as preparation methods for country meat in the North are different than retail meats available for purchase, and these transmission pathways are not well understood. Nonetheless, in this study it was not possible to determine whether country food-related factors were direct contributors to AGI, or if these variables were proxy indicators for other risk factors. For instance, in Rigolet, households who spend more money on obtaining country foods might also spend more time hunting, trapping, fishing, and gathering foods, and some research has linked AGI with reduced hygiene practices during camping [44]. More targeted research should explore more specifically how country food practices and consumption may affect exposure to, and infection by, agents of AGI in the Arctic, especially in the context of climatic change.

In Rigolet, not visiting a cabin recently significantly increased the odds of AGI in the statistical models, which is likely a proxy indicator for a related risk factor (e.g. socio-economic status as discussed below). For many Inuit communities, visiting cabins includes permanent or semi-permanent structures (e.g. small buildings that can be disassembled and relocated, as well as tents, snow-houses) that are located on land typically not owned by an individual and not accessible by road and thus reached by boat or snowmobile. Factors that impact on an individual's access to a cabin include having someone in the home that has the skills to navigate the land and ice, weather conditions (as well as climate changes), fuel and maintenance costs for boats or snowmobiles, and geographic location of the cabin. Past research in the Arctic has documented the importance of going to cabins to connect with the land as critical to good physical and emotional health [26, 27]. Several studies have described a relationship between stress and

decreased gastrointestinal health [45, 46], which could perhaps also extend to AGI outcomes. Due to the cross-sectional nature of our study, however, we cannot assess the temporality of this statistically significant finding, nor did we attempt to measure stress in participants. It is possible that people who visit a cabin more frequently have lower levels of stress and thus reduced susceptibility to AGI outcomes; however, this finding could also suggest that those who had AGI were not able to visit the cabin because of their illness. Therefore, more research examining the physical health and wellbeing benefits of visiting cabins and connecting with the land is warranted and would be valuable for Indigenous public health planning and programming globally.

In Rigolet and Iqaluit, exposure to cats and puppies was statistically associated with AGI outcomes, which was also reported in a similar study in Canada [3–5, 29, 30] and Chile [11]. Cats and dogs are known to sometimes asymptotically carry several pathogens that could cause AGI symptoms in humans [47–49]. We did not test cats or dogs in this study for fecal shedding of AGI pathogens, nor were data on cat and dog exposure captured in other burden of AGI studies in North America [3, 4, 7, 29, 50]. Therefore, we do not know whether cats and dogs were actually involved in exposure of participants of this study to agents of AGI, or if this exposure factor is a surrogate for other risk factors. Further research should examine the potential for cats, dogs, and other domestic pets to be reservoirs of agents of human AGI-related infection in Inuit communities, but also in North America in general.

In Iqaluit, not washing the kitchen countertop with soap after preparing retail or country meat increased the odds of AGI. This finding is consistent with past research that identified kitchen

countertops as an important source of foodborne illness [51, 52], which suggests that washing the counter top after preparing meat reduces the risk of exposure to foodborne pathogens [51, 53]. Not washing the countertop with soap might be explained by the transition in diet and food handling practices in the Arctic. A rapid shift in diet from predominantly eating country foods (e.g., caribou, seal, walrus, whale) to food purchased at retail stores (e.g., chicken, beef, pork) has been documented in most Inuit communities [54]. This transition in diet also requires a shift in food preparation and handling techniques. For example, in Iqaluit, country meats are commonly prepared on pieces of cardboard on the floor and consumed raw or frozen; after which, the cardboard is disposed of rather than washing and reusing it. Conversely, chicken, beef, and pork is prepared on the countertop and eaten cooked. Further research is required to adequately understand the knowledge, attitudes, and behaviours regarding country and retail food handling and safety in the North to develop and implement effective public health planning and programming.

We also found that the odds of AGI significantly increased if the person responsible for food preparation in the house was employed (either part-time or full-time), which could be explained by employment limiting time available for proper food safety practices, higher income leading to more risky food consumption behaviours [55–57], or employment limiting time available to access the land for country food resulting in increased retail food consumption. Past research suggests that those with higher levels of income are less likely to wash countertops after preparing meat and practice other good food preparation techniques in the kitchen than those with lower levels of education and income [55–57]. These studies posit that those with higher levels of income are more likely to have higher knowledge regarding safe food handling

practices and behaviours, but still partake in the risky behaviour. The reasons for these trends might be due to cultural influences or social norms [55], but are not clear and warrant future research.

Several Inuit communities face water-related challenges in terms of the provision of safe municipal drinking water [58], effects of climate change and high impact weather events on drinking water safety [20, 59], and the preference of many residents to seek alternative drinking water sources including brooks, streams, ice, and snow [58]. In using these alternative sources of drinking water, residents could be at higher risk for AGI, which could explain why drinking tap water in Rigolet was associated with significantly reduced odds of AGI. Alternatively, residents in Rigolet have reported perceived serious concerns about the safety of tap water [60]; thus, this finding could be explained by AGI cases perceiving tap water as the source of their illness, and thus choosing not to drink tap water after being ill. Further research should specifically investigate the fraction of AGI attributable to drinking water, as well as the temporal relationship between weather patterns, water quality, and AGI in Northern communities.

As discussed by other researchers, this study had several limitations that are not uncommon among burden of AGI studies [6, 11, 12, 61]. First, this study was done in only two Inuit communities, and Inuit communities are very heterogeneous in culture, language, mobility, and geographical locations. While we attempted to sample two communities to capture information from a small and large Inuit settlement, urban and remote setting, and from two Inuit regions, caution should be exercised in generalizing the results to the Nunatsiavut region, Nunavut Territory, or Inuit communities in general. Second, this study was conducted at two points in

time during the year, which could result in an over or under-estimate of the annual incidence of AGI. Collecting data longitudinally over an entire year or multiple years would enable more accurate estimates of AGI incidence. Third, while we excluded cases of AGI that were due to diagnosed chronic conditions, those cases with undiagnosed AGI-related chronic conditions might have been misclassified as a case. Fourth, considering the structure of the data and environmental modes of AGI transmission, in Rigolet we were surprised to find no significant clustering at the household level in the September survey; this warrants further research. Fifth, the outcome measure in this study used a self-reported case definition; pathogen testing was beyond the scope of this study. Our case definition was non-specific; however, we chose this outcome measure and case definition to match other Canadian studies to facilitate comparisons and inform decision making processes that are based on national surveillance systems, as well as to capture community-level burden of illness (compared to burdens estimated from surveillance data). There are biases inherent in self-reported case definitions, including recall bias, interviewer bias, and misclassification bias. We have assumed that these biases are similar between Rigolet and Iqaluit, and with other burden of AGI studies using similar methods. Finally, while we attempted a census sample in Rigolet, the sample size was still relatively small for statistical modelling. The high response rate and no significant differences between demographics between the survey population and the Canadian census suggests a representative sample, which is unique and an important contribution as representative samples are not common in Indigenous health research, and can be challenging to achieve in many Indigenous communities [62].

CONCLUSION

This study estimated AGI incidence at the community level in Rigolet and Iqaluit, and found a higher estimated incidence of AGI in these communities compared to other similar studies in Canada and abroad. Several factors were associated with increased odds of AGI, including food sources, water sources, animal exposure, over-crowding, and food safety practices. Information on this increased incidence of AGI and these potential AGI risk factors offers important information for public health planning, prioritization, and programming in Inuit regions. While this research estimates the incidence and potential risk factors of AGI in Rigolet and Iqaluit, the improved understanding of AGI experienced at the community-level in a Canadian Indigenous community sheds light on the need to better understand the burden of AGI in sub-sets of the population that might be at higher risk of AGI, including Indigenous populations.

ACKNOWLEDGEMENTS:

First and foremost, we thank the residents in Iqaluit and Rigolet for welcoming us into their community, engaging with us throughout the project, and contributing to the larger IHACC research project. We are thankful to all the surveyors in Rigolet (M. Baikie, I. Shiwak, and D. Wolfrey) and Iqaluit (M. Naglingniq, S. Kownirk, G. Akpik, M. Atagoyuk, N. Atagoyuk, T. Atagoyuk, J. Campbell, N. Campbell, P. Davidee, A. Kilabuk-Degrassé, N. Lewis, S. Lewis, L. Mark, D. McGlade, E. Nevin, O. Nowdlak, I. Oopakak, M. Petooloosie, M. Pillaktuaq, M. Unahah). This survey would not have been possible without their diligent work and expertise. We would like to thank M. Baikie, P. Workman, K. Hutchinson, W. Joy, and many others in the Department of Health and Social Services, Government of Nunavut; J. Shirley and M.E. Thomas and others at the Nunavut Research Institute; M. Wood, G. Turner, M. Kinney, T. Buckle and others at the Department of Health and Social Development, Nunatsiavut Government; M. LeBlanc-Havard, C. Brice-Bennett, G. Elliott and others at the Labrador Grenfell Health Authority; C. Wolfrey, S. Blake, S. Wolfrey, and M. Williams at the Rigolet Inuit Community Government; and S. Maclachlan, N. Markwick, C. Huet, J. Ostapchuk, M.E. MacDonald, Y. Guo, and J. Fraser at the University of Guelph. Thanks to S. Majowicz for her assistance and burden of AGI expertise. Particular thanks to A. Bunce and P. Arpin for their dedication throughout the project. These people played instrumental roles in contributing to the study design, data collection, data interpretation, and ultimately the success of this project. This research is part of an international project entitled the “Indigenous Health Adaptation to Climate Change” project, with parallel field study sites in Uganda and Peru. The study design and conduct were independent of the funding source. This work was supported by the Public Health Agency of Canada; the Nasivvik Centre for Inuit Health and Changing Environments; the

International Research Initiative on Adaptation to Climate Change (IRIACC) funded by the International Development Research Centre (IDRC), Canadian Institutes for Health Research (CIHR), Social Sciences and Humanities Research Council (SSHRC), and National Science and Engineering Research Council (NSERC); and a Vanier Canada Graduate Scholarship to S. Harper (CIHR).

REFERENCES

1. WHO. Global Health Risk Summary Tables. Geneva, Switzerland: Health Statistics and Informatics Department, World Health Organization; 2009.
2. Costello A, Abbas M, Allen A, Ball S, Bell S, Bellamy R, et al. Managing the Health Effects of Climate Change. *Lancet*. 2009;373(9676):1693–733.
3. Majowicz SE, Doré K, Flint J, Edge VL, Read S, Buffett MC, et al. Magnitude and Distribution of Acute, Self-Reported Gastrointestinal Illness in a Canadian Community. *Epidemiology and Infection*. 2004;132(4):607–17.
4. Sargeant JM, Majowicz SE, Snelgrove J. The Burden of Acute Gastrointestinal Illness in Ontario, Canada, 2005-2006. *Epidemiology and Infection*. 2008;136(4):451–60.
5. Thomas MK, Majowicz SE, MacDougall L, Sockett PN, Kovacs SJ, Fyfe M, et al. Population Distribution and Burden of Acute Gastrointestinal Illness in British Columbia, Canada. *BMC Public Health*. 2006;6:307.
6. Scallan E, Majowicz SE, Hall G, Banerjee A, Bowman CL, Daly L, et al. Prevalence of Diarrhoea in the Community in Australia, Canada, Ireland, and the United States. *International Journal of Epidemiology*. 2005;34(2):454–60.
7. Jones TF, McMillian MB, Scallan E, Frenzen PD, Cronquist AB, Thomas S, et al. A Population-Based Estimate of the Substantial Burden of Diarrhoeal Disease in the United States; FoodNet, 1996-2003. *Epidemiology and Infection*. 2007;135(2):293–301.
8. MacDougall L, Majowicz S, Doré K, Flint J, Thomas K, Kovacs S, et al. Under-Reporting of Infectious Gastrointestinal Illness in British Columbia, Canada: Who Is Counted in Provincial Communicable Disease Statistics? *Epidemiology and Infection*. 2008;136(2):248–56.
9. Baumann-Popczyk A, Sadkowska-Todys M, Rogalska J, Stefanoff P. Incidence of Self-Reported Acute Gastrointestinal Infections in the Community in Poland: A Population-Based Study. *Epidemiology and Infection*. 2012;140(7):1173–84.

10. Scavia G, Baldinelli F, Busani L, Caprioli A. The Burden of Self-Reported Acute Gastrointestinal Illness in Italy: A Retrospective Survey, 2008-2009. *Epidemiology and Infection*. 2012;140(7):1193–206.
11. Thomas MK, Perez E, Majowicz SE, Reid-Smith R, Olea A, Diaz J, et al. Burden of Acute Gastrointestinal Illness in the Metropolitan Region, Chile, 2008. *Epidemiology and Infection*. 2011;139(4):560–71.
12. Thomas MK, Perez E, Majowicz SE, Reid-Smith R, Albil S, Monteverde M, et al. Burden of Acute Gastrointestinal Illness in Gálvez, Argentina, 2007. *Journal of Health, Population, and Nutrition*. 2010;28(2):149–58.
13. Cheng AAC, McDonald JR, Thielman NM. Infectious Diarrhea in Developed and Developing Countries. *Journal of Clinical Gastroenterology*. 2005;39(9):757–73.
14. Pardhan-Ali A, Wilson J, Edge VL, Furgal C, Reid-Smith R, Santos M, et al. Community-Level Risk Factors for Notifiable Gastrointestinal Illness in the Northwest Territories, Canada, 1991-2008. *BMC Public Health*. 2013;13(1):63.
15. Pardhan-Ali A, Berke O, Wilson J, Edge VL, Furgal C, Reid-Smith R, et al. A Spatial and Temporal Analysis of Notifiable Gastrointestinal Illness in the Northwest Territories, Canada, 1991-2008. *International Journal of Health Geographics*. 2012;11(1):17.
16. Pardhan-Ali A, Wilson J, Edge VL, Furgal C, Reid-Smith R, Santos M, et al. A Descriptive Analysis of Notifiable Gastrointestinal Illness in the Northwest Territories, Canada, 1991-2008. *BMJ Open*. 2012;2(4).
17. Hruddy SE, Hébert PC, Stanbrook MB, Sibbald B, Flegel K, MacDonald N, et al. Safe Water? Depends on Where You Live! *Canadian Medical Association Journal*. 2008;178(8):975.
18. Hruddy SE, Hruddy EJ. Published Case Studies of Waterborne Disease Outbreaks—Evidence of a Recurrent Threat. *Water Environment Research*. 2007;79(3):233–45.
19. Adelson N. The Embodiment of Inequity: Health Disparities in Aboriginal Canada. *Canadian Journal of Public Health*. 2005;96:S45–61.
20. Harper SL, Edge VL, Schuster-Wallace CJ, Berke O, McEwen SA. Weather, Water Quality and Infectious Gastrointestinal Illness in Two Inuit Communities in Nunatsiavut, Canada: Potential Implications for Climate Change. *EcoHealth*. 2011;8(1):93–108.
21. King M, Smith A, Gracey M. Indigenous Health Part 2: The Underlying Causes of the Health Gap. *Lancet*. 2009;374(9683):76–85.

22. Smylie J, Anderson M. Understanding the Health of Indigenous Peoples in Canada: Key Methodological and Conceptual Challenges. *Canadian Medical Association Journal*. 2006;175(6):602–5.
23. Auer A, Andersson R. Canadian Aboriginal Communities and Medical Service Patterns for the Management of Injured Patients: A Basis for Surveillance. *Public Health*. 2001;115:44–50.
24. Harper SL, Edge VL, Schuster-Wallace CJ, Ar-Rushdi M, McEwen SA. Improving Aboriginal Health Data Capture: Evidence from a Health Registry Evaluation. *Epidemiology and Infection*. 2011;139(11):1774–83.
25. Cunsolo Willox A, Harper SL, Ford JD, Edge VL, Landman K, Houle K, et al. Climate Change and Mental Health: An Exploratory Case Study from Rigolet, Nunatsiavut, Canada. *Climatic Change*. 2013;121(2):255–70.
26. Cunsolo Willox A, Harper SL, Ford JD, Landman K, Houle K, Edge VL, et al. “From This Place and of This Place:” Climate Change, Sense of Place, and Health in Nunatsiavut, Canada. *Social Science & Medicine*. 2012;75(3):538–47.
27. Cunsolo Willox A, Harper SL, Edge VL, Landman K, Houle K, Ford JD, et al. The Land Enriches the Soul: On Climatic and Environmental Change, Affect, and Emotional Health and Well-Being in Rigolet, Nunatsiavut, Canada. *Emotion, Space and Society*. 2013;6:14–24.
28. Statistics Canada. Rigolet, Newfoundland and Labrador. Census Profile. 2011 Census. Statistics Canada Catalogue No. 98-316-XWE. Ottawa. Released October 24, 2012. [Internet]. 2012.
29. Thomas MK, Majowicz SE, Pollari F, Sockett PN. Burden of Acute Gastrointestinal Illness in Canada, 1999-2007: Interim Summary of NSAGI Activities. *Canada Communicable Disease Report*. 2008;34(4):4–9.
30. Majowicz SE, Horrocks J, Bocking K. Demographic Determinants of Acute Gastrointestinal Illness in Canada: A Population Study. *BMC Public Health*. 2007;7:162.
31. Gaziano C. Comparative Analysis of within-Household Respondent Selection Techniques. *Public Opinion Quarterly*. 2005;69(1):124–57.
32. Research CI of H. CIHR Guidelines for Health Research Involving Aboriginal People. 2007;
33. Pearce TD, Ford JD, Laidler GJ, Smit B, Duerden F, Allarut M, et al. Community Collaboration and Climate Change Research in the Canadian Arctic. *Polar Research*. 2009;28(1):10–27.

34. Saudny H, Leggee D, Egeland G. Design and Methods of the Adult Inuit Health Survey 2007-2008. *International Journal of Circumpolar Health*. 2012;71:1–9.
35. Adlam SB, Perera S, Lake RJ, Campbell DM, Williman J, Baker MG. Acute Gastrointestinal Illness in New Zealand: A Community Study. *Epidemiology and Infection*. 2011;139(2):302–8.
36. Doorduyn Y, Van Pelt W, Havelaar H, Havelaar AH. The Burden of Infectious Intestinal Disease (IID) in the Community: A Survey of Self-Reported IID in The Netherlands. *Epidemiology and Infection*. 2012;140(7):1185–92.
37. Hall G, Kirk M, Ashbolt R, Stafford R, Lalor K, Health P, et al. Frequency of Infectious Gastrointestinal Illness in Australia, 2002: Regional, Seasonal and Demographic Variation. *Epidemiology and Infection*. 2006;134(1):111–8.
38. Gauci C, Gilles H, O'brien S, Mamo J, Stabile I, Ruggeri FM, et al. The Magnitude and Distribution of Infectious Intestinal Disease in Malta: A Population-Based Study. *Epidemiology and Infection*. 2007;135(8):1282–9.
39. Prieto PA, Finley RL, Muchaal PK, Guerin MT, Isaacs S, Domínguez AC, et al. Burden of Self-Reported Acute Gastrointestinal Illness in Cuba Study Design. *Journal of Health, Population, and Nutrition*. 2009;27(3):345–57.
40. Müller L, Korsgaard H, Ethelberg S. Burden of Acute Gastrointestinal Illness in Denmark 2009: A Population-Based Telephone Survey. *Epidemiology and Infection*. 2012;140(2):290–8.
41. Ho SC, Chau PH, Fung PK, Sham A, Nelson E, Sung J. Acute Gastroenteritis in Hong Kong: A Population-Based Telephone Survey. *Epidemiology and Infection*. 2010;138(7):982–91.
42. Parkinson AJ, Butler JC. Potential Impacts of Climate Change on Infectious Diseases in the Arctic. *International Journal of Circumpolar Health*. 2005;64(5):478–86.
43. Pufall EL, Jones AQ, McEwen SA, Lyall C, Peregrine AS, Edge VL. Perception of the Importance of Traditional Country Foods to the Physical, Mental, and Spiritual Health of Labrador Inuit. *Arctic*. 2011;64(2):242–50.
44. Boulware DR. Influence of Hygiene on Gastrointestinal Illness among Wilderness Backpackers. *Journal of Travel Medicine*. 2004;11(1):27–33.
45. Drossman D, Creed FH, Olden KW, Svedlund J, Toner BB, Whitehead WE. Psychosocial Aspects of the Functional Gastrointestinal Disorders. *Gut*. 1996;25(1):21–34.
46. Bhatia V, Tandon R. Stress and the Gastrointestinal Tract. *Journal of Gastroenterology and Hepatology*. 2005;20:332–9.

47. Procter TD, Pearl DL, Finley RL, Leonard EK, Janecko N, Reid-Smith RJ, et al. A Cross-Sectional Study Examining Campylobacter and Other Zoonotic Enteric Pathogens in Dogs That Frequent Dog Parks in Three Cities in South-Western Ontario and Risk Factors for Shedding of Campylobacter Spp. *Zoonoses and Public Health*. 2014;61(3):208–18.
48. Leonard EK, Pearl DL, Finley RL, Janecko N, Reid-Smith RJ, Peregrine AS, et al. Comparison of Antimicrobial Resistance Patterns of Salmonella Spp. and Escherichia Coli Recovered from Pet Dogs from Volunteer Households in Ontario (2005-06). *The Journal of Antimicrobial Chemotherapy*. 2012;67(1):174–81.
49. Lefebvre SL, Reid-Smith RJ, Waltner-Toews D, Weese JS. Incidence of Acquisition of Methicillin-Resistant Staphylococcus Aureus, Clostridium Difficile, and Other Health-Care-Associated Pathogens by Dogs That Participate in Animal-Assisted Interventions. *Journal of the American Veterinary Medical Association*. 2009;234(11):1404–17.
50. Febriani Y, Levallois P, Gingras S, Gosselin P, Majowicz SE, Fleury MD. The Association between Farming Activities, Precipitation, and the Risk of Acute Gastrointestinal Illness in Rural Municipalities of Quebec, Canada: A Cross-Sectional Study. *BMC Public Health*. 2010;10:48.
51. Gorman R, Bloomfield S, Adley CC. A Study of Cross-Contamination of Food-Borne Pathogens in the Domestic Kitchen in the Republic of Ireland. *International Journal of Food Microbiology*. 2002;76(1-2):143–50.
52. Kendall P, Elsbernd A, Sinclair K, Schroeder M, Chen G, Bergmann V, et al. Observation versus Self-Report: Validation of a Consumer Food Behavior Questionnaire. *Journal of Food Protection*. 2004;67(11):2578–86.
53. Medrano-Félix A, Martínez C, Castro-del Campo N, León-Félix J, Peraza-Garay F, Gerba CP, et al. Impact of Prescribed Cleaning and Disinfectant Use on Microbial Contamination in the Home. *Journal of Applied Microbiology*. 2011;110(2):463–71.
54. Sharma S, Cao X, Roache C, Buchan A, Reid R, Gittelsohn J. Assessing Dietary Intake in a Population Undergoing a Rapid Transition in Diet and Lifestyle: The Arctic Inuit in Nunavut, Canada. *The British Journal of Nutrition*. 2010;103(5):749–59.
55. Yang S, Leff MG, McTague D, Horvath KA, Jackson-Thompson J, Murayi T, et al. Multistate Surveillance for Food-Handling, Preparation, and Consumption Behaviors Associated with Foodborne Diseases: 1995 and 1996 BRFSS Food-Safety Questions. *Morbidity and Mortality Weekly Report*. 1998;47(SS-4):33–54.
56. Li-Cohen AE, Bruhn CM. Safety of Consumer Handling of Fresh Produce from the Time of Purchase to the Plate: A Comprehensive Consumer Survey. *Journal of Food Protection*. 2002;65(8):1287–96.

57. Altekruise SF, Street DA, Fein SB, Levy AS. Consumer Knowledge of Foodborne Microbial Hazards and Food-Handling Practices. *Journal of Food Protection*. 1996;59(3):287–94.
58. Martin D, Bélanger D, Gosselin P, Brazeau J, Furgal C, Déry S. Drinking Water and Potential Threats to Human Health in Nunavik: Adaptation Strategies under Climate Change Conditions. *Arctic*. 2007;60(2):195–202.
59. White DM, Craig Gerlach S, Loring P, Tidwell AC, Chambers MC. Food and Water Security in a Changing Arctic Climate. *Environmental Research Letters*. 2007;2(4):045018.
60. Goldhar C, Bell T, Wolf J. Rethinking Existing Approaches to Water Security in Remote Communities: An Analysis of Two Drinking Water Systems in Nunatsiavut, Labrador, Canada. *Water Alternatives*. 2013;6(3):462–86.
61. Majowicz SE, Hall G, Scallan E, Adak GK, Gauci C, Jones TF, et al. A Common, Symptom-Based Case Definition for Gastroenteritis. *Epidemiology and Infection*. 2008;136(7):886–94.
62. Ford JD. Indigenous Health and Climate Change. *American Journal of Public Health*. 2012;102(7):1260–6.

TABLES

Table 1. Demographics of Rigolet and Iqaluit based on the 2011 Census, as well as the September 2012 and May 2013 survey respondents in Rigolet, Nunatsiavut, and Iqaluit, Nunavut, Canada in September 2012 and May 2013.

Variable	RIGOLET			IQUALUIT		
	Rigolet 2011 Census No. (%)	September Survey Participants No. (%)	May Survey Participants No. (%)	Iqaluit 2011 Census No. (%)	September Survey Participants No. (%)	May Survey Participants No. (%)
Sex	<i>n</i> = 305	<i>n</i> = 226	<i>n</i> = 236	<i>n</i> = 6,695	<i>n</i> = 532	<i>n</i> = 522
Male	160 (52.5)	109 (48.2)	117 (49.6)	3,400 (50.8)	207 (38.9)*	221 (42.3)*
Female	145 (47.5)	117 (51.8)	119 (50.4)	3,295 (49.2)	325 (61.1)*	302 (57.7)*
Age	<i>n</i> = 305	<i>n</i> = 226	<i>n</i> = 236	<i>n</i> = 6,710	<i>n</i> = 529	<i>n</i> = 522
0-9	40 (13.1)	38 (16.8)	43 (18.2)	1,150 (17.1)	44 (8.2)*	52 (10.0)*
10-14	15 (4.9)	17 (7.5)	19 (8.1)	450 (6.7)	13 (2.5)*	25 (4.8)
15-19	15 (4.9)	10 (4.4)	12 (5.1)	515 (7.7)	20 (3.8)*	21 (4.0)*
20-24	25 (8.2)	10 (4.4)	10 (4.2)	560 (8.2)	32 (6.1)	38 (7.3)
25-64	180 (59.0)	134 (59.3)	139 (58.9)	3,870 (57.7)	382 (72.2)*	364 (69.7)*
65-69	10 (3.3)	6 (2.7)	6 (2.5)	80 (1.2)	17 (3.2)*	9 (1.7)
70+	20 (6.6)	11 (4.9)	7 (3.0)	85 (1.3)	21 (4.0)*	13 (2.5)*
Indigenous	<i>n</i> = 270	<i>n</i> = 226	<i>n</i> = 236	<i>n</i> = 6,085	<i>n</i> = 531	<i>n</i> = 522
Indigenous	250 (92.6)	216 (95.6)	226 (95.8)	3650 (60.0)	362 (68.2)*	332 (63.6)
Non-Indigenous	20 (7.4)	10 (4.4)	10 (4.2)	2435 (40.0)	169 (31.8)*	190 (36.4)

* proportion per category significantly different ($p < 0.05$) from all other categories combined using Pearson chi-square test. For age, we compared each category to all other categories collapsed.

Table 2. Estimates of acute gastrointestinal illness (AGI) estimated incidence in Rigolet, Nunatsiavut and Iqaluit, Nunavut, Canada in September 2012 and May 2013.

AGI Estimated Incidence	RIGOLET		IQALUIT	
	September	May	September	May
<i>Study AGI Case Definition* - 14-day recall</i>				
	n=30	n=32	n=72	n=53
14-day Period Prevalence	13.5% (8.9-18.0%)	13.8% (9.4-18.3%)	13.7% (10.7-16.6%)	10.6% (7.7 – 13.0%)
Estimated Annual Incidence Rate (episodes/person/year)	3.8 (2.4-5.2)	3.9 (2.5-5.3)	3.8 (2.9-4.8)	2.8 (2.1-3.7)
Estimated Age-adjusted Annual Incidence Rate (episodes/person/year)	3.6 (2.3-5.0)	4.1 (2.7-5.6)	4.1 (3.0-4.9)	2.8 (2.1-3.6)
Estimated Annual Incidence Proportion	97.7%	97.9%	97.8%	94.2%
<i>Study AGI Case Definition* - 28-day recall</i>				
		n=47		n=105
28-day Period Prevalence	-	20.5% (15.3-25.8%)	-	20.9% (17.3-24.5%)
Estimated Annual Incidence Rate (episodes/person/year)	-	3.0 (2.1-3.9)	-	3.1 (2.5-3.7)
Estimated Age-adjusted Annual Incidence Rate (episodes/person/year)	-	3.0 (2.1-3.9)	-	3.1 (2.5-3.7)
Estimated Annual Incidence Proportion	-	95.0%	-	95.3%
<i>International AGI Case Definition** - 14-day recall</i>				
	n=17	n=23	n=64	n=43
14-day Period Prevalence	7.6% (4.1-11.1%)	10.0% (6.1-13.8%)	12.1% (9.3-14.9%)	8.4% (6.0-10.8%)
Estimated Annual Incidence Rate (episodes/person/year)	2.2 (1.1-3.3)	2.7 (1.6-4.0)	2.8 (2.0-3.5)	2.3 (1.6-3.0)
Estimated Age-adjusted Annual Incidence Rate (episodes/person/year)	1.9 (0.9-3.0)	3.1 (1.9-4.4)	3.6 (2.7-4.5)	2.2 (1.5-2.9)
Estimated Annual Incidence Proportion	88.9%	93.5%	93.7%	89.8%
<i>International AGI Case Definition** - 28-day recall</i>				
		n=34		n=86
28-day Period Prevalence	-	14.8% (10.2-19.5%)	-	17.1% (13.8-20.4%)
Estimated Annual Incidence Rate (episodes/person/year)	-	2.1 (1.4-2.8)	-	2.4 (1.9-3.0)
Estimated Age-adjusted Annual Incidence Rate (episodes/person/year)	-	3.0 (2.1-3.9)	-	3.1 (2.5-3.7)
Estimated Annual Incidence Proportion	-	87.7%	-	91.4%

*Study AGI Case Definition: Any diarrhea and/or vomiting not due to pregnancy, drugs or alcohol, medication or diagnosed chronic conditions.

** International AGI Case Definition: 3+ diarrhea and/or vomiting not due to pregnancy, drugs or alcohol, medication or diagnosed chronic conditions [61].

- A 28 day recall period was not captured in the September 2012 questionnaire.

Table 3. Univariable exact logistic regression* and final multivariable logistic regression (controlling for household clustering) model [□]* results, examining the effects of predictor variables on the odds of acute gastrointestinal illness in Rigolet, Nunatsiavut, in September 2012.

RIGOLET SEPTEMBER MODEL	n	Univariable Results			Multivariable Model Results		
		Odds Ratio	P	Confidence Interval (95%)	Odds Ratio	P	Confidence Interval (95%)
Homeless person staying in the house							
No	182	<i>referent</i>			<i>referent</i>		
Yes	38	6.17	0.022	1.30 - 29.23	4.76	0.021	1.27 - 17.90
Visited cabin in past month							
No	74	1.94	0.100	0.89 - 4.23	3.33	0.039	1.06 - 10.44
Yes	152	<i>referent</i>			<i>referent</i>		
Weekly amount spent on obtaining country food							
Low (under \$150)	89	<i>referent</i>			<i>referent</i>		
Medium (\$150-\$300)	87	1.68	0.320	0.60 - 4.70	2.65	0.130	0.75 - 9.36
High (over \$300)	35	3.08	0.062	0.94 - 10.06	7.18	0.010	1.62 - 31.90
Storage of drinking water							
Container in the fridge	122	<i>referent</i>			-	-	- -
Container outside of the fridge	51	2.71	0.077	0.90 - 8.20	-	-	- -
No storage	47	3.76	0.028	1.15 - 12.24	-	-	- -
Exposure to cats in past month							
No	159	<i>referent</i>			-	-	- -
Yes	62	1.83	0.200	0.71 - 4.73	-	-	- -
Perceived quality of drinking water							
Very poor or poor	32	2.85	0.157	0.67 - 12.17	-	-	- -
Fair, good, or very good	187	<i>referent</i>			-	-	- -

* The results from the univariable analysis are presented for those variables with $p < 0.20$.

[□]* Likelihood ratio test comparing the model with and without the household-level variable: variance = 0.51, 95% CI = 0.007- 35.138, $p = 0.300$; Intra-class correlation coefficient (ICC) = 0.14, 95% CI = 0.002 – 0.914. Note that due to the structure of the Rigolet data, a random intercept was forced in the model to control for household-level clustering.

Table 4. Univariable exact logistic regression* and final multivariable logistic regression (controlling for household clustering) results, □* examining the effects of predictor variables on the odds of acute gastrointestinal illness in Rigolet, Nunatsiavut, in May 2013.

RIGOLET MAY MODEL	n	Univariable Results			Multivariable Model Results		
		Odds Ratio	P	Confidence Interval (95%)	Odds Ratio	P	Confidence Interval (95%)
Exposure to a puppy in past month							
No	219	<i>referent</i>			<i>referent</i>		
Yes	13	6.20	0.002	1.93 - 19.87	16.16	0.029	1.32 - 197.63
Tap water consumption in past two weeks							
No	125	2.77	0.015	1.22 - 6.30	3.74	0.036	1.09 - 12.81
Yes	111	<i>referent</i>			<i>referent</i>		
Age Group							
0-19	74	<i>referent</i>			<i>referent</i>		
20-55	119	2.50	0.061	0.96 - 6.54	4.57	0.023	1.23 - 16.97
Over 55	43	1.57	0.478	0.45 - 5.51	2.89	0.242	0.49 - 16.93
Consumption of raw or undercooked retail eggs in the past month							
No	147	<i>referent</i>			-	-	- -
Yes	85	2.17	0.045	1.02 - 4.60	-	-	- -
Visited a cabin in past month							
No	110	2.60	0.088	0.87 - 7.82	-	-	- -
Yes	126	<i>referent</i>			-	-	- -
Visited another community in past two weeks							
No	161	<i>referent</i>			-	-	- -
Yes	75	2.15	0.049	1.00 - 4.58	-	-	- -
Weekly amount spent on retail food							
Less than \$150	44	<i>referent</i>			-	-	- -
\$150-\$300	154	1.32	0.634	0.42 - 4.13	-	-	- -
Over \$300	31	4.18	0.030	1.15 - 15.21	-	-	- -
Primary source of drinking water in past two weeks							
Tap water	93	<i>referent</i>			-	-	- -
Brook water	14	2.86	0.160	0.66 - 12.43	-	-	- -
Bottled water	124	2.23	0.070	0.94 - 5.29	-	-	- -
Consumption of brook water in past two weeks							
No	178	<i>referent</i>			-	-	- -
Yes	53	1.96	0.103	0.87 - 4.38	-	-	- -
Storage of drinking water							
Container in the fridge	100	0.44	0.064	0.18 - 1.05	-	-	- -
Container outside of the fridge	56	<i>referent</i>			-	-	- -
No storage	75	0.33	0.029	0.12 - 0.89	-	-	- -
Quantity of tap water consumed yesterday							
None of water consumed	125	3.05	0.021	1.19 - 7.88	-	-	- -
Some or most of water consumed	26	1.88	0.399	0.43 - 8.16	-	-	- -
All of the water consumed	85	<i>referent</i>			-	-	- -
Exposure to dogs in past month							
No	62	<i>referent</i>			-	-	- -
Yes	172	2.15	0.136	0.79 - 5.85	-	-	- -

* The results from the univariable analysis are presented for those variables with $p < 0.20$.

□* Likelihood ratio test comparing the model with and without the household-level variable: variance = 2.54, 95% CI = 0.55-11.65, $p = 0.011$; Intra-class correlation coefficient = 0.44, 95% CI = 0.14-0.78.

Table 5. Univariable* and multivariable exact logistic regression model results, examining the effects of predictor variables on the odds of acute gastrointestinal illness in Iqaluit, Nunavut, in September 2012.

IQALUIT SEPTEMBER MODEL	<i>n</i>	Univariable Logistic Results			Multivariable Exact Logistic Model Results		
		Odds Ratio	<i>P</i>	Confidence Interval (95%)	Odds Ratio	<i>P</i>	Confidence Interval (95%)
Country fish consumption							
Less than half of meals	458	<i>referent</i>			<i>referent</i>		
Half or more of meals	64	2.44	0.006	1.30 - 4.60	3.26	0.001	1.54 - 6.70
Exposure to Cats							
No	436	<i>referent</i>			<i>referent</i>		
Yes	90	2.35	0.003	1.33 - 4.16	2.07	0.023	1.06 - 3.95
Employment status of the person responsible for food preparation							
Not Employed	217	<i>referent</i>			<i>referent</i>		
Employed	306	2.13	0.008	1.22 - 3.73	2.12	0.016	1.13 - 4.13
Washing counter with soap							
No	102	2.38	0.002	1.37 - 4.10	1.95	0.024	1.03 - 3.59
Yes	418	<i>referent</i>			<i>referent</i>		
Car ownership							
No	262	<i>referent</i>			-	-	- -
Yes	256	1.76	0.032	1.05 - 2.94	-	-	- -
Perceived quality of drinking water							
Very poor or poor	37	2.14	0.061	0.96 - 4.74	-	-	- -
Fair, good, or very good	482	<i>referent</i>			-	-	- -
Number of days leftovers are kept							
1 day	71	<i>referent</i>					
2 days	300	1.23	0.060	0.96 - 6.57			
3 days or more	148	1.02	0.205	0.70 - 5.44			
Type of house							
Public housing	194	<i>referent</i>			-	-	- -
Rented	199	2.02	0.024	1.10 - 3.71	-	-	- -
Owned	126	1.85	0.078	0.93 - 3.65	-	-	- -
Monthly living expenses							
Low (\$500 or less)	135	<i>referent</i>			-	-	- -
Medium (\$501-\$1,700)	166	2.32	0.032	1.08 - 5.01	-	-	- -
High (\$1,701 or more)	226	2.37	0.022	1.13 - 4.95	-	-	- -
Education level of person responsible for food preparation in the home							
Elementary or lower	202	<i>referent</i>			-	-	- -
High school or higher	330	1.95	0.021	1.11 - 3.43	-	-	- -

* The results from the univariable analysis are presented for those variables with $p < 0.10$; however, those with $p < 0.20$ were considered in the multivariable model building process.

Table 6. Univariable exact logistic results (for those variables with $p < 0.20$) and final multivariable Exact logistic regression model results, examining the effects of predictor variables on the odds of acute gastrointestinal illness in Iqaluit, Nunavut, in May 2013.

IQALUIT MAY MODEL	<i>n</i>	Univariable Exact Logistic Results			Multivariable Exact Logistic Model Results		
		Odds Ratio	<i>P</i>	Confidence Interval (95%)	Odds Ratio	<i>P</i>	Confidence Interval (95%)
Homeless person staying in the house							
No	305	<i>referent</i>					
Yes	205	2.29	0.005	1.24 - 4.29	2.05	0.016	1.09 - 3.88
Pet Ownership							
No	314	<i>referent</i>			-	-	- -
Yes	196	1.49	0.181	0.81 - 2.75	-	-	- -
Over-crowding							
No	388	<i>referent</i>			-	-	- -
Yes	124	1.55	0.175	0.78 - 2.97	-	-	- -
Pet Ownership x Over-crowding							
Pets + Over-crowding		-	-	- -	<i>referent</i>		
Pets + No over-crowding		-	-	- -	3.19	0.010	1.32 - 7.72
Pet Ownership x Over-crowding							
Pets + Over-crowding					<i>referent</i>		
No Pets + No over-crowding					2.87	0.010	1.29 - 6.38
Pet Ownership x Over-crowding							
Pets + Over-crowding					<i>referent</i>		
No pets + Over-crowding					4.40	0.010	1.42 - 13.64
Country meat consumption							
Less than half of meals	414	<i>referent</i>			-	-	- -
Half or more of meals	95	1.91	0.059	0.93 - 3.77	-	-	- -
Consumption of dried meat							
No	347	<i>referent</i>			-	-	- -
Yes	164	1.72	0.086	0.92 - 3.19	-	-	- -
Sex							
Male	218	<i>referent</i>			-	-	- -
Female	294	1.65	0.109	0.87 - 3.23	-	-	- -
Perceived quality of drinking water							
Very poor or poor	32	2.22	0.118	0.71 - 5.92	-	-	- -
Fair, good, or very good	468	<i>referent</i>			-	-	- -
Exposure to Working Dog							
No	493	<i>referent</i>			-	-	- -
Yes	19	2.41	0.123	0.56 - 7.98	-	-	- -
Municipal Water Supply							
Tap Water	420	2.14	0.126	0.82 - 7.10	-	-	- -
Trucked Water	88	<i>referent</i>			-	-	- -
Weekly amount spent on retail food							
Under \$300	266	<i>referent</i>			-	-	- -
\$301+	231	1.61	0.132	0.85 - 3.08	-	-	- -
Exposure to cats							
No	423	<i>referent</i>			-	-	- -
Yes	89	1.64	0.178	0.77 - 3.31	-	-	- -
Washing hands before food preparation							
No	140	1.55	0.192	0.80 - 2.92	-	-	- -
Yes	370	<i>referent</i>			-	-	- -

FIGURES




Figure 1. A map of Labrador and Nunavut, highlighting the communities in Nunavut and the Nunatsiavut Land Claim Settlement Region.

Acute Gastrointestinal Illness	Demographic Information	Socio-Economic Levels	Food Consumed	Water Consumed	Exposure to Animals
<ul style="list-style-type: none"> • Primary symptoms • Secondary symptoms • Severity of symptoms • Healthcare seeking behaviours 	<ul style="list-style-type: none"> • Gender • Age • Aboriginal Identity • Household ID 	<ul style="list-style-type: none"> • Education Level • Overcrowding • Employment Status • Household Expenses • Household Items • Food Security Level 	<ul style="list-style-type: none"> • Type of Food (e.g. meat, fruits and vegetables, ready-to-eat foods, dairy) • Source of Food (e.g. retail vs country food) • Frequency of Food Consumed • Preparation method (e.g. cooked, raw, frozen, dried, fermented) • Food Handling Behaviours 	<ul style="list-style-type: none"> • Primary and Secondary Sources of Drinking Water (e.g. tap, brook, bottled, melted snow/ice) • Piped vs. Trucked Water • Amount consumed • Secondary Treatment • Storage of Water (e.g. counter-top vs. fridge) • Perceived Quality 	<ul style="list-style-type: none"> • Pets • Working Animals • Wild Animals • Animals inside the home

Figure 2. A summary of the information captured in the cross-sectional surveys in Rigolet, Nunatsiavut, and Iqaluit, Nunavut (2012, 2013).




Figure 3. The estimated annual incidence of acute gastrointestinal illness in Hamilton,[29] British Columbia,[29] Ontario,[4] Rigolet, and Iqaluit, Canada, using a case definition of self-reported diarrhea and/or vomiting in the past 28-days not due to pregnancy, medication, drugs/alcohol, or chronic conditions