



Tecnociencia 2015, Vol. 17, N° 1.

THE PANAMANIAN CLIMBING RAT, MAMMALIA, RODENTIA, CRICETIDAE, *TYLOMYS PANAMENSIS* (GRAY, 1873): NEW REPORT IN DARIEN

Pedro Méndez-Carvajal^{1,2,3}, Marcos Peñafiel³, Alvin Zapata¹, and Guido Berguido¹

¹Fundación Pro-Conservación de los Primates Panameños (FCPP), 0816-07905 Panamá, República de Panamá.

²Department of Anthropology, Durham University, Durham, UK, South Road, DH1 3LE.

³Sociedad Panameña de Biología (SPB), Panamá, República de Panamá.

*Corresponding author Email: fcprimatespanama@gmail.com

ABSTRACT

This study reports a new record for *Tylomys panamensis*, increasing its range by 121 km, 66° NW from the original endemic distribution. It has been considered as Data Deficient by IUCN as no information has been published for almost 64 years. Based on our long term surveillance at the Darien's canopy, using Orion Camera System (OCS), we have been able to obtain information about their circadian and monthly activity over the course of one year. The presence of *T. panamensis* remarks the importance of the largest National Park of the Mesoamerican region and claim urgency to be protected from its actual devastation, particularly the montane forest in the rest of the provinces of Darien and Panama.

KEYWORDS

Climbing rat, Darien, mammals, Panama, *Tylomys panamensis*.

LA RATA ESCALADORA PANAMEÑA, MAMMALIA, RODENTIA, CRICETIDAE, *TYLOMYS PANAMENSIS* (GRAY, 1873): NUEVO AVISTAMIENTO EN DARIÉN

RESUMEN

El estudio informa un nuevo avistamiento para *Tylomys panamensis*, incrementando su rango a 121 km, 66° NW de su distribución original. Esta especie ha sido considerada como Data Deficiente por UICN dado que no habría información publicada por casi 64 años. Basados en nuestra vigilancia de dosel en Darién a largo plazo utilizando el Sistema de Cámaras Orión (SCO), hemos podido obtener información de su actividad circadiana y actividad mensual por espacio de un año. La presencia de *T. panamensis* confirma la importancia del mayor Parque Nacional de la región Mesoamericana clamando con urgencia a ser protegido de su actual devastación, y en particular los bosques montañosos del resto de las provincias de Darién y Panamá.

PALABRAS CLAVES

Darién, Mamíferos, Panamá, Rata escaladora, *Tylomys panamensis*.

INTRUCTION

Tylomys panamensis: Panamanian Climbing Rat (Gray, 1873), have been classified as Data Deficient by IUCN Red list (Reid & Woodman, 2008), and Vulnerable according to the Environmental Authority of Panama (ANAM, 2012), due to its regional endemism and restricted distribution to Cana, Jaqué, Paya, Pirre, and the Sapo Mountain Chain, Darien province (Alston, 1882; Reid & Woodman, 2008). This rat was first classified as *Neomys panamensis* by Gray (1873), then changed to *Hesperomys panamensis*, and finally to *Tylomys panamensis* in 1879 (Miller Jr., 1911). It has been discussed in terms of similitude with the other climbing rats from this genus, *T. fulviventer*, the later has argus brownish coloration, sides lighter in tone than the back, and hands and feet dusky above (Anthony, 1916). However, it was suggested that *T. fulviventer* be considered junior synonyms of *T. panamensis* (Handley, 1966; Musser & Carleton, 2005). In a later review of Goldman (1920), *T. panamensis* is reported as an arboreal rat, uncommon for low lands, reported only for Cana and Paya River in Darien province at 600 m.a.s.l. (Handley, 1966). So far there is little information related to the actual distribution and general ecology of *T. panamensis* populations. The only information published is on coloration characteristics: white-

bellied, sides, fore and hind limbs with same coloration brownish or slate gray (Anthony, 1916). The head is similar in coloration as the body, with dark color extending to the ears and bicolor tail (Anthony, 1916). Genetic studies based in two possible specimens identified as *T. panamensis* determined that this species is the highest diploid number of this group recorded, with $2n=52$ (Pathak *et al.*, 1973).

METHODS

The Fundación Pro-Conservación de los Primates Panameños (FCPP), together with the Sociedad Panameña de Biología (SPB) have been surveying different areas in Panama to generate a current distribution database, following Graham *et al.* (2004) and Guisan & Wilfried (2005). Several methods of mammal detection have been applied, such as using understory and canopy techniques to set camera traps, mistnets and *ad-libitum* observations in Panama.

Since March 4, 2013 to the present, we have been collecting information from five Bushnell-Trophy CamTM camera traps, settled at the canopy at Chucanti Nature Reserve, Darien province, Panama, with an average level of eight to 12 m high, using Orion Camera System (OCS), a new method to deploy camera traps at canopy (Méndez-Carvajal, 2014). The cameras are part of a long term project of surveillance to study the Darien Black spider monkey *Ateles fusciceps rufiventris* and related fauna at the tree top led by FCPP, installed into the highest point of the Majé Mountain Chain, Chucanti Nature Reserve, with an extensive mature forest of 3.5 km² (Méndez-Carvajal, 2012). The study site is located in the frontier border between the Panama Province and Darien Province with (08° 47'16.5" N, 078° 27'01.4"W). Its higher zone reaches the 1,350 m.a.s.l., with temperatures between 24-27 °C, and annual rainfall of 1,940.5 mm (Laurance, 2008).

RESULTS AND DISCUSSION

We obtained a total of 5,976 hours of trap footage, with continued data collected over the course of 249 days from five cameras. Two cams detected *T. panamensis*: Cam 1 (08°47'50.3" N, 078°27'43.8"W), located at 800 m.a.s.l., which obtained an index of appearance of 0.004

ind/year and Cam 3 (08°47'51.2" N, 078°28'3.1"W), located at 850 m.a.s.l., with 0.5 ind/year (Fig. 1). We found *T. panamensis* to be more active in the rainy season between June and September, showing a peak in August (Fig. 2), coinciding with the breeding period reported for the Peromyscini tribe (Eisenberg & Redford, 1999). During data collection, this climbing rat presented strict nocturnal-crepuscular, arboreal activity, from approximately 21:00 to 06:00 hrs, and presenting more activity at midnight (Fig. 2). For Cam 3, we observed two different individuals at 12 m high, using a microhabitat composed by a liana *Bauhinia spp.* (Fabaceae) which was used for the rat as a link to connect with the palms *Oenocarpus spp.*, and *Attalea spp.* (Palmae), Orchids (Orchidaceae), Bromeliads (Bromeliaceae), ferns, other epiphytic plants from Cychlanthaceae family, and *Anthurios spp.* (Araceae). *T. panamensis* passes almost unnoticed, with a speed average of 17 cm/s (n = 5), we applied $d = v \cdot t$ (d = distance, v = velocity, t = time) to calculate the approximate distance traveled from the camera position elsewhere. Using the maximum time expended to return as four minutes, we calculated that this rat could go as far as 20 m from its possible nest if traveling at relatively constant velocity over time and space. Based upon several cam trap photographs taken of two different individuals, and the relation with the dimensions of the surrounding vegetation following Lambert *et al.* (2012), we calculated a total head-body size of 226.6 mm (n=10, SD±24.3) and tail of 199.1 mm (n=10, SD±29.7), and total length 423.3 mm (n=6, SD±22.5). Unpublished data were found reviewing information at the National Museum of Natural History (NMNH) from United States and the Instituto Conmemorativo Gorgas de Estudios de la Salud (ICGES) from Panama, and its mammal collection, the Colección Zoológica Eustorgio Méndez (CoZEM). We found specimens of *T. panamensis* collected from NMNH and CoZEM coincide with the average calculated for body size of the specimen captured in camera (Figs. 3 and 4). General average body size calculated from museum specimens and photos were: TL 423(380-470); HB 218 (166-255); T 205(144-226); RF 38(32-42) and EN 18(10-25) (Fig. 4). Other mammals that appeared in the same cameras were: *Sciurus granatensis*, *Potos flavus*, *Tamandua mexicana*, and *Ateles fusciceps rufiventris* (Méndez-Carvajal, 2014). Using our criteria of high elevation, cloudy forest, and remnant forested zone, we consider the actual distribution range for this species should be expanded using the mountain chain forest that is

left in the area of Darien until Chucanti Nature Reserve and Majé Mountain Chain from Panama province.




Fig. 1. *Tylomys panamensis* photographed at Chucanti Nature Reserve, Darien. A) Bicolor tail, B) Gray dorsal coloration, C) White ventral coloration D) *T. panamensis* using *Bauhinia spp.*, as a way to move from the top to the down side E) *T. panamensis* jumping position, F) Vertical movement.




Fig. 2. Activity pattern for *T. panamensis* in terms of a round year camera trap study; and circadian activity for *T. panamensis* at Chucanti Nature Reserve, Darien, Panama.




Fig. 3. *T. panamensis*, previous range and new sight at Chucanti Nature Reserve, Darien province, Republic of Panama.




Fig. 4. Specimens courtesy of the National Museum of Natural History (NMNH): A) 314585 B) 314586 C) 314587 and D) 314587. Specimens courtesy of the Colección Zoológica Eustorgio Méndez (CoZEM): E) H-0284 F) lateral view of H-0284 in black and white to compare with our individuals photo-captured with infrared sensor. TL: Total Length; HB: Head-Body; T: Tail; HF: Hind Foot; EN: Ear Notch.

CONCLUSIONS

This report is the first obtained since the 1950's (Musser & Carleton, 2005; Reid & Woodman, 2008), offering data that expand *T. panamensis* information and distribution at 121 km, 66° NW from the original distribution taking as a reference Pirre Mountain Chain. The lack of information could be related with the cryptic behavior of this arboreal and nocturnal rat, as well as the altitudinal range where this rat is found. We consider Chucanti Nature Reserve as a valuable forested zone, and one of the few areas into the Darien province that could be protecting the existing endemic fauna and flora of its cloudy forest in the Majé Mountain Chain, which connects the Bayano forest with the Panama province, Republic of Panama.

ACKNOWLEDGEMENTS

We thank the collaboration of the Panamanian Environmental Authority (ANAM), Department of Wildlife Management, and the Colección Zoológica Eustorgio Méndez (CoZEM) from the Instituto Conmemorativo Gorgas de Estudios de la Salud (ICGES), with special thanks to Aydee Cornejo and Nohelys Alvarado, for their great support with the specimens. We really appreciate the collaboration of Stephanie Canington from the National Museum of Natural History at Smithsonian Institution, who kindly helped with specimens' information. We appreciate the help of Joanna Setchell and Russell Hill for their comments that greatly improved the manuscript. We are grateful with Milvia Sánchez, Yayson Péres, Arcadio Castillo, Elena Méndez, and Ricardo Moreno, who helped in the field in the scientific station of Chucanti. We deeply appreciate the support of Idea Wild, Mohamed bin-Zayed Species Conservation Fund (projects # 1025476 and 12055182), and II Rufford Small Grant (project #16021-2). Our special appreciation to Virginia Commonwealth University (VCU), for their first camera traps donated, to Cathy Viverette, Edward Crawford, and especially to Kensey Barker and Ryan Weaver who kindly reviewed the English of the manuscript. Pedro Méndez-Carvajal would like to thank the government of Panama through the Senacyt-IFARHU for the doctoral scholarship that will allow analyzing the long term data at the University of Durham, UK.

REFERENCES

- Alston, E.R. 1882. *Biologia centrali-americana*. Mammalia. R.H. Porter, 220 p.
- ANAM (Autoridad Nacional del Ambiente de Panamá). 2012. Taller de actualización de especies amenazadas y endémicas de Panamá (mamíferos). Anexo.
- Anthony, H.E. 1916. Panama Mammals Collected in 1914-1915. *American Museum of Natural History*. Art Vol. XXXV, Art. XX, New York, USA. 357-376.
- Eisenberg, J. F. & K.H. Redford. 1999. Mammals of the Neotropics: The central Neotropics. Ecuador, Peru, Bolivia, Brazil. The University of Chicago Press, Chicago, USA. 609p.
- Goldman, E. A. 1920. Mammals of Panama. *Smithsonian Misc. Coll* 69: 1-309.
- Graham, C.H. S. Ferrier, F. Huettman, C. Moritz, & A.T. Peterson. 2004. New developments in museum-based informatics and applications in biodiversity analysis. *Trends. Ecol. Evol.* 19: 497-505.
- Gray. 1873. *Tylomys panamensis*. *Ann. & Mag. Nat. Hist. ser.* 4(12): 417.
- Guisan, A. & T. Wildfried. 2005. Predicting species distribution: offering more than simple habitat models. *Ecology letters* 8:993-1009.
- Handley, C.O. 1966. Checklist of the Mammals of Panama; pp. 753-795, in: R.L. Wenzel and J. Tipton (eds.). *Ectoparasites of Panama*, Field Mus. Nat. Hist., Chicago. USA.
- Lambert, M.R., M.C. Yasuda, & B.D. Todd. 2012. Evaluation of a photographic technique for estimating body size in lizards from distance. *Herpetological Conservation and Biology*. 7(1): 83-88.
- Laurance, W.F. 2008. Adopt a Forest. *Biotropica*. 40(1): 3-6.

Méndez-Carvajal, P.G. 2012. Preliminary Primate Survey at the Chucanti Nature Reserve, Darien Province. *Mesoamericana* 16 (3): 22-29.

Méndez-Carvajal, P.G. 2014. The Orion Camera System, a new method for deploying camera traps in tree canopy to study arboreal primates and other mammals: A case study in Panama. *Mesoamericana*, 18(1): 9-23.

Miller, G.S. Jr. 1911. Genus *Tylomys* Peters. *List of North American land mammals in the United States National Museum*. 179 pp.

Musser, G.G. & M.D. Carleton. 2005. Rodentia: *Tylomys panamensis*; pp. 2,142, in: D.E. Wilson and D.M. Reeder (eds). *Mammal Species of the World. A Taxonomic and Geographic Reference* (3rd ed), Johns Hopkins University Press, USA.

Pathak, S.H., T.C. Hsu, L. Shirley & J.D. Helm III. 1973. Chromosome homology in the climbing rats, genus *Tylomys* (Rodentia: Cricetidae). *Chromosoma* 42: 215-228.

Reid, R. & N. Woodman. 2008. *Tylomys panamensis*. In: IUCN 2011. IUCN Red List of Threatened Species. Version 2014. www.iucnredlist.org. Retrieved on 28 May 2014.

Recibido marzo de 2015, aceptado abril de 2015.