



ISSN: 2091-2749 (Print)
2091-2757 (Online)

Correspondence

Dr. Alka Singh
Associate Professor, Department
of Obstetrics and Gynaecology,
Patan Academy of Health
Sciences
Email:singhalka@hotmail.com

Peer Reviewed By

Dr. Jay N Shah
Patan Academy of Health
Sciences

Peer Reviewed By

Dr. Ashis Shrestha
Patan Academy of Health
Sciences

Prevalence of hepatitis B in pregnant women and management of babies born to Hepatitis B positive mother: A criterion based clinical audit

Alka Singh,¹ Suchita Joshi,² Yubaraj Sharma,³

¹Associate Professor, Department of Obstetrics and Gynaecology,
² Assistant Professor, Department of Pediatrics, ³Associate Professor,
Department of Medicine, Patan Academy of Health Sciences,
Lalitpur, Nepal

ABSTRACT

Introductions: This study aims at identifying the prevalence of Hepatitis B virus infection in pregnant women and adherence to the management of the newborns of these women as per the international guidelines.

Methods: This is a cross sectional study conducted at Patan Hospital from January 1, 2008 to December 30, 2014. Prevalence of HBsAg in pregnant women in Patan Hospital and management of babies born to these women, the immunization status, breast feeding status and follow up advise for the babies were studied. These data were analyzed to see whether we were following international guidelines for preventing mother to child transmission of hepatitis B infection.

Results: There were total of 58,917 deliveries and 59,438 births during the seven year study period. During this period, total 148 cases of pregnant ladies with HBsAg positive status were recorded. The prevalence of HBsAg positive case was 0.25% or 1 in every 398 deliveries. Fifty four babies (87%) received hepatitis B vaccine (HBV) and hepatitis B immunoglobulin (HBIG) within 12 hours of birth. Eight babies (22%) failed to receive HBV and HBIG within 12hours of birth. Fifty eight (96%) babies were breastfed after birth. Mother's HBsAg positive status was not the reason for formula feeding in any of the babies. Follow up HBsAg status was not advised in any of the babies.

Conclusions: At Patan hospital, we seem to be following most of the guidelines in the management of HBV infection, except for checking the HBsAg status of the babies at 9-18 months of age.

Keywords: chronic liver disease, HBsAg positive, hepatitis B vaccine, Hepatitis B immunoglobulin, mother to child transfer

INTRODUCTIONS

Newborn infants acquiring HBV infection by perinatal transmission have a greater than 95% chance of becoming chronic HBV carriers.¹ Therefore, it is very important to establish protocols in institutes to prevent mother to child transfer (MTCT) and decrease the burden of chronic HBV infection.

Neonates of all HBsAg positive mothers should receive immunoprophylaxis treatment with HBIG and HBV given within 12 hours of birth to decrease MTCT of hepatitis B virus infection.² This is 85 to 95 percent effective in reducing the infection from vertical transmission.³ The guidelines also advise to have the HBsAg status of the babies checked at 9-18 months of age.⁴

The objectives of this study was to audit if the recommendation for preventing MTCT of hepatitis B and guidelines for babies born to HBsAg positive mothers were followed in Patan Hospital.

METHODS

This was a cross sectional study done from January 1, 2008 to December 30, 2014, conducted in Patan Hospital. Delivery record files were checked to confirm that HBsAg status was recorded in all women. All HBsAg positive results of pregnant women recorded in the hospital laboratory record book were selected. The record files of these patients and their babies were retrieved from the record filing section. The mode of delivery and indication for cesarean sections were noted. The immunization status and breast feeding status of the babies were also recorded from the baby's hospital record and drug cardex. These data were analyzed to see whether we were following international guidelines for preventing mother to child transmission of hepatitis B infection which included administration of HBIG and HBV vaccine to the neonates of HBsAg-positive mother within 12 hours of delivery.

Women who delivered outside Patan Hospital and mother or baby's record file that were missing were excluded from this study.

The files were reviewed and the following answers were sought for.

1. Were all pregnant women screened for hepatitis B?
2. Was HBsAg positive status wrongly used as an indication for Cesarean section?
3. Did all babies born to HBsAg positive women receive HBV and HBIG within 12 hours of birth?
4. Was HBsAg status wrongly used as a contraindication for breastfeeding?
5. Was HBsAg status checked at 9-18 months in all babies born to HBsAg positive mothers?

RESULTS

There were total of 58,917 deliveries and 59,438 births during the seven year study period. From the maternity record, it was seen that all the women were tested for HBsAg before delivery except for five women who had no antenatal checkups.

During this period, 148 cases of pregnant ladies with HBsAg positive status were recorded. The prevalence of HBsAg positive case was 0.25% or 1 in every 398 deliveries. Among these 148 cases 80(54.1%) cases were excluded from the study because their hospital records could not be traced. The reason behind not being able to find the hospital records among 45 was because they had temporary hospital numbers. In our hospital, those with temporary numbers are not filed in the hospital record section. Among all the pregnant ladies with HBsAg positive status, 68(45.9%) hospital record files were successfully traced. Fifteen(22%) of these cases had delivery by cesarean section and none of them were for HBsAg status. Hepatitis B infection was not an indication for cesarean section in any of the cases. The indications were: breech(2), meconium(4), failed induction(3), previous cesarean with

unfavorable cervix(2), cephalopelvic disproportion(1) and non progress of labor(3).

Six out of the remaining 68 cases were excluded from the study because four babies were not delivered in Patan Hospital and therefore, they did not have hospital record files and two baby records were missing.

Finally, the total cases included in this study were 62. The average maternal age was 27 years. None of the cases were positive for HIV and VDRL which is also done routinely for all antenatal clients.

The genders of the babies born to these mothers, 35(56.4%) were male and 27(43.5%) were female. The average gestational age was 38 weeks (four were premature births). Fifty four(87%) received HBV and HBIG within 12 hours of birth. Eight babies(22%) failed to receive HBV and HBIG within 12hours of birth as per the hospital pediatric department guidelines. Fifty eight(94%) babies were breastfed after birth. Two(3.2%) mothers did not breastfeed because the babies were on intravenous drip, and the other two mothers did not breast feed because they had cesarean delivery and felt too weak. Mother's HBsAg positive status was not the reason for formula feeding in any of the babies. None of the babies were advised to have their HBsAg status checked at 9-18 months of age.

DISCUSSIONS

A hospital based retrospective study was carried out. The prevalence of HBsAg positive was 0.25% in this study. This is similar to a US study, where the prevalence of chronic hepatitis B virus infection in pregnancy was 0.2 to 6%, with rates varying by race and ethnicity.⁵

In Nigeria the incidence of seroprevalence of HBsAg was 8.3%⁶, much higher than ours. The CDC recommends that all pregnant women should be screened for the presence of HBsAg at diagnosis of pregnancy.⁷ This is also supported by U.S. Preventive Services Task

Force, American Academy of Family Physicians, American College of Obstetricians and Gynecologists, the American Academy of Pediatrics, and the Centers for Disease Control and Prevention. Those pregnant women with unknown HBsAg status or with risk factors should be screened upon admission for delivery.⁸

Although it is recommended that all pregnant women should be tested for HBsAg status to prevent MTCT of Hepatitis B infection, this is not yet practiced in most developing nations like Nigeria, hence the high prevalence. In developed countries, the increased awareness, identification of mothers who are Hepatitis B surface antigen and adequate prophylaxis reduce the prevalence of HBV infection.⁹ In a large population-based study from Florida involving nearly 1.7 million pregnant women, the prevalence of HBV was approximately 27 times higher among Asian-Americans and 5 times higher among African-Americans as compared with whites.¹⁰

This emphasizes the importance of screening of pregnant women for HBsAg status in our country. It was very reassuring to note that in Patan Hospital, we have been successful in screening all pregnant women for HBsAg status. Prevalence rate in this study is comparable to western countries and this is probably because the antenatal population in our hospital is mostly not drug abusers and sexual workers and most of their previous deliveries were done in hospital setup.

Pregnancy is not a contraindication for vaccination to hepatitis B virus.¹¹ Pregnant women who are not immune to hepatitis B virus should be vaccinated, because premature delivery may be increased if acute hepatitis B is acquired in the last trimester and because MTCT occurs in over 60% of pregnancies associated with acute hepatitis B infection at or near term.¹¹ Offering HBV during pregnancy is not a routine practice in our hospital and this can be considered.

Current guidelines do not recommend elective cesarean delivery for mothers with chronic

HBV infection, since this does not reduce the risk of mother to child transmission of HBV. Therefore, elective cesarean section should not be used in HBsAg positive pregnant women to prevent mother to child transmission of HBV.¹² It was reassuring to note that in our hospital, HBsAg positive status alone was not being practiced as an indication for elective cesarean delivery.

Multiple studies have shown breastfeeding to be safe for children with mothers with chronic HBV infection.¹³ The American Academy of Pediatrics states that breastfeeding is not contra-indicated in women who are HBsAg positive.¹⁴ Although breast milk contains HBsAg, breast feeding does not increase the risk of MTCT of hepatitis B virus, especially if the baby receives HBV and HBIG within 12 hours of birth.¹⁵ In Patan hospital, we do not discourage breast feeding in these mothers and this has been reflected in this audit.

HBIG and HBV can effectively prevent mother-to-child transmission of hepatitis B virus.¹⁶ From this study we can see that in Patan Hospital giving HBIG and HBV to babies born to HBsAg positive mother is a routine practice. Seven babies that missed this were because their mother's HBsAg status was not written in neonatal record sheet and one baby was discharged before mother's HbAg test result was available.

Unfortunately this study shows that none of the babies were advised to have their HBsAg status tested at 9-18 months as the guidelines suggest.

The limitation of this study was that we were unable to trace the record files of those patients who did not have permanent hospital identity.

CONCLUSIONS

We at Patan Hospital seem to be following most of the guidelines such as screening all women for HBsAg antenatally, not making HBsAg positive an indication for cesarean

section, giving HBIG and HBV vaccines before 12 hours after birth and breastfeeding. But we are not checking the HBsAg status of the babies at 9-18 months of age as also suggested by the guidelines. Therefore, this can be introduced in our practice. In order not to miss any baby whose mother is HBsAg positive, it would be a good idea to have a designated space to write the mother's HBsAg status in the neonatal record sheet as we already have for the VDRL and HIV status. Besides, the junior doctors carrying out newborn examination should be trained to check maternal notes as well so as not to miss any baby who needs HBIG and HBV vaccine. As for checking HBsAg and HBAb status of the babies at 9-18 months, this is not yet stated in our hospital guidelines. As this is now a standard recommendation, we should amend this in our own guideline and re-audit to see if the changes have improved our practice.

REFERENCES

1. Beasley RP, Trepo C, Stevens CE, Szmuness W. The e antigen and vertical transmission of hepatitis B surface antigen. *Am J Epidemiol* 1977;105(2):94–98.
2. Centers for Disease Control and Prevention (CDC). Assessing completeness of perinatal hepatitis B virus infection reporting through comparison of immunization program and surveillance data—United States. *MMWR Morb Mortal Wkly Rep* 2011; 60(13): 410–413. *J Perinatol* 1991; 8(3): 227–232.
3. Mast EE, Margolis HS, Fiore AE, et al. A comprehensive immunization strategy to eliminate transmission of hepatitis B virus infection in the United States: recommendations of the Advisory Committee on Immunization Practices (ACIP) part 1: immunization of infants, children, and adolescents [published corrections appear in *MMWR Morb Mortal Wkly Rep*. 2006;55(6):158-159 and *MMWR Morb Mortal Wkly Rep*. 2007;56(48):1267]. *MMWR Recomm Rep*. 2005;54(RR-16):1-31.

4. Centers for disease control and prevention. Recommendations for Routine Testing and Follow-up for chronic Hepatitis B virus (HBV) Infection. *MMWR* 2008; 57 (No. RR-Gambarin-Gelwan M. Hepatitis B in pregnancy. *Clin Liver Dis* 2007; 11(4):945–963.
5. Anaedobe CG, Fowotade A, Omoruyi CE, Bakare RA. Prevalence , socio-demographic features and risk factors of Hepatitis B virus infectioin among pregnanat women in southwestern Nigeria. *Pan African Medical Journal* 2015; 20:406.
6. ACOG educational bulletin. Viral hepatitis in pregnancy. Number 248, July 1998 replaces No. 174, November 1992). American College of Obstetricians and Gynecologists. *Int J Gynaecol Obstet* 1998; 63(2): 195–202.
7. U.S. Preventive Services Task Force. Screening for Hepatitis B virus Infection: Recommendation statement [Internet]. Rockville, MD: Agency for Healthcare Research and Quality; 2004 February [cited 2010 September 5]. Available from: <https://www.uspreventiveservicestaskforce.org/Home/GetFile/1/645/hepbvs/pdf/>.
8. Jones MN, Hepatitis B and Pregnancy. An underestimated issue. *Liv Int.* 2009;29:133-139. Conell LE, Salihu HM, Salemi JL, August EM, Weldeselasse H, Mbah AK. Maternal hepatitis B and hepatitis C carrier status and pernatal outcomes. *Liverint* 2011;31(8):1163-1170.
9. Levy M, Koren G. Hepatitis B vaccine in pregnancy: maternal and fetal safety. *Am J Perinatol* 1991; 8(3): 227–232.
10. Hu Y, Chen J, Wen J, Xu C, Zhang S, Xu B, Zhou YH. Effect of elective cesarean section on the risk of mother to child transmission of hepatitis B virus. *BMC Pregnancy Childbirth* 2013 May 24; 13:119 doi:10.1186/1471-2393-13-119.
11. Shi Z, Yang Y, Wang H, Ma L, Schreiber A, Li X, Sun W, Zhao X, Yang X, Zhang L, Lu W, Teng J, An Y. Breastfeeding of newborns by mothers carrying hepatitis B virus: a meta-analysis and systematic review. *Arch Pediatr Adolesc Med.* 2011 Sep;165(9):837-56.
12. Gartner LM, Morton J, Lawrence RA, Naylor AJ, O'Hare D, Schanler RJ et al. Breastfeeding and the use of human milk. *Pediatrics* 2005; 115(2): 496–506.
13. Wong VC, Lee AK, Ip HM. Transmission of hepatitis B antigens from symptom free carrier mothers to the fetus and the infant. *Br J Obstet Gynaecol* 1980; 87(11): 958–965.
14. Zhang L, Gui XE, Teter C, Zhong H, Pang Z, Ding L, Li F, Zhoi Y, Zhang L. Effect of hepatitis B immunization on prevention of mother-to-infant transmission of hepatitis B virus and on the immune response of infants towards hepatitis B vaccine. *Vaccine.* 2014 Oct 21;32(46):6091-7.