



ISSN: 2091-2749 (Print)
2091-2757 (Online)

Correspondence

Dr. Gajendra Mani Shah
Department of Orthopaedics
and Trauma Surgery, Patan
Hospital, Patan Academy of
Health Sciences, Lalitpur, Nepal
Email:
skgajendramani65@gmail.com

Peer Reviewers

Prof. Dr. Jay N Shah
Patan Hospital, PAHS

Asst. Prof. Dr. Ashis Shrestha
Patan Hospital, PAHS

Outcome of anterior cruciate ligament reconstruction using bone-patellar tendon-bone auto graft

Gajendra Mani Shah,¹ Nabees Man Singh Pradhan,² Rojan Tamrakar,¹ Bidur Gyawali,¹ Toya Raj Bhatta,¹ Balakrishnan M Acharya,³ Sagun Basnyat,¹ Suman Shrestha²

¹Lecturer, ²Professor, ³Associate Professor, Department of Orthopaedics and Trauma Surgery, Patan Hospital, Academy of Health Sciences, Lalitpur, Nepal

ABSTRACT

Introductions: A rupture of the Anterior Cruciate Ligament (ACL) is the most common ligament injury resulting in instability of the knee which can cause secondary articular injury and early osteoarthritis (OA). Aim of this study was to evaluate the short term outcomes of ACL reconstruction using Bone Patellar Tendon Bone Auto Graft (BPTB) auto graft and factors that might contribute to poor results.

Methods: Patients with Anterior Cruciate Ligament injury during the period of July 2013 to May 2014 were enrolled to observe the outcome of the ACL reconstruction using BPTB without arthroscopy. Lysholm Knee Scoring Scale was used to determine the clinical outcome. Outcome parameters were evaluated using Statistical Package for Social Science Version 20 (SPSS).

Results: Total 30 normal or near normal function outcome of knee was reported in 29 (97%) patients after ACL reconstruction using BPTB without arthroscopy. There was no failure of grafts. Good static knee stability achieved with increased patient's satisfaction. Lysholm Score was excellent in 27 (90%) and good in 3 (10 %) cases. Anterior knee pain persisted in 6 cases (20%).

Conclusions: Excellent functional outcome with BPTB auto graft in Anterior Cruciate Ligament injury was achieved.

Keywords: ACL reconstruction, bone-patellar tendon-bone auto graft, clinical outcome

INTRODUCTIONS

Up to 300,000 Anterior Cruciate Ligament (ACL) reconstructions are performed annually in the United States. An estimated 80% of these reconstructions are done with auto grafts, with the remainder being performed with allograft from various sources.¹ An ideal graft would replicate the anatomy and biomechanics of the native ACL, with rapid incorporation and low donor site morbidity.²⁻⁴ This study was conducted to evaluate the short term outcomes of ACL reconstruction using BPTB auto graft and associated factors that might contribute to poor results.

METHODS

A prospective observational study was done in patients with ACL injury during the period of July 2013 to May 2014 to observe the outcome using BPTB reconstruction without arthroscopy at Patan Hospital, Lagankhel, Lalitpur; Shree Birendra Hospital, Chhauni and NAMS, Bir Hospital, Kathmandu from July 2013 to May 2014. Lysholm Knee Scoring Scale⁴ was used to determine the clinical outcome.

Ethical clearance was taken from the Institutional Review Board of NAMS. Written consent was obtained from all the patients. Inclusion criteria was age 20 to 50 years with positive Lachman test, pivot-shift test, and closed or nearly closed growth plates based on radiographs. Excluded from the study were: open knee injury and associated lower limb, pelvic or spinal fracture with vascular injuries; neurological involvement and compartment syndrome; bilateral ACL injuries; prior ACL reconstruction of the knee requiring a revision procedure; associated

posterior cruciate ligament (PCL) tears or multiple-ligament injuries; pre-existing deformity in either of the knee; spinal disorders with neurological deficits and associated osteoarthritic changes.

Functional outcome was assessed 3, 6, and 9 months after the surgery with measurement of Lysholm Knee Scoring Scale⁴ knee flexion and extension; patients satisfaction by visual analogue score (VAS) score; and anterior laxity was assessed by anterior drawer test, Lachman test and pivot shift test.

The SPSS version 20 was used for data analysis. Parametric data was analyzed using student t-test and nonparametric data by Chi-square. Confidence interval was set to be 95% and p value <0.05 was considered significant.

RESULTS

There were 30 patients, with mean age 30.93 ± 6.23 years, majority 20-30 years and right knee involvement. The mean knee flexion preoperatively and postoperatively at 3, 6 and 9 months improved to near normal, (Table 1). The means flexion lag was 1.5 degree compared to normal side and mean extension lag was 1.72 degree.

There was no statistically significant difference in mean value of preoperative flexion and extension ($P > 0.05$). Lysholm Knee Score at 9 months was excellent (>90) in 22 cases, good (84-90) in 8, and none in fair (65-83) or poor (<65) category.

The means of preoperative flexion and extension was statistically not significant ($P > 0.05$), (Table 2).

Table 1. Knee Flexion and extension before and after ACL reconstruction using BPTB without arthroscopy (n=30)

Mean \pm SD	Normal Side	Preoperatively	3 Months	6 Months	9 Months
Flexion (degrees)	131.5 \pm 23	130.5 \pm 9	104.80 \pm 5.57	118.73 \pm 3.39	130 \pm 1.01
Extension (degrees)	8.9 \pm 0.6	4.90 \pm 0.76	5.17 \pm 0.59	6.40 \pm 0.56	7.18 \pm 0.51

Table 2. Mean knee Flexion/Extension comparison between preoperatively and postoperatively (9 Months)

	Preoperatively	Postoperatively (9 Months)	Std Deviation	P
Mean Knee Flexion (degrees)	130.5	130	1.01	>0.05
Mean Knee Extension (degrees)	4.90	7.18	0.59	>0.05

Table 3. Lysholm Knee Score before and after ACL reconstruction using BPTB without arthroscopy (n=30)

	Preoperatively	At 3 Months	At 6 Months	At 9 Months
Mean Lysholm Knee Score \pm SD	62.50 \pm 7.13	56.43 \pm 6.13	81.36 \pm 6.80	91.70 \pm 2.79

Table 4. Mean VAS Score after ACL reconstruction using BPTB without arthroscopy (n=30)

	3 Months	6 Months	9 Months
Mean VAS Score \pm SD	3.66 \pm 0.76	1.53 \pm 0.50	0.30 \pm 0.46

Preoperatively anterior drawer test, Lachman test and pivot shift test were positive for all cases but postoperatively at 9 months follow up, the anterior drawer test and Lachman test significantly improved and pivot shift test was negative in all cases.

DISCUSSIONS

In our study, there was male predominance with the ratio of 4:1, similar to Lars Ejerhed, MD (male 70%),⁵ Taylor DC⁶ (M:F 4.8:1) and Mountcastle⁷ (M:F 3.3:1). The male predominance in most of the study may be because male is more active and frequently involved in outdoor activities and contact sports.

Our mean flexion lag of 1.5 degree was better than other studies. Dong Jiang⁸ reported mean flexion lag 2 degree in double blind group and 2.5 in single blind group. In a similar study by Kanniraj⁹ average flexion loss was <5 degree in 92% and <15 degree in 8%.

The average extension loss in our study was 1.72 degree, similar to Dong Jiang⁸ (0.8 degree in double blind group and 0.9 in single blind group), Kanniraj⁹ <3 degree in 76 (96%) and 3 to 5 degree in 3 (4%).

Anterior laxity of joints assessed by anterior drawer test, Lachman test and pivot shift test were positive for all cases preoperatively but

postoperatively at 9 months all tests significantly improved.

In a study by Shuzhen Li¹⁰ the postoperative Lachman test was negative in 44 out of 78 patients in the BPTB group and postoperative pivot test negative in 158 out of 179.

Similar study conducted by Matthew, 80.8% (autograft) and 85.1% (allograft) had negative pivot shift test.¹¹ But in our study at 9 months follow up anterior drawer test and Lachman test was significantly improved and pivot shift test was negative in all cases.

In our study, preoperative mean Lysholm Knee Score was 62.50 \pm 7.13 degrees and postoperative at 3 months 56.43 \pm 6.13, 6 months 81.36 \pm 6.80 and 9 months 91.70 \pm 2.79. Similarly, Lysholm Knee Score at 9 months was excellent (>90) in 22 cases, good (84-90) in 8 cases. In similar study by Kanniraj⁹ the mean Lysholm knee score improved from 48 (range, 24–77) to 92 (range, 81–96) (p<0.0001); scores were excellent (>90) in 46 (58%) patients and good (83-90) in 32 (40.50%) patients.

In a study by Lars Ejerhed⁵ the mean Lysholm Knee Score preoperative was 70 (14-95) and postoperatively it was 95 which is similar to our study. Dong Jiang⁸ too reported similar findings.

In our study patient satisfaction mean VAS improved (Table 4), similar to Matthew.¹¹

The risk of anterior knee pain, patellar tendon rupture and patellar fracture is higher¹² but we did not have such complications of BPTB auto graft at 9 months follow up. Similarly, we did not have any post-operative infection and graft failure.

Present study has limitations of small sample size and shorter follow up. We analyzed only Lysholm Knee Scoring Scale. Inclusion of objective parameters like quadriceps strength, hamstring strength may have added value to the analysis.

CONCLUSIONS

Excellent clinical outcome was achieved in anterior cruciate ligament (ACL) reconstruction with bone-patellar tendon-bone (BPTB) autograft without arthroscopy.

REFERENCES

- Cohen SB, Sekiya JK. Allograft safety in anterior cruciate ligament reconstruction. *Clin Sports Med.* 2007;26(4):597-605. DOI: <https://doi.org/10.1016/j.csm.2007.06.003>
- Baer GS, Harner CD. Clinical outcomes of allograft versus autograft in anterior cruciate ligament reconstruction. *Clin Sports Med.* 2007;26(4):661-81. DOI: <https://doi.org/10.1016/j.csm.2007.06.010>
- Aglietti P, Giron F, Cuomo P, Losco M, Mondanelli N. Single- and double-incision double-bundle ACL reconstruction. *Clin Orthop Relat Res.* 2007;454:108-13. DOI: [10.1097/BLO.0b013e31802baaf4](https://doi.org/10.1097/BLO.0b013e31802baaf4) PMID: [17202919](https://pubmed.ncbi.nlm.nih.gov/17202919/)
- Bengtsson J, Möllborg J, Werner S. A study for testing the sensitivity and reliability of the Lysholm knee scoring scale. *Knee Surg Sports Traumatol Arthroscopy.* 1996;4(1):27-31. PMID: [8819060](https://pubmed.ncbi.nlm.nih.gov/8819060/). DOI: <https://doi.org/10.1007/BF01565994>
- Ejerhed L, Kartus J, Sernert N, et al. Patellar tendon or semitendinosus tendon autografts for anterior cruciate ligament reconstruction? a prospective randomized study with a two-year follow-up. *The American Journal of Sports Medicine.* 2003;31(1):19-25. DOI: <https://doi.org/10.1177/03635465030310011401>
- Taylor DC, Posner M, Curl WW, Feagin JA: Isolated tears of the anterior cruciate ligament: over 30-year follow-up of patients treated with arthrotomy and primary repair. *Am J Sports Med.* 2009, 37:65–71. PMID: [19029313](https://pubmed.ncbi.nlm.nih.gov/19029313/) DOI: [10.1177/0363546508325660](https://doi.org/10.1177/0363546508325660)
- Mountcastle SB, Posner M, Kragh JF, et al. Gender differences in anterior cruciate ligament injury vary with activity: epidemiology of anterior cruciate ligament injuries in a young, athletic population. *Am J Sports Med.* 2007;35:1635–42. PMID: [17519438](https://pubmed.ncbi.nlm.nih.gov/17519438/) DOI: [10.1177/0363546507302917](https://doi.org/10.1177/0363546507302917)
- Jiang D, Ao YF, Gong X, Wang YJ, Luo H, Chen LX, Wang HJ, Xie X, Zhang JY, Yu JK. Double-Bundle Anterior Cruciate Ligament Reconstruction Using Bone–Patellar Tendon–Bone Allograft: Technique and 2-to 5-Year Follow-up. *The American journal of sports medicine.* 2012 May;40(5):1084-92. PMID: [22472270](https://pubmed.ncbi.nlm.nih.gov/22472270/) DOI: [10.1177/0363546512440686](https://doi.org/10.1177/0363546512440686)
- Kanniraj Marimuthu, Narendra Joshi, Mrinal Sharma, Chandra Sekar Sharma, Rakesh Bhargava, Ajay Singh Rajawat, Sushil Subash Rangdal. Anterior cruciate ligament reconstruction using the medial third of the patellar tendon. *Journal of Orthopaedic Surgery.* 2011;19(2):221-5. PMID: [21857050](https://pubmed.ncbi.nlm.nih.gov/21857050/) DOI: [10.1177/230949901101900219](https://doi.org/10.1177/230949901101900219)
- Shuzhen Li, Yueping Chen, Zonghan Lin, Wei Cui, Jingmin Zhao, Wei Su. A systematic review of randomized controlled clinical trials comparing hamstring autografts versus bone-patellar tendon-bone autografts for the reconstruction of the anterior cruciate ligament. *Arch Orthop Trauma Surg.* 2012;132:1287–97. PMID: [22661336](https://pubmed.ncbi.nlm.nih.gov/22661336/) DOI: [10.1007/s00402-012-1532-5](https://doi.org/10.1007/s00402-012-1532-5)
- Matthew J. Kraeutler, Jonathan T. Bravman, Eric C. McCarty. Bone–Patellar Tendon–Bone Autograft Versus Allograft in Outcomes of Anterior Cruciate Ligament Reconstruction : A Meta-analysis of 5182 Patients. *Am J Sports Med.* 2013;41(10):2439-48. PMID: [23585484](https://pubmed.ncbi.nlm.nih.gov/23585484/) DOI: [10.1177/0363546513484127](https://doi.org/10.1177/0363546513484127)
- Xie X., Xiao Z., Li Q. Increased incidence of osteoarthritis of knee joint after ACL reconstruction with bone-patellar tendon-bone autografts than hamstring autografts: A meta-analysis of 1,443 patients at a minimum of 5 years. *Eur J Orthop Surg Traumatol.* 2015;25:149–59. PMID: [24748500](https://pubmed.ncbi.nlm.nih.gov/24748500/). DOI: [10.1007/s00590-014-1459-3](https://doi.org/10.1007/s00590-014-1459-3)