

1989

Geologic Evolution of Trail Ridge Eolian Heavy-Mineral Sand and Underlying Peat, Northern Florida

Eric Force
U.S. Geological Survey

Fredrick J. Rich
Georgia Southern University, frich@georgiasouthern.edu

Follow this and additional works at: <https://digitalcommons.georgiasouthern.edu/geo-facpubs>



Part of the [Geography Commons](#), and the [Geology Commons](#)

Recommended Citation

Force, Eric, Fredrick J. Rich. 1989. "Geologic Evolution of Trail Ridge Eolian Heavy-Mineral Sand and Underlying Peat, Northern Florida." *U.S. Geological Survey Professional Paper 1499*: 1-23 Washington, DC: United States Government Printing Office. source: <https://pubs.er.usgs.gov/publication/pp1499>
<https://digitalcommons.georgiasouthern.edu/geo-facpubs/117>

This article is brought to you for free and open access by the Geology and Geography, Department of at Digital Commons@Georgia Southern. It has been accepted for inclusion in Geology and Geography Faculty Publications by an authorized administrator of Digital Commons@Georgia Southern. For more information, please contact digitalcommons@georgiasouthern.edu.

Geologic Evolution of Trail Ridge Eolian Heavy-Mineral Sand and Underlying Peat, Northern Florida

U.S. GEOLOGICAL SURVEY PROFESSIONAL PAPER 1499



AVAILABILITY OF BOOKS AND MAPS OF THE U.S. GEOLOGICAL SURVEY

Instructions on ordering publications of the U.S. Geological Survey, along with prices of the last offerings, are given in the current-year issues of the monthly catalog "New Publications of the U.S. Geological Survey." Prices of available U.S. Geological Survey publications released prior to the current year are listed in the most recent annual "Price and Availability List." Publications that are listed in various U.S. Geological Survey catalogs (see back inside cover) but not listed in the most recent annual "Price and Availability List" are no longer available.

Prices of reports released to the open files are given in the listing "U.S. Geological Survey Open-File Reports," updated monthly, which is for sale in microfiche from the U.S. Geological Survey, Books and Open-File Reports Section, Federal Center, Box 25425, Denver, CO 80225. Reports released through the NTIS may be obtained by writing to the National Technical Information Service, U.S. Department of Commerce, Springfield, VA 22161; please include NTIS report number with inquiry.

Order U.S. Geological Survey publications by mail or over the counter from the offices given below.

BY MAIL

Books

Professional Papers, Bulletins, Water-Supply Papers, Techniques of Water-Resources Investigations, Circulars, publications of general interest (such as leaflets, pamphlets, booklets), single copies of Earthquakes & Volcanoes, Preliminary Determination of Epicenters, and some miscellaneous reports, including some of the foregoing series that have gone out of print at the Superintendent of Documents, are obtainable by mail from

**U.S. Geological Survey, Books and Open-File Reports
Federal Center, Box 25425
Denver, CO 80225**

Subscriptions to periodicals (Earthquakes & Volcanoes and Preliminary Determination of Epicenters) can be obtained ONLY from the

**Superintendent of Documents
Government Printing Office
Washington, D.C. 20402**

(Check or money order must be payable to Superintendent of Documents.)

Maps

For maps, address mail orders to

**U.S. Geological Survey, Map Distribution
Federal Center, Box 25286
Denver, CO 80225**

Residents of Alaska may order maps from

**Alaska Distribution Section, U.S. Geological Survey,
New Federal Building - Box 12
101 Twelfth Ave., Fairbanks, AK 99701**

OVER THE COUNTER

Books

Books of the U.S. Geological Survey are available over the counter at the following Geological Survey Public Inquiries Offices, all of which are authorized agents of the Superintendent of Documents:

- WASHINGTON, D.C.--Main Interior Bldg., 2600 corridor, 18th and C Sts., NW.
- DENVER, Colorado--Federal Bldg., Rm. 169, 1961 Stout St.
- LOS ANGELES, California--Federal Bldg., Rm. 7638, 300 N. Los Angeles St.
- MENLO PARK, California--Bldg. 3 (Stop 533), Rm. 3128, 345 Middlefield Rd.
- RESTON, Virginia--503 National Center, Rm. 1C402, 12201 Sunrise Valley Dr.
- SALT LAKE CITY, Utah--Federal Bldg., Rm. 8105, 125 South State St.
- SAN FRANCISCO, California--Customhouse, Rm. 504, 555 Battery St.
- SPOKANE, Washington--U.S. Courthouse, Rm. 678, West 920 Riverside Ave.
- ANCHORAGE, Alaska--Rm. 101, 4230 University Dr.
- ANCHORAGE, Alaska--Federal Bldg., Rm. E-146, 701 C St.

Maps

Maps may be purchased over the counter at the U.S. Geological Survey offices where books are sold (all addresses in above list) and at the following Geological Survey offices:


- ROLLA, Missouri--1400 Independence Rd.
- DENVER, Colorado--Map Distribution, Bldg. 810, Federal Center
- FAIRBANKS, Alaska--New Federal Bldg., 101 Twelfth Ave.

Geologic Evolution of Trail Ridge Eolian Heavy-Mineral Sand and Underlying Peat, Northern Florida

By ERIC R. FORCE *and* FREDRICK J. RICH

U.S. GEOLOGICAL SURVEY PROFESSIONAL PAPER 1499

The peat underlying ilmenite ore sand is a synchronous facies and records the approach of the eolian dune



DEPARTMENT OF THE INTERIOR

MANUEL LUJAN, JR., *Secretary*

U.S. GEOLOGICAL SURVEY

Dallas L. Peck, *Director*

Any use of trade, product, or firm names in this publication is for
descriptive purposes only and does not imply endorsement by the
U.S. Government

Library of Congress Cataloging in Publication Data

Force, Eric R.

Geologic evolution of Trail Ridge eolian heavy-mineral sand and underlying peat, northern Florida.

(U.S. Geological Survey professional paper ; 1499)

Bibliography: p.

Supt. of Docs. no.: I 19.16: 1499

1. Heavy minerals—Florida—Trail Ridge. 2. Sand—Florida—Trail Ridge. 3. Peat—Florida—Trail Ridge. 4. Eolian processes—Florida—Trail Ridge. 5. Geology—Florida—Trail Ridge. I. Rich, Fredrick J. II. Title. III. Series.

QE364.2.H4F67

1989

553.4'6

88-607931

For sale by the Books and Open-File Reports Section, U.S. Geological Survey,
Federal Center, Box 25425, Denver, CO 80225

CONTENTS

| | Page | | Page |
|----------------------------------------------------------|------|-----------------------------------------------------------|------|
| Abstract..... | 1 | Nature of Trail Ridge Lignitic Peat | 7 |
| Introduction..... | 1 | General description | 7 |
| Previous work..... | 2 | Petrography | 8 |
| Nature of the Trail Ridge ilmenite ore sand | 3 | Palynology..... | 9 |
| Areal distribution and geologic relations | 3 | Age | 10 |
| Sequence and structure | 4 | The sand component of Trail Ridge peat..... | 10 |
| Sand morphology and grain size | 5 | Depositional relation of Trail Ridge peat and sands | 14 |
| Heavy minerals | 6 | Paleogeographic reconstruction | 15 |
| Heavy-mineral variation with sedimentary structure | 6 | Acknowledgments | 15 |
| Origin | 7 | References | 15 |

ILLUSTRATIONS

| | Page |
|------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| FIGURE 1. Location of Trail Ridge shown on a satellite thermal image | 2 |
| 2. Cross-sectional profile of Trail Ridge..... | 2 |
| 3. Diagrams showing lithologic features, percent leucoxenized ilmenite, and percent TiO ₂ in altered ilmenite, Trail Ridge ore body | 3 |
| 4. Photograph showing humate pseudobedding in Trail Ridge sand..... | 4 |
| 5. Photograph showing Trail Ridge ilmenite ore sand | 5 |
| 6. Circular diagram of azimuths and magnitude of crossbed dip in Trail Ridge ilmenite ore sand | 5 |
| 7. Topographic map showing Trail Ridge localities discussed in the text | 6 |
| 8. Scanning electron microscope photographs of grains from the Trail Ridge ore sand..... | 7 |
| 9. Photomicrograph of thin section of Trail Ridge ilmenite ore sand | 8 |
| 10. Photograph of laminae alternately rich and poor in heavy minerals | 8 |
| 11. Graphs comparing adjacent laminae rich and poor in heavy minerals | 9 |
| 12. Graphs showing palynological and maceral constituents of peat in core RTR2 | 11 |
| 13. Photomicrograph of thin section of sandy lignitic peat from core RTR2 | 12 |
| 14. Scanning electron microscope photograph of sand-bearing lignitic peat sample from core RTR2 | 12 |
| 15. Graphs showing grain-size distributions of inorganic fraction of sandy Trail Ridge peat from core RTR2..... | 13 |
| 16. Scanning electron microscope photographs of sand grains in Trail Ridge peat from core RTR2 | 13 |
| 17. Photograph showing Holocene coastal transgressive longwall dune near Newcastle, New South Wales, Australia..... | 14 |
| 18. Schematic paleogeography of Trail Ridge at the time of dune formation..... | 15 |

TABLE

| | Page |
|------------------------------------------------------------------------------------|------|
| TABLE 1. Organic and sand constituents of core samples from Trail Ridge peat | 10 |

GEOLOGIC EVOLUTION OF TRAIL RIDGE EOLIAN HEAVY-MINERAL SAND AND UNDERLYING PEAT, NORTHERN FLORIDA

By Eric R. FORCE and Fredrick J. RICH

ABSTRACT

The southern portion of Trail Ridge in Clay County, Fla., has been an important domestic source of altered ilmenite, zircon, and other minerals since 1949. The ridge as a physiographic feature extends in a north-south direction over 160 km and is 1 to 2 km wide. The ore is a fine- to medium-grained sand body that extends down from the crest of the ridge about 20 m. Average heavy-mineral content is about 4 percent. Over half of the ilmenite and zircon is contained in thin, dark laminae dipping 26° to 41° southwest; these laminae extend deep within the ore and outline the slip faces of a great eolian dune complex. The subordinate dark laminae differ in heavy mineralogy and grain size from the encasing light-colored sand. The dark laminae have a modal grain size of 0.2 mm and average about 6 percent heavy minerals, among which altered ilmenite, zircon, and rutile predominate. The light-colored laminae have a coarser modal grain size of 0.3 mm and average only 1 percent heavy minerals, among which the lighter heavy minerals staurolite, sillimanite, and tourmaline form half. Grain size variables and grain surfaces are appropriate for eolian sand. Overprinted on these original features of the dune are a surficial weathering zone over 3 m thick, where tan leucoxene takes the place of black altered ilmenite, and several underlying humate-cemented zones that probably represent water-table stillstands.

The immediately underlying unit is a lignitic peaty layer 1.5 m thick, here referred to as peat. The organic fraction is derived entirely from freshwater plants. In-place tree stumps have been noted, but the predominant component of the peat is fragments of wood and other transported plant debris. The peat and its constituents indicate deposition in a swamp environment, and local horizons enriched with charcoal and fungal remains indicate periodic subaerial exposure. Vegetation varied from open shrub swamp to cypress forest.

The age of the peat has been determined palynologically as post-Miocene. Its carbon-14 age is greater than 4.5×10^4 years; that is, pre-latest Pleistocene.

The upper portion of the peat layer contains admixtures of sand. This sand is present as isolated grains embedded in laminated organic matrix and is dominantly well rounded and frosted. As a grain population, the sand in the peat matches the overlying Trail Ridge ore sand in mineralogy and grain morphology but is slightly finer in grain size. This sand we regard as an important clue to the history of the area. The sand was apparently deposited from aerial suspension, and its characteristics were acquired in the adjacent high-energy eolian environment. Upward increase of sand in peat records the approach of the dune that eventually prograded the swamp. The peat and the

overlying sand are essentially the same age. Fine sand found in peat represents the sand fraction remaining suspended in flow separation at the top of the slip face of the dune. Ore represents the traction-load fraction. The Trail Ridge dune itself is probably the drainage dam that impounded the swamp it later overrode.

The sand embedded in peat is also an important clue to the weathering history of Trail Ridge heavy minerals, as entombment in peat probably arrested oxidation. The presence of leucoxene, and other features of the mineral assemblage, shows that the minerals were already weathered at deposition. This evidence is in accord with grain-surface features and grain-size-density relations. Thus, mineral alteration at Trail Ridge occurred in two stages, one before and one after deposition.

Trail Ridge apparently represents a coast-parallel transgressive dune complex, analogous to younger dunes elsewhere that have become completely decoupled from parental shorelines. The Trail Ridge body was probably composite, made up of individual parabolic dunes, each migrating southwestward. The base of the Trail Ridge body was probably originally higher at the southern end than at the northern end.

INTRODUCTION

Trail Ridge is a linear north-south-trending ridge about 1 to 2 km wide, which extends at least 160 km from northeastern Florida to southeastern Georgia. It is about 65 km inland from the coast, and its crest increases in elevation from about 46 m (150 ft) in the north to as much as 77 m (252 ft) in the south. Because the ridge typically stands tens of meters above adjacent areas and impounds the Okefenokee Swamp, it is one of the most conspicuous linear landforms in the coastal plain of the Southeastern States (fig. 1). A profile is shown in figure 2.

The ridge is composed of a sorted fine-grained to medium-grained quartzose sand body up to 20 m thick. In a 29-km portion of its length in Florida, the sand throughout its thickness averages about 4 percent heavy minerals,¹ predominantly altered ilmenite, and lesser zircon, rutile, and aluminosilicates. This sand is, or was

¹ All listed mineral percentages are by weight, unless otherwise specified.




FIGURE 1.—Location of Trail Ridge shown on a satellite thermal image.




FIGURE 2.—Approximate cross-sectional profile of Trail Ridge, with vertical exaggeration of 200. Shaded bodies are heavy-mineral sands. Line of section A-A' is shown in figure 1.

before mining, probably the most valuable single resource of titanium minerals in the United States.

Since 1949, E.I. Du Pont de Nemours and Company has mined this body in Clay County, Fla. Exploration holes and cores in this area have shown that the central to western portions of the south end of the Trail Ridge

sand body rest on about 1.5 m of lignitic peat that typically contains large wood fragments. In this paper we examine the relation between the lignitic peat and the overlying sand body (henceforth referred to as ilmenite ore sand) to clarify their original paleogeographic relation and their origin.

PREVIOUS WORK

The earliest geologic descriptions of Trail Ridge were entirely geomorphic. Cooke (1925, 1939) envisioned the ridge as a large Pleistocene marine sand spit that was attached to a mainland near Jesup, Ga. (Cooke, 1925, plate Xb). Cooke believed that the sand body possibly was tilted northward after deposition and that the sand was partly eolian in origin. MacNeil (1950) extended this view to include the Okefenokee Swamp as a former lagoon behind a Trail Ridge marine shoreline and bar.

By the mid 1940's, Florida was the site of several small heavy-mineral mining operations. Discovery of economic heavy minerals at Trail Ridge coincided with the first descriptions of its sands. Spencer (1948) reported on a U.S. Bureau of Mines drilling program on Trail Ridge that encountered valuable deposits of altered ilmenite, zircon, and rutile in sand containing 3 to 10 percent heavy minerals to depths of over 10 m. A heavy-mineral assemblage of altered ilmenite, zircon, rutile, staurolite, tourmaline, sillimanite, kyanite, corundum, monazite, and spinel was described, as was the nature of ilmenite alteration. Credit for the discovery is shared by Spencer's team and the Du Pont company, under J.L. Gillson. Thoenen and Warne (1949) extended the drilling investigations along and adjacent to Trail Ridge by using the newly developed jet drill.

The Trail Ridge mine was opened by Du Pont near the southern end of the ridge in 1949. At first the plant was operated by Humphrey Gold Corporation. The operation was described in several mining journals in the early 1950's. The most thorough such description, by Carpenter and others (1953), was the first to describe the humate-cemented "hardpan" in the ilmenite ore sand and the woody peat layer underlying it. They found wood fragments, up to 5 cm in diameter, and pinecones in this peat layer. They reported the thickness of the peat to be about 1.5 m and that of the overlying ilmenite ore sand to be up to 20 m. Average heavy-mineral grade was listed as 4 percent.

Recognition by Creitz and McVay (1948) that Trail Ridge ilmenite is altered to an iron-deficient, less-dense compound containing microcrystalline rutile, and by Cannon (1950) that the heaviest ilmenite alteration at Trail Ridge occurs above the present water table, spawned many worldwide comparative studies from 1954




FIGURE 3.—Leucoxene zonation in the Trail Ridge ore body. Percent leucoxenized ilmenite is calculated as a moving average from E.C. Pirkle and Yoho (1970, cores TR 1 and 2); TiO₂ in altered ilmenite is from Temple (1966).

to 1966 of ilmenite alteration in sand matrix. Of these studies, the one by Temple (1966) is most concerned with Trail Ridge per se. Garnar (1972) showed the chemical alteration path for Trail Ridge ilmenite.

The papers by E.C. Pirkle and Yoho (1970) and E.C. Pirkle and others (1970) constitute the first thorough descriptions of the deposit and emphasize the vertical sequence shown in two cores. They present the first records of (1) the remarkably consistent grain size of the deposit, the median value being about 0.3 mm, (2) sedimentary structures and other original depositional features in the ore sand, and (3) configuration of the base of the ore body and the relation of ore sand to peats and other underlying lithologies. The "Trail Ridge Sequence" of these authors consists of the ore sand and underlying peat. Figure 3 shows their data on leucoxene zonation within the ore body. They hypothesized that the ore sand was deposited in a complex of beach ridges and associated eolian dunes.

Subsequent geologic publications on Trail Ridge have continued to be almost entirely by the Pirkle family and their colleagues. E.C. Pirkle and others (1971) described an additional core through Trail Ridge, adjacent to the Florida-Georgia line. F.L. Pirkle (1975) compared grain sizes, heavy-mineral suites, and grain shapes among Trail Ridge sands and proposed possible source regions. E.C. Pirkle and others (1977) described the portion of the Trail Ridge deposit northwest of Highway 301 in Florida. F.L. Pirkle and Czel (1983) reported marine fossils of probable Pleistocene age at elevations of 39 to 49 m (130 to 160 feet) in sands just west of and stratigraphically above a Georgia segment of

the ridge. W.A. Pirkle and E.C. Pirkle (1984) and F.L. Pirkle (1984) summarized the evidence for a beach-ridge origin of Trail Ridge, the latter paper concluding that some grain-size characteristics of the deposit are consistent only with an eolian origin.

The most recent published work is by the authors. Rich (1985) described the palynology of the peat bed, and Force and Garnar (1985) described eolian crossbedding of the ilmenite ore sand.

NATURE OF THE TRAIL RIDGE ILMENITE ORE SAND

AREAL DISTRIBUTION AND GEOLOGIC RELATIONS

The nature of the sand composing Trail Ridge is well documented only in the portion having high heavy-mineral contents, which is at the southern end of the body, in Clay and Bradford Counties, Fla. Even there the areal distribution and geologic relations of the sand body are not adequately reported in the literature. Most of our knowledge comes from descriptions of the ilmenite ore sand in profile, in substantial part from four cored drill holes. The mining operations only rarely expose the main part of the ore body. Unpublished company reports, to which we have limited access, constitute the largest body of information that can be used to define the geometry of this sand body.

At the southern end of the ridge near Starke, Fla., the central and perhaps western parts of the sand body are underlain by lignitic peat. The nature of the contact is a



FIGURE 4.—Humate pseudobedding in Trail Ridge sand, photographed in wall of dredge pond, in interval 1.6 to 2.6 m below land surface.

major focus of this paper. The contact surface apparently dips about 3 m/km (15 feet per mile) to the west-southwest (E.C. Pirkle and Yoho, 1970, fig. 3). Underlying the peat are the "post-Hawthorn clastics" of E.C. Pirkle and others (1970), consisting in part of marine clay. North of this area, the sand body is underlain by fine-grained sands and (or) clays that locally contain wood (root?) fragments and show mineralogical indications of subaerial weathering (E.C. Pirkle and others, 1977).

The ilmenite ore sand extends from this subhorizontal base to the ground surface. Thus the ore is thickest where the elevation of the ridge is greatest (fig. 2). Maximum thickness is about 20 m.

SEQUENCE AND STRUCTURE

The ore is more or less homogeneous except for a rootlet-mottled leached zone extending down 2 to 3 m from the ground surface and grading downward into a variably indurated humate-cemented zone. Humate induration increases irregularly downward to a hardpan that can make a hammer ring. Several humate-indurated horizons occur in some localities, to as much as 6.5 m

below the land surface, forming pseudobedding (figs. 4, 5) overprinted on weathered sand. This humate cement must represent precipitation at the present and former water tables. Below 6.5 m, humate cementation is weak.

The surficial leached zone and the upper part of the humate-cemented zone coincide approximately with the interval in which leucoxene is enriched at the expense of less-altered ilmenite (fig. 3).

Force and Garnar (1985) reported that the weakly cemented ilmenite ore sand underlying the humate-cemented zone shows high-angle eolian crossbedding outlined by heavy-mineral laminae dipping southwest at 26° to 41° (fig. 6). This underlying zone was exposed during a 1984 lowering of the dredge pond level at the Trail Ridge mine (figs. 5,7). A 1968 lowering had also exposed such crossbeds (E.C. Pirkle and Yoho, 1970, fig. 6; Force and Garnar, 1985, fig. 2), and a core shows the crossbeds from a deeper interval (E.C. Pirkle and others, 1970, p. 39, PTR1). These three localities represent a relatively small area (fig. 7), so it is not yet possible to define the areal extent of eolian deposits. All three localities are in the area where the sand body is underlain by peat.



FIGURE 5.—Trail Ridge ilmenite ore sand exposed in 1984. The basal interval (about 0.9 m thick) is eolian sand with southwest-dipping crossbeds, overprinted by thin humate-cemented zones. Overlying lithologies are weakly cemented sand with crossbedding still visible (0.6 m thick), massive resistant humate-cemented sand (1.2 m thick), and at the top a rootlet-mottled bleached zone (the remainder). Location is shown on map in figure 7.

SAND MORPHOLOGY AND GRAIN SIZE

The ilmenite ore sand is remarkably homogeneous in the four profiles that have been described from cores (E.C. Pirkle and Yoho, 1970; E.C. Pirkle and others, 1970, 1977). This homogeneity is understandable in the context of an eolian origin and deposition on steep slip faces. Median grain size of the sand ranges only from 0.24 to 0.38 mm, and the distribution of median grain size values within the sand body appears random.

Skewness values for grain-size distribution of the sands have never been published. Our calculations (based on samples of E.C. Pirkle and Yoho, 1970; E.C. Pirkle and others 1970, 1977) show the sands that are low in postdepositional clay give positive values, consistent with Friedman's (1961) conclusions for eolian sand.




FIGURE 6.—Stereographic projection of azimuths and magnitude of crossbed dip in Trail Ridge ilmenite ore sand.

These positive values persist to the base of the ilmenite ore sand.

Grain sorting by size is good. Sorting coefficients listed by E.C. Pirkle and Yoho (1970) for the ilmenite ore sand in the two southern drill holes range only from 1.16 to 1.32, again with irregular vertical distribution.² There is no apparent relation to median grain size. These sorting coefficients indicate the sands to be well sorted, in the terminology of Trask (1932).

The sands are well rounded also, though this description has never been rigorously quantified. However, F.L. Pirkle (1975) presents average axial ratios of intermediate and maximum axes of zircon and aluminosilicates from one size fraction of sands from three of the drill holes. This measure of sphericity ranges from 0.64 to 0.68.

Quartz sand grains from the Trail Ridge ilmenite ore sand are somewhat frosted, and grains of altered ilmenite show a polish. Figure 8 shows scanning electron microscope photographs of quartz and ilmenite, collected in 1984 from a crossbedded sand exposed below the lowest humate hardpan layer. The surfaces are analogous to those described elsewhere for eolian sands (for example, Krinsley and Margolis, 1971).

² Our calculations from E.C. Pirkle and Yoho (1970) data give a range of 1.23 to 1.35. Apparently their values are for sand fractions, free of postdepositional matrix.




FIGURE 7.—Part of the Starke quadrangle map (1949 edition) illustrating Trail Ridge localities discussed in the text. PTR localities refer to E.C. Pirkle and Yoho (1970); RTR localities refer to Rich (1985).

HEAVY MINERALS

The heavy-mineral content of the ilmenite ore sand varies widely even in the interval samples analyzed by E.C. Pirkle and Yoho (1970) and E.C. Pirkle and others (1970, 1977). The reported heavy-mineral content ranges from less than 1 percent to 16.29 percent (in tetrabromoethane). Variation is high even within a single core. Among the drill holes, average heavy-mineral content of

the body varies from 6 percent toward the south to 3 percent to the north.

Average percentages of the altered ilmenite and leucoxene in the heavy-mineral assemblage have been published by E.C. Pirkle and Yoho (1970) and E.C. Pirkle and others (1970, 1977). These minerals together range from 47 to 64 percent in the northern part of the body and from 16 to 67 percent toward the south. Values below about 50 percent are mostly restricted to the uppermost parts of the ilmenite ore sand; that is, the same zone in which leucoxene has formed at the expense of less-altered ilmenite.³

The remainder of the heavy ($\rho > 2.85$) suite at Trail Ridge consists on the average of about 15 percent zircon, 2 to 3 percent rutile, 15 percent staurolite, 3 percent kyanite, 5 to 6 percent sillimanite, 5 percent tourmaline, and minor amounts of spinel, corundum, and monazite (from Spencer, 1948). The grain sizes of various heavy-mineral species from commercial concentrates are given by Garnar (1980). The median size for altered ilmenite is about 0.15 mm and for zircon about 0.10 mm. Staurolite is coarser than either (Garnar, 1980, fig. 8).

HEAVY-MINERAL VARIATION WITH SEDIMENTARY STRUCTURE

Force and Garnar (1985) found that high-angle eolian crossbeds are outlined by laminae up to 1 cm in thickness that are enriched in heavy minerals (and preferentially cemented by iron-organic compounds; see fig. 9). Intervening light-colored sands are coarser (about 0.3 mm) than the dark layers (about 0.2 mm). The light-colored sands in one layer contained 1.0 percent heavy minerals, of which only 36 percent was titanium minerals and 15 percent was zircon. This light-colored layer has high proportions of the lighter heavy minerals staurolite (24 percent), tourmaline (13 percent), and aluminosilicates (13 percent). As can be seen in figure 9, these are present as relatively coarse grains.

An adjacent layer of dark-colored sand contained 6.3 percent heavy minerals. The light mineral quartz, as well as the heavy minerals, is finer in these layers, although some coarser quartz is present (fig. 9). The heavy-mineral assemblage in the dark layer is dominated by titanium minerals (54 percent) and zircon (28 percent); staurolite, tourmaline, and aluminosilicates are each less than 10 percent of the heavy-mineral assemblage.

³ The nature of altered ilmenite at Trail Ridge needs some explanation here. Below the surficial leucoxene zone, altered ilmenite consists of a microcrystalline aggregate of pseudorutile and other less-abundant phases (Temple, 1966) with little ilmenite detectable to X-ray diffraction. The TiO_2 contents of these aggregates average about 65 percent. Magnetic and density properties and color are still enough like ilmenite that the name ilmenite has customarily been retained.




FIGURE 8.—Scanning electron microscope photographs of grains from the Trail Ridge ore sand. *A*, Quartz. *B*, Ilmenite. The structure of this grain suggests a concentric weathering zonation developed prior to deposition.

For this study, the grain-size distributions of mineral fractions in adjacent light-colored and dark-colored laminae were determined for us by Jeanette (Hilbish) Thomas. The sampled laminae are shown in figure 10, and the results in figure 11. The light-colored lamina contained 0.6 percent heavy ($\rho > 2.85$) minerals, whereas the dark-colored one contained 12.3 percent. The modal grain sizes of each are similar to those analyzed by Force and Garnar (1985). In the dark-colored layer the modal grain size of both light-mineral and heavy-mineral fractions are 0.5 to 0.75 ϕ finer than in the adjacent light-colored layer. In each layer, the heavy minerals are 0.5 to 0.75 ϕ finer than the light minerals. The dark lamina contains a heavy-mineral assemblage having a much higher percentage of the densest heavy minerals, ilmenite and zircon (78 percent); the light-colored layer contains an assemblage dominated by sillimanite, staurolite, and tourmaline (55 percent). In the dark-colored bands, the size distribution of individual heavy-mineral species is a sensitive function of mineral density, with the median size of sillimanite ($\rho = 3.25$) almost 1 ϕ coarser than that of zircon ($\rho = 4.6$). Figure 11 suggests that ilmenite ($\rho = 4.7$ if fresh) behaved during deposition like a mineral having specific gravity of 4.0 to 4.6 and hence must have already lost density by leaching. Leucoxene apparently had its present specific gravity of about 4.0 at deposition.

Our calculations, assuming that these layers are typical and that dark layers constitute 10 percent of the

deposit, indicate that 50 to 75 percent of the titanium minerals of the deposit, and 55 to 90 percent of its zircon, are contained in these thin, dark laminae. The dark layers contain only 10 percent of the tourmaline of the deposit, however, and comparably low percentages of other minerals of intermediate density.

ORIGIN

The Trail Ridge ilmenite ore sand in the area where it is best known is clearly eolian. Steep slip faces dip southwest; dune migration must therefore have been in this direction.

The separation of grains by size into laminae richer and poorer in heavy minerals suggests that the grains are entrainment equivalents (in the sense of Slingerland, 1977). However, this equivalence probably was not established on the slip face, where the laminae formed, but on the windward side of the dune, where sorting power is great. Heavy-mineral-enriched material may have been supplied from the windward side in daily pulses (as in Hunter and Richmond, 1986).

NATURE OF TRAIL RIDGE LIGNITIC PEAT

GENERAL DESCRIPTION

The lower stratigraphic unit of the Trail Ridge sequence is referred to as peaty or sapropelic sediment




FIGURE 9.—Photomicrograph of thin section of Trail Ridge ilmenite ore sand, from impregnated box core taken toward base of the sequence shown in figure 5. Orientation is right-side-up. Field of view is 6 mm; plane light. Note dark iron-organic cement in heavy-mineral lamina.

by E.C. Pirkle and Yoho (1970) and is described as containing over 92 percent woody substances in a mat 1.5 m thick. We find that Trail Ridge peat when dried is medium brown in color and has recognizable woody fragments in an amorphous organic matrix, which was deformed by compaction into a fissile to blocky-fracturing mat. A laminated appearance is imparted by flattened and reoriented woody fragments. The considerable variation in peat structure includes unconsolidated peaty sand and solid plugs of wood.

The peat layer marks the lower limit of the ilmenite ore sand, and the buried stumps and branches in it are a mining nuisance. E.C. Pirkle and others (1977) note in passing that some of these stumps are in growth position. Pine cones were noted by Carpenter and others (1953).

Our samples of this organic layer, which we shall refer to as Trail Ridge peat, come from two cores of sediment. They were acquired through a joint drilling project managed by E.I. Du Pont de Nemours and Company and the Florida Bureau of Geology. The cores are referred to



FIGURE 10.—Laminae alternately rich and poor in heavy minerals, below humate-cemented material (at top of photograph) at locality of figure 5. (See fig. 7 for location.) Sample of heavy-mineral-rich laminae analyzed for figure 11 is shown by arrow. The heavy-mineral-poor laminae of figure 11 immediately overlie it. Field of view is 0.8 m.

as RTR1 and RTR2 and come from the area of the Trail Ridge ore body in Clay County, Fla. (fig. 7).

PETROGRAPHY

Rich and others (1978) showed that Trail Ridge peat is actually intermediate between peat and lignite in character. The light color of palynomorphs, pronounced solubility of the sediment in alkali, and certain ultraviolet-emission spectral characteristics support its designation as a peat. On the other hand, the morphology of petrographic constituents (for example, the fact that samples can be polished), the huminite reflectance, and other ultraviolet-emission spectral properties indicate that the sediment is a lignite.

Microscopic analysis of 22 polished pellets of Trail Ridge peat is summarized in table 1 and shown with




FIGURE 11.—Comparison of adjacent laminae rich and poor in heavy minerals. The upper graphs show histograms of the whole sample, the middle graphs show histograms of the heavy-mineral fraction, and the bottom graph shows cumulative curves for individual heavy-mineral species. Analyses by Jeanette (Hilbish) Thomas.

palynologic information in figure 12. The dominant constituent in all samples is detrital particles of various degraded humic materials (humodetrinite). Other constituents (defined in table 1) are textinite, ulminite, and gelinite. Local horizons of charcoal (fusinite) and fungal remains are present. Dominance by humodetrinite (fig. 13) suggests an analogy with highly degraded shrub peats of the modern Okefenokee Swamp (Rich, 1979; Cohen and others, 1984).

PALYNOLOGY

Pollen and spores in the Trail Ridge peat were studied by Rich (1985). Stratigraphic trends in microflora for

core RTR2 are shown in figure 12 for comparison with petrographic constituents. In most samples, the dominant microflora is of the shrubs and small trees, especially holly (*Ilex*), wax myrtle/bayberry (*Myrica*), ti-ti (*Cyrilla*), hazel (*Corylus*) and loblolly bay (*Gordonia*), and herbaceous forms such as sedges (*Cyperaceae*), ferns (*Osmunda*), and moss (*Sphagnum*). All of these forms are consistent with or diagnostic of a freshwater origin, and some suggest standing water.

A stratigraphically lower zone of peat in RTR1 contains more abundant cypress (*Taxodium*) pollen and pollen of shrubby forms consistent with a bald cypress swamp community. Ti-ti (*Cyrilla*) shows a peak in the

TABLE 1.—Organic and sand constituents of core samples from Trail Ridge peat
[n.d. = no data; locations of cores are shown in figure 7]

Ulminite. Biochemically and (or) geochemically gelified tissues having little cell structure remaining.
Humodetrinite. Detrital plant fragments and bits of gelified plant tissue.
Gelinites. A broad range of structureless organic materials derived exclusively by precipitation of colloidal humic suspensions within cell cavities and interparticulate spaces.
Other huminites. A variety of morphologically dissimilar organic substances, generally derived from colloidal humic suspensions (for example, corpohuminite). Includes textinite (coalified cell walls).
Fusinite. Charred tissues, usually having well-preserved cell structure.
Other inertinites. Highly oxidized byproducts of biochemical and geochemical alteration, as well as fungal remains and minute fragments of charcoal.
Liptinites. Resins, waxes, spores and pollen, algae, and plant cuticles.
Textinite. See Other huminites.

| Sampled interval (m depth) | Macerals, expressed as volume percent of organic matter ¹ | | | | | | | Sand, expressed as approximate volume percent of peat ² | Ash, expressed as weight percent dry basis |
|-------------------------------|----------------------------------------------------------------------|---------------|-----------|-----------------|----------|-------------------|------------|-----------------------------------------------------------------------|-----------------------------------------------|
| | Ulminite | Humodetrinite | Gelinites | Other huminites | Fusinite | Other inertinites | Liptinites | | |
| Core RTR1 | | | | | | | | | |
| 15.6 | 30.4 | 54.3 | 1.6 | 4.5 | 2.8 | 4.0 | 2.4 | n.d. | 30.7 |
| 15.7 | 63.3 | 16.3 | 2.1 | 2.5 | 4.8 | 6.8 | 4.2 | n.d. | 4.3 |
| 15.8 | 22.0 | 64.2 | .8 | 4.0 | 1.2 | 3.3 | 4.5 | n.d. | 4.7 |
| 16.0 | 13.0 | 74.5 | 1.3 | 2.0 | 1.3 | 5.8 | 2.1 | n.d. | 4.9 |
| 16.2 | 19.7 | 58.8 | 2.0 | 2.6 | 8.2 | 6.7 | 2.0 | 4 percent | 36.3 |
| 16.3 | 15.5 | 71.1 | 2.5 | 5.1 | 1.8 | 2.8 | 1.2 | .1 percent | 14.0 |
| 16.5 | 8.7 | 83.7 | 1.0 | 2.8 | .0 | 2.2 | 1.6 | .1 percent | 4.5 |
| 16.6 | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | <.1 percent | 39.0 |
| 16.7–17.4 | 13.8 | 77.8 | .7 | 4.5 | .4 | 1.5 | 1.3 | 2 percent | 65–89 |
| 17.4–17.5 | 44.9 | 44.0 | .5 | 5.2 | .5 | 2.5 | 2.4 | n.d. | 33.0 |
| Core RTR2 | | | | | | | | | |
| 16.7–16.9 | 38.6 | 52.1 | 1.9 | 3.4 | 0.1 | 2.0 | 1.9 | 4 percent | 47.8 |
| 16.9 | 19.8 | 69.8 | 1.4 | 2.5 | .9 | 2.6 | 3.0 | n.d. | 57.2 |
| 17.1 | 27.1 | 62.6 | .9 | 3.8 | .3 | 3.0 | 2.3 | 7 percent | 78.2 |
| 17.3 | 14.5 | 66.0 | 3.0 | 4.2 | 4.6 | 5.6 | 2.1 | n.d. | 32.1 |
| 17.4–17.6 | 11.0 | 75.1 | 2.6 | 6.5 | 1.4 | 2.9 | .5 | n.d. | 7.1 |
| 17.6–17.8 | 17.3 | 69.9 | 1.7 | 6.3 | 1.3 | 2.1 | 1.4 | <.1 | 4.8 |
| 17.8–18.0 | 4.6 | 79.8 | 2.7 | 2.2 | 3.6 | 5.9 | 1.2 | n.d. | 8.8 |
| 18.0–18.3 | 10.0 | 77.6 | 4.2 | 1.9 | 3.0 | 2.6 | .7 | n.d. | 6.1 |
| 18.4 | 12.4 | 76.0 | 3.0 | 1.3 | 3.2 | 3.2 | .9 | <.1 | 8.4 |
| 18.5 | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | n.d. | <.1 | 8.6 |
| 18.7 | 16.6 | 74.6 | 1.8 | 2.4 | 2.0 | 1.2 | 1.4 | <.1 | 4.9 |
| 18.8 | 27.4 | 65.9 | 1.0 | 2.7 | .1 | 2.0 | .9 | <.1 | 5.0 |

¹ Maceral abundance was determined by 1,000-point counts of polished pellets in reflected white light.

² Sand abundance was determined in thin section by use of visual comparison charts.

uppermost part of both cores. Oak (*Quercus*) and pine (*Pinus*), although common constituents, are less abundant than in modern swamps of the same region.

Palynology of many samples is closely related to petrography (fig. 12). The pollen was apparently produced by a local flora, as some of these pollen are known to have restricted dispersal patterns (Rich and Spackman, 1977). Analogous correlations have been found elsewhere by Teichmuller (1958), Corvinus and Cohen (1979), and others.

AGE

Palynology by Rich (1985) established only a post-Miocene age for the peat. Some relative-abundance evidence suggested a Pliocene rather than a Pleistocene age, but control by environmental change could not be ruled out for this evidence.

Two samples of lignitic peat from core RTR2 were analyzed for ¹⁴C. Both samples, taken from the upper peat between 16.76 to 17.09 m of depth, proved to be "dead," indicating an age greater than 45,000 years B.P.

(Meyer Rubin, USGS, written commun., 1985). Thus the peat is not of latest Pleistocene or Holocene age. The stratigraphic relations described below suggest that the overlying ilmenite ore sands, likewise, are of post-Miocene, pre-latest Pleistocene age.

THE SAND COMPONENT OF TRAIL RIDGE PEAT

Toward the top of the peat body, several narrow intervals of Trail Ridge PTR cores (fig. 7) have been aptly described as sandy peat (E.C. Pirkle and others, 1970, p. 50). We studied this sandy peat in thin section, as a separated bulk sample, and via analysis for ash content and chemistry. The 12 thin sections, representing most of the peat thickness in the two RTR cores (fig. 7), show that the uppermost peat layers characteristically contain about 1 percent or more of sand (table 1) that is dominantly well rounded, especially the coarser grains (fig. 13). The entire heavy-mineral suite of the overlying Trail Ridge ore body is present. Underlying peats most commonly contain little sand or more angular sand poor in heavy minerals. Ash values (table 1) partly




FIGURE 12.—Palynological (top) and maceral (bottom) constituents of peat in core RTR2. Palynofloral constituents are expressed as a percentage of total microflora (from Rich, 1985). Maceral constituents are expressed as a percentage of total macerals.

reflect the distribution of sand; ash from this peat commonly contains 1 to 2 percent TiO_2 and 0.1 to 0.4 percent zirconium (Jesse Yeakel, written commun., 1980).

Figures 13 and 14 show that rounded sand grains in the upper peat are isolated and disseminated and embedded in laminated organic matrix, with deformed organic material draped around the grains. The organic matrix of




FIGURE 13.—Thin-section photomicrograph of exceptionally sandy lignitic peat from core RTR2, 17.09-m depth. Field of view is 6 mm; plane light. Sand is imbedded in an organic matrix of humodetrinite.

this upper sandy peat and peaty sand is dominated by humodetrinite (table 1) with admixed charcoal and (or) fungal remains, suggesting subaerial exposure of the peat during sand accumulation. It appears unlikely that the sand represents either traction-load detritus or material reworked from the overlying sand mass after deposition; in the former case the sand would be bedded, and in the latter case the sand would occupy veins. In either case, the sand grains would be in contact.

We were able to separate 1.1 g of sand from the RTR2 sample and subjected it to one of the most rigorous textural and mineralogic analyses possible for a sample of this size. Figure 15A shows the grain-size distribution of the sand. The modal grain size is about 0.2 mm, sorting is rather poor, and a fine tail is present in the grain-size population. Skewness is probably slightly positive. Compared to the overlying Trail Ridge ilmenite ore sand (fig. 11), there is a great deal of overlap in grain-size distri-



FIGURE 14.—Scanning electron microscope photograph of sand-bearing lignitic peat sample from core RTR2, 16.76 to 16.99 m. Note the isolation of sand grains in an organic matrix of plant fragments.

bution, but modal grain size of sand in peat is finer and sorting is poorer.

Heavy-mineral content of this sand is 6.7 percent, slightly higher than in the overlying ilmenite ore sand. Heavy minerals show a modal grain size slightly finer than that of light minerals (fig. 15B).

The mineralogy of the sand in peat compares closely with that of overlying ilmenite ore sand. The light fraction of the sand is over 99 percent quartz. No feldspar was identified. The heavy minerals are dominated by ilmenite (46.5 percent by volume), zircon (16 percent), staurolite (8.5 percent), and sillimanite+kyanite (7.5 percent) (fig. 15). Minor but characteristic constituents are tan or gray leucoxene (5 percent), rutile (4 percent), tourmaline (2 percent), and monazite, spinel, corundum, and garnet (all less than 1 percent). Size distributions of the heavy minerals (fig. 15C) are apparently a function of mineral density and are related to each other much as in the overlying ilmenite ore sand.

Grain shapes and surface characteristics of the sand grains are also similar to those of the ilmenite ore sand (fig. 16). Rounding and sphericity are moderate to superb. The sand grains are frosted and show microscopic impact and abrasion features suggestive of an eolian origin. (See Krinsley and Margolis, 1971.)






FIGURE 15.—Grain-size distributions of inorganic fraction of sandy Trail Ridge peat from core RTR2, 16.76- to 16.99-m depth. A, Grain-size histogram of inorganic fraction, with heavy minerals ($\rho > 2.85$) shaded. B, Grain-size histogram of heavy minerals. C, Cumulative distributions of staurolite, ilmenite, zircon, and sillimanite.

FIGURE 16.—Scanning electron microscope photographs of sand grains in Trail Ridge peat from RTR2, 16.76- to 16.99-m depth. A, Quartz. B, Ilmenite. C, Zircon.



FIGURE 17.—Holocene coastal transgressive longwall dune near Newcastle, New South Wales, Australia, taken from the landward side. This dune, an economic heavy-mineral deposit, has migrated away from the coast but remains parallel to it and now rests in part on Holocene peats (Thom and others, 1981).

Certainly the sand did not acquire these features in the depositional environment, a swamp. The distribution of sand suggests that the transport agent that moved it there apparently did so in some form of suspension, not as bed load. In a swamp, the appropriate form of suspension transport is eolian. An eolian origin for the sand component of peat is an additional tie to the overlying ilmenite ore sand, which is independently known to be eolian.

DEPOSITIONAL RELATION OF TRAIL RIDGE PEAT AND SANDS

The Trail Ridge ilmenite ore sand represents a large southwest-migrating eolian dune complex. Trail Ridge peat represents a freshwater swamp, originally formed west of the dune but later overridden by it. The sand component of the upper peat was deposited from aerial suspension and acquired its characteristics in the adjacent eolian environment. The sand serves as a sensitive recorder of the approach of the dune; it represents the finer fraction that remained in suspension during eolian flow separation at the top of the slip face of the dune. The

ilmenite ore sand represents the fraction that moved down the slip face as traction load. Thus the peat and the ilmenite ore sand are essentially the same age.

The existence of a large active dune, probably little-vegetated, upwind of the swamp explains the scarcity of pine and oak pollen in the peat. The portions of the swamp nearest the dune were largely subaerially exposed and vegetated partly by *ti-ti*. Some other portions of the swamp were subaqueous, as indicated by some of the palynoflora of the peat.

Oxidation is the first stage of ilmenite alteration (Temple, 1966) and is important in the weathering alteration of other minerals also. Thus the sand entombed in reducing peat should serve also as a recorder of the original composition of ilmenite and other minerals supplied to the dune before postdepositional weathering. The presence of leucoxene and the absence or trace amounts of labile constituents such as feldspar show that the dune was supplied by a sand already highly weathered. This observation is consistent with the previously discussed surface polish and grain-size-density characteristics in the ore sand. Since postdepositional weathering at Trail Ridge is already well established, a two-stage weathering history is implied.




FIGURE 18.—Schematic paleogeography of Trail Ridge at the time of dune formation.

Eolian dunes very commonly block natural drainage channels, and it seems likely that the active Trail Ridge dune thus impounded the swamp represented by Trail Ridge peat. In the same way, the inactive Trail Ridge landform now impounds the modern Okefenokee Swamp.

No evidence of marine influence in deposition of the Trail Ridge sequence has ever been encountered. All observed sands are eolian, and all observed sheltered deposits were deposited in fresh water. Marine fossils collected by F.L. Pirkle and Czel (1983) were thought by them to postdate Trail Ridge deposition.

PALEOGEOGRAPHIC RECONSTRUCTION

The lack of marine facies coeval with Trail Ridge deposition suggests that the Trail Ridge dune had become decoupled from parent shoreline facies and moved inland while remaining roughly parallel to shore, similar to the formation of transgressive longwall dunes of eastern Australia (Thom and others, 1981). These

bodies, of both Pleistocene and Holocene age, are similarly decoupled from parallel coeval shorelines and rest in part on marsh and swamp deposits (fig. 17). An eolian body as large as Trail Ridge would probably have been composite, composed of individual transgressive parabolic dunes oriented according to prevailing wind direction (fig. 18), analogous to Pleistocene dunes of the southern Queensland coast (Ward, 1977, 1978; Thompson and Ward, 1975) that are even larger than Trail Ridge. Many of the Australian transgressive dunes are also heavy-mineral ore bodies (McKellar, 1975).

Transgressive dunes may move inland considerable distances and climb onto irregular surfaces well above coeval sea levels (Australia—Hesp, 1986; Short, 1987; Brazil—Bigarella, 1972). Our conception of paleogeography at the time of formation of the Trail Ridge dune is shown in figure 18. The transgressive dune moved west, roughly parallel to the coast, but migrated by successive accretions of parabolic dunes moving upslope toward the southwest, thus forming slip facies oriented as shown in figure 6.

ACKNOWLEDGMENTS

The authors would like to thank E.I. Du Pont de Nemours Company and the Florida Bureau of Geology for help in the field and with core acquisition and T.E. Garnar and Fred Pirkle for useful advice. Jesse Yeakel of Exxon Corporation and William Spackman of Pennsylvania State University provided thin sections and petrographic data. John Evans (USGS, Reston) took the scanning electron microscope photographs. Jeanette (Hilbish) Thomas generated the data for figure 11. Meyer Rubin (USGS) provided the radiometric dates on peat. The Department of Geology and Geological Engineering at South Dakota School of Mines and Technology is acknowledged for extending support to F.J. Rich.

REFERENCES

- Bigarella, J.J., 1972, Eolian environments: their characteristics, recognition, and importance: Society Economic Paleontologists and Mineralogists Special Publication 16, p. 12–62.
- Cannon, H.B., 1950, Economic minerals in the beach sands of the Southeastern United States, in Snyder, F.G., ed., Symposium on mineral resources of the Southeastern United States: University Tennessee Press, p. 202–210.
- Carpenter, J.H., Detweiler, J.C., Gillson, J.T., Weichel, E.C., and Wood, J.P., 1953, Mining and concentration of ilmenite and associated minerals at Trail Ridge, Fla.: Mining Engineering, August, p. 1–7.
- Cohen, A.D., Andejko, M.J., Spackman, W., and Corvinus, D.A., 1984, Peat deposits of the Okefenokee Swamp, in Cohen, A.D., ed., The Okefenokee Swamp: Wetlands Surveys, Los Alamos.
- Cooke, C.W., 1925, The coastal plain, in Physical geography of Georgia: Georgia Geological Survey Bulletin 42, p. 19–57.

- 1939, Scenery of Florida interpreted by a geologist: Florida Department Conservation Geological Bulletin 17, 118 p.
- Corvinus, D.A., and Cohen, A.D., 1979, Petrographic characteristics of carbonaceous sediments from the Snuggedy Swamp of South Carolina [abs.]: 9th International Congress Carboniferous Stratigraphy and Geology, Urbana, Abstracts, p. 45.
- Creitz, E.E., and McVay, T.N., 1948, A study of opaque minerals in Trail Ridge, Florida, dune sands: American Institute Mining Metallurgical Engineers Technical Publication 2426, 7 p.
- Force, E.R., and Garnar, T.E., 1985, High-angle aeolian crossbedding at Trail Ridge, Florida: *Industrial Minerals*, August, p. 55-59.
- Friedman, G.M., 1961, Distinction between dune, beach, and river sands from their textural characteristics: *Journal Sedimentary Petrology*, v. 31, p. 514.
- Garnar, T.E., Jr., 1972, Economic geology of Florida heavy mineral deposits: Florida Bureau of Geology Special Publication 17, p. 17-21.
- 1980, Heavy minerals industry of North America, *in*, Coope, B.M., ed., Proceedings of the 4th Industrial Minerals International Congress Industrial Minerals, London, p. 29-42.
- Hesp, P.A., 1986, A review of Australian coastal vegetated dune evolution, dynamics, geomorphology, and internal structures [abs.]: 12th International Sedimentological Congress (Canberra) Abstracts, p. 139-140.
- Hunter, R.E., and Richmond, B.M., 1986, Daily cycles in coastal dunes [abs.]: 12th International Sedimentological Congress (Canberra) Abstracts, p. 147-148.
- Krinsley, D.H., and Margolis, S.V., 1971, Surface grain texture, *in* Carver, R.E., ed., Procedures in sedimentary petrology: New York, Wiley-Interscience, p. 151-180.
- MacNeil, F.S., 1950, Pleistocene shore lines in Florida and Georgia: U.S. Geological Survey Professional Paper 221-F, p. 95-108.
- McKellar, J.B., 1975, The eastern Australia rutile province: Australasian Institute Mining Metallurgy Monograph 5, p. 1055-1061.
- Pirkle, E.C., Pirkle, W.A., and Yoho, H.H., 1974, The Green Cove Springs and Boulougne heavy-mineral sand deposits of Florida: *Economic Geology*, v. 67, p. 1129-1137.
- 1977, The Highland heavy-mineral sand deposit on Trail Ridge in northern peninsular Florida: Florida Bureau of Geology Report Investigation 84, 50 p.
- Pirkle, E.C., and Yoho, W.H., 1970, The heavy mineral ore body of Trail Ridge, Florida: *Economic Geology*, v. 65, p. 17-30.
- Pirkle, E.C., Yoho, W.H., and Hendry, C.W., 1970, Ancient sea level stands in Florida: Florida Bureau of Geology, Geological Bulletin 52, 61 p.
- 1971, North Florida setting for fifteenth field conference, Southeastern Geological Society: Geological Review of Some North Florida Mineral Resources, Southeastern Geological Society, Tallahassee, p. 1-15.
- Pirkle, F.L., 1975, Evaluation of possible source regions of Trail Ridge sands: *Southeastern Geology*, v. 17, p. 93-114.
- 1984, Environment of deposition of Trail Ridge sediments as determined from factor analysis, *in* Cohen, A.D., ed., The Okefenokee Swamp: Los Alamos, Wetlands Surveys, p. 629-650.
- Pirkle, F.L., and Czel, L.J., 1983, Marine fossils from region of Trail Ridge, a Georgia-Florida landform: *Southeastern Geology*, v. 24, p. 31-38.
- Pirkle, W.A., and Pirkle, E.C., 1984, Physiographic features and field relations of Trail Ridge in northern Florida and southeastern Georgia, *in* Cohen, A.D., ed., The Okefenokee Swamp: Los Alamos, Wetlands Surveys, p. 613-628.
- Rich, F.J., 1979, The origin and development of tree islands in the Okefenokee Swamp, as determined by peat petrography and pollen stratigraphy: unpublished Ph. D. dissertation, the Pennsylvania State University, 301 p.
- 1985, Palynology and paleoecology of a lignitic peat from Trail Ridge, Florida: Florida Bureau of Geology Information Circular 100, 15 p.
- Rich, F.J., and Spackman, W., 1977, A comparison of modern and ancient pollen: sedimentation around tree islands in the Okefenokee Swamp [abs.]: American Association Stratigraphic Palynology, 10th Annual Meeting, p. 25.
- Rich, F.J., Yeakel, J.D., and Gutzler, R.Q., 1978, Trail Ridge buried peat: its significance as a coalification intermediate [abs.]: Geological Society America Abstracts with Programs, v. 10, no. 7, p. 478.
- Short, A.D., 1987, Modes, timing and volume of Holocene cross-shore and aeolian sediment transport, southern Australia, *in* Kraus, N.C., ed., Coastal sediments '87: New York, N.Y., American Society of Civil Engineers, v. 2, p. 1925-1937.
- Slingerland, R.L., 1977, The effects of entrainment on the hydraulic equivalence relationships of light and heavy minerals in sands: *Journal Sedimentary Petrology*, v. 47, p. 753-770.
- Spencer, R.V., 1948, Titanium minerals in Trail Ridge, Fla.: U.S. Bureau of Mines Report of Investigations 4208, 21 p.
- Teichmuller, M., 1958, Rekonstruktion verschiedener Moortypen des Hauptflozes der niederheinischer Braunkohle: *Fortschrift Geologie Rheinland und Westfalen*, v. 2, p. 599-612.
- Temple, A.K., 1966, Alteration of ilmenite: *Economic Geology*, v. 61, p. 695-714.
- Thoenen, J.R., and Warne, J.D., 1949, Titanium minerals in central and northeastern Florida: U.S. Bureau of Mines Report of Investigations 4515, 62 p.
- Thom, B.G., Bowman, G.M., and Roy, P.S., 1981, Late Quaternary evolution of coastal sand barriers, Port Stephens-Myall Lakes area, central New South Wales: *Quaternary Research*, v. 15, p. 345-364.
- Thompson, C.H., and Ward, W.T., 1975, Soil landscapes of North Stradbroke Island: Proceedings Royal Society Queensland, v. 86, p. 9-14.
- Trask, P.D., 1932, Origin and environment of source sediments of petroleum: Houston, Tex., Gulf Publishing Co., 323 p.
- Ward, W.T., 1977, Sand movement on Fraser Island: a response to changing climates: University Queensland, Occasional Papers in Anthropology no. 8, p. 113-126.
- 1978, Notes on the origin of Stradbroke Island: University Queensland Geologic Department Papers, v. 8, p. 97-104.

SELECTED SERIES OF U.S. GEOLOGICAL SURVEY PUBLICATIONS

Periodicals

- Earthquakes & Volcanoes (issued bimonthly).
- Preliminary Determination of Epicenters (issued monthly).

Technical Books and Reports

Professional Papers are mainly comprehensive scientific reports of wide and lasting interest and importance to professional scientists and engineers. Included are reports on the results of resource studies and of topographic, hydrologic, and geologic investigations. They also include collections of related papers addressing different aspects of a single scientific topic.

Bulletins contain significant data and interpretations that are of lasting scientific interest but are generally more limited in scope or geographic coverage than Professional Papers. They include the results of resource studies and of geologic and topographic investigations; as well as collections of short papers related to a specific topic.

Water-Supply Papers are comprehensive reports that present significant interpretive results of hydrologic investigations of wide interest to professional geologists, hydrologists, and engineers. The series covers investigations in all phases of hydrology, including hydrogeology, availability of water, quality of water, and use of water.

Circulars present administrative information or important scientific information of wide popular interest in a format designed for distribution at no cost to the public. Information is usually of short-term interest.

Water-Resources Investigations Reports are papers of an interpretive nature made available to the public outside the formal USGS publications series. Copies are reproduced on request unlike formal USGS publications, and they are also available for public inspection at depositories indicated in USGS catalogs.

Open-File Reports include unpublished manuscript reports, maps, and other material that are made available for public consultation at depositories. They are a nonpermanent form of publication that may be cited in other publications as sources of information.

Maps

Geologic Quadrangle Maps are multicolor geologic maps on topographic bases in 7 1/2- or 15-minute quadrangle formats (scales mainly 1:24,000 or 1:62,500) showing bedrock, surficial, or engineering geology. Maps generally include brief texts; some maps include structure and columnar sections only.

Geophysical Investigations Maps are on topographic or planimetric bases at various scales; they show results of surveys using geophysical techniques, such as gravity, magnetic, seismic, or radioactivity, which reflect subsurface structures that are of economic or geologic significance. Many maps include correlations with the geology.

Miscellaneous Investigations Series Maps are on planimetric or topographic bases of regular and irregular areas at various scales; they present a wide variety of format and subject matter. The series also includes 7 1/2-minute quadrangle photogeologic maps on planimetric bases which show geology as interpreted from aerial photographs. Series also includes maps of Mars and the Moon.

Coal Investigations Maps are geologic maps on topographic or planimetric bases at various scales showing bedrock or surficial geology, stratigraphy, and structural relations in certain coal-resource areas.

Oil and Gas Investigations Charts show stratigraphic information for certain oil and gas fields and other areas having petroleum potential.

Miscellaneous Field Studies Maps are multicolor or black-and-white maps on topographic or planimetric bases on quadrangle or irregular areas at various scales. Pre-1971 maps show bedrock geology in relation to specific mining or mineral-deposit problems; post-1971 maps are primarily black-and-white maps on various subjects such as environmental studies or wilderness mineral investigations.

Hydrologic Investigations Atlases are multicolored or black-and-white maps on topographic or planimetric bases presenting a wide range of geohydrologic data of both regular and irregular areas; principal scale is 1:24,000 and regional studies are at 1:250,000 scale or smaller.

Catalogs

Permanent catalogs, as well as some others, giving comprehensive listings of U.S. Geological Survey publications are available under the conditions indicated below from the U.S. Geological Survey, Books and Open-File Reports Section, Federal Center, Box 25425, Denver, CO 80225. (See latest Price and Availability List.)

"**Publications of the Geological Survey, 1879-1961**" may be purchased by mail and over the counter in paperback book form and as a set of microfiche.

"**Publications of the Geological Survey, 1962-1970**" may be purchased by mail and over the counter in paperback book form and as a set of microfiche.

"**Publications of the U.S. Geological Survey, 1971-1981**" may be purchased by mail and over the counter in paperback book form (two volumes, publications listing and index) and as a set of microfiche.

Supplements for 1982, 1983, 1984, 1985, 1986, and for subsequent years since the last permanent catalog may be purchased by mail and over the counter in paperback book form.

State catalogs, "List of U.S. Geological Survey Geologic and Water-Supply Reports and Maps For (State)," may be purchased by mail and over the counter in paperback booklet form only.

"**Price and Availability List of U.S. Geological Survey Publications**," issued annually, is available free of charge in paperback booklet form only.

Selected copies of a monthly catalog "New Publications of the U.S. Geological Survey" available free of charge by mail or may be obtained over the counter in paperback booklet form only. Those wishing a free subscription to the monthly catalog "New Publications of the U.S. Geological Survey" should write to the U.S. Geological Survey, 582 National Center, Reston, VA 22092.

Note.--Prices of Government publications listed in older catalogs, announcements, and publications may be incorrect. Therefore, the prices charged may differ from the prices in catalogs, announcements, and publications.