



metabolism and so on.

## **Exercise Biochemistry Review**

Proceedings of IBEC 2018, Beijing, China, October 23-25 P0-194

## Urinary metabolomics study on effects of Rhodiola on Marathon Amateurs after Quantitative Exercise Load

Yumei Han¹,Yi Jia¹,Junsheng Tian²,Bo Zhang¹,Xiong Xue¹
1.School of PE, Shanxi University
2.Modern Research Center for Traditional Chinese Medicine, Shanxi University,

**Objective** To study the effect of Chinese medicine Rhodiola on oxidative stress injury in amateur marathon runners after quantitative exercise load (20 km) and explore its mechanism. **Methods** Eight marathon amateurs were divided into four groups according to different test time, including before and after quantitative exercise load (group C and group CE), before and after quantitative exercise load after taking a month of the Rhodiola (group MC and group ME). The participants had serum superoxide dismutase (SOD), glutathione peroxidase (GSH-Px), total antioxidant capacity (T-AOC) and malondialdehyde (MDA), as well as myocardial enzyme index - creatine kinase (CK), lactate dehydrogenase (LDH), creatine kinase isoenzyme (CK-MB) and glutamic oxaline aminotransferase (AST/GOT) activity evaluated. In order to further explore the mechanism of action of Rhodiola, the urine was analyzed by ¹H-HMR metabonomics technique. **Results** 1) Compared with group C, the activity of serum CK, LDH and AST/GOT of group CE increased significantly (P<0.01 or P<0.05), MDA content and SOD activity also increased significantly (P<0.05). A total of fifteen potential biomarkers were found in group CE, such as valine, lactic acid, 2-hydroxy isobutyric acid and so on (P<0.05 and VIP>1). 2) Compared with group CE, the activity of serum CK and AST/GOT and the content of MDA of group ME decreased significantly (P<0.05), eleven

**Conclusions** Rhodiola can enhance the antioxidant capacity and improve myocardial damage of marathon amateurs after quantitative exercise load, which may be due to increased synthesis and utilization of aminoacyl-tRNA and other amino acids.

metabolites among the fifteen potential biomarkers reverted significantly (P<0.01 or P<0.05), which

mainly involved in 4 metabolic pathways including alanine, aspartic acid and glutamic acid