



Missouri University of Science and Technology
Scholars' Mine

Physics Faculty Research & Creative Works

Physics

01 Jun 2012

Triple Compton Effect: A Photon Splitting into Three upon Collision with a Free Electron

Erik Loetstedt

Ulrich D. Jentschura

Missouri University of Science and Technology, ulj@mst.edu

Follow this and additional works at: https://scholarsmine.mst.edu/phys_facwork

 Part of the [Physics Commons](#)

Recommended Citation

E. Loetstedt and U. D. Jentschura, "Triple Compton Effect: A Photon Splitting into Three upon Collision with a Free Electron," *Physical Review Letters*, vol. 108, no. 23, pp. 233201-1-233201-5, American Physical Society (APS), Jun 2012.

The definitive version is available at <https://doi.org/10.1103/PhysRevLett.108.233201>

This Article - Journal is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Physics Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.

Triple Compton Effect: A Photon Splitting into Three upon Collision with a Free Electron

Erik Lötstedt* and Ulrich D. Jentschura

Department of Physics, Missouri University of Science and Technology, Rolla, Missouri 65409-0640, USA
(Received 22 February 2012; published 6 June 2012; corrected 24 July 2012)

The process in which a photon splits into three after the collision with a free electron (triple Compton effect) is the most basic process for the generation of a high-energy multiparticle entangled state composed out of elementary quanta. The cross section of the process is evaluated in two experimentally realizable situations, one employing gamma photons and stationary electrons, and the other using keV photons and GeV electrons of an x-ray free electron laser. For the first case, our calculation is in agreement with the only available measurement of the differential cross section for the process under study. Our estimates indicate that the process should be readily measurable also in the second case. We quantify the polarization entanglement in the final state by a recently proposed multiparticle entanglement measure.

DOI: [10.1103/PhysRevLett.108.233201](https://doi.org/10.1103/PhysRevLett.108.233201)

PACS numbers: 34.50.-s, 03.65.Ud, 12.20.Ds

Introduction.—The triple Compton effect is a fundamental process of light-matter interaction, in which a photon splits into three upon collision with an electron,

$$e^- + \gamma \rightarrow e^- + \gamma + \gamma + \gamma. \quad (1)$$

The cross section of this QED process is of fourth order in the fine-structure constant $\alpha \approx 1/137.036$. Currently in the literature there is no general theoretical treatment of the triple Compton effect. In principle, there is no limit of the number of photons that can be coherently emitted when an electron interacts with a photon, but the only processes that have been discussed so far are the usual, second-order (in α) single Compton effect [1] and the third-order double Compton effect [2]. Only the single Compton effect has a classical limit. To the contrary, both the double and the triple Compton effects are intrinsically quantum processes that cannot be described by classical electrodynamics. With strong lasers becoming available, the nonlinear generalization of Compton scattering in which several laser photons are absorbed have been vigorously discussed (see [3,4] for recent investigations of the nonlinear Compton effect, and [5–7] for a discussion on the nonlinear double Compton effect, where two photons are coherently emitted).

The three photons emitted in the process (1) originate from the same initial photon, are emitted at the same time, and are therefore quantum mechanically entangled. The creation of an entangled state of three photons is an important goal in quantum information. The conventional way of generating entangled triple-photon states is by employing nonlinear crystals [8–10], but one can imagine other sources, not yet experimentally realized, such as electron-positron annihilation [11,12]. In this Letter, we propose the triple Compton effect as an alternative source of entangled photon triplets, which by suitable optimization could be competitive both in terms of production rate and the degree of entanglement. Especially interesting is the possibility of creating correlated photons with high

energy in the GeV range, via triple Compton backscattering on a relativistic electron beam. We will show that an advantageous setup for such an experiment is an x-ray free electron laser (XFEL), providing a high-flux, high-energy photon beam together with a GeV electron beam.

The only previous experimental study of the triple Compton effect that we are aware of is Ref. [13], where the differential cross section was measured for a well-defined interval of the solid angle, as defined by the detectors that were arranged in a symmetric angular configuration. On the theoretical side, the only preceding investigation is [14], where the total cross section in the limit of ultrarelativistic initial photon energy was studied. In our treatment of the problem, we can evaluate the differential cross section for arbitrary values of the directions, energies, and polarizations of the emitted photons. On the other hand, the double Compton effect is rather well studied, both experimentally [15–18] and theoretically [2,19,20]. In addition, several other processes have been studied where two photons are produced in the final state, such as double bremsstrahlung [21–23], bound-state decay [24,25], and laser-induced photon splitting [26], but comparatively little is known about triple photon production.

Theoretical formulation.—Unless stated otherwise, natural units with $\hbar = c = 1$ are used, and m denotes the electron mass. Four-vector products are denoted with a dot (i.e., we have $a \cdot b \equiv a^\mu b_\mu = a^0 b^0 - \mathbf{a} \cdot \mathbf{b}$ for two four-vectors a and b). The contraction with the Dirac matrices γ^μ is written with a hat, $\hat{a} = a^0 \gamma^0 - \mathbf{a} \cdot \boldsymbol{\gamma}$.

We label the four-momenta of the incoming electron and photon with $p_i = (E_i, \mathbf{p}_i)$ and $k_0 = \omega_0 n_0 = \omega_0(1, \mathbf{n}_0)$, respectively, and the four-momenta of the outgoing electron and photons with $p_f = (E_f, \mathbf{p}_f)$, and $k_j = \omega_j n_j = \omega_j(1, \mathbf{n}_j)$, $j = 1, 2, 3$, respectively. When explicitly evaluating the cross section, we take the vectors $\mathbf{k}_j = \omega_j \mathbf{n}_j$ in the lab frame, in spherical coordinates with the polar axis directed along \mathbf{k}_0 , i.e., $\mathbf{k}_0 = \omega_0(0, 0, 1)$ and $\mathbf{k}_j = \omega_j(\sin\theta_j \cos\phi_j, \sin\theta_j \sin\phi_j, \cos\theta_j)$. The amplitude M for




FIG. 1. One of the $4! = 24$ contributing Feynman diagrams of the triple Compton effect. To obtain the total amplitude, we have to add the diagrams corresponding to the permutations of the insertion order of the photons k_0 , k_1 , k_2 , and k_3 .

the triple Compton effect is formed from a coherent sum of $4! = 24$ Feynman diagrams, one of which is shown in Fig. 1.

We have

$$M = m^3 \sum_{\xi} u_r^\dagger(p_f) \gamma^0 \hat{\epsilon}_{\xi(3)} \frac{\hat{q}_3(\xi) + m}{q_3^2(\xi) - m^2} \hat{\epsilon}_{\xi(2)} \frac{\hat{q}_2(\xi) + m}{q_2^2(\xi) - m^2} \times \hat{\epsilon}_{\xi(1)} \frac{\hat{q}_1(\xi) + m}{q_1^2(\xi) - m^2} \hat{\epsilon}_{\xi(0)} u_{r_i}(p_i), \quad (2)$$

where the sum runs over all the 24 permutations ξ of $(0, 1, 2, 3)$. The momenta q_n entering the propagators are defined as $q_n(\xi) = p_i + \sum_{j=0}^{n-1} (-1)^{\delta_{0\xi(j)}+1} k_{\xi(j)}$, where δ_{ij} is Kronecker's delta function. (One adds to p_i the four-momentum of the incoming photon k_0 and subtracts the four-momenta of the outgoing photons, according to the relevant permutation ξ). In the expression for the amplitude (2), we have introduced the positive-energy bispinor $u_r(p)$, with the spin index $r = 1, 2$, which we use in the conventions of Chap. 2 of [27] [the normalization is $\bar{u}_r(p)u_r(p) = u_r^\dagger(p)\gamma^0 u_r(p) = 1$]. In Eq. (2), r_i and r_f denote the initial and final electron spins, respectively. The polarization four-vectors $\epsilon_j = (0, \mathbf{\epsilon}_j)$ satisfy $\epsilon_j \cdot k_j = 0$ for $j = 0, 1, 2, 3$. As an explicit set of basis vectors for ϵ_j , we take $\epsilon_j^1 = (\cos\theta_j \cos\phi_j, \cos\theta_j \sin\phi_j, -\sin\theta_j)$ and $\epsilon_j^2 = (-\sin\phi_j, \cos\phi_j, 0)$. The differential cross section follows from the usual rules of QED [28] as

$$\frac{d\sigma}{d\omega_1 d\omega_2 d\Omega_1 d\Omega_2 d\Omega_3} \equiv \sigma_5 = \frac{\alpha^4}{(2\pi)^4} \frac{1}{m^4} \frac{\omega_1 \omega_2 \omega_3}{E_f p_i \cdot k_0} \frac{|M|^2}{|K|} \times \Theta(\omega_3) \Theta(E_f - m), \quad (3)$$

where $\Theta(\cdot)$ is the step function, $d\Omega_j = \sin\theta_j d\theta_j d\phi_j$ is the differential solid angle of photon j , and we have introduced σ_5 as an abbreviation of the fivefold differential cross section. In Eq. (3), the final four-momentum of the electron is fixed by energy-momentum conservation as $p_f = p_i + k_0 - k_1 - k_2 - k_3$, and the energy of photon three is

$$\omega_3 = \frac{p_i \cdot (k_0 - k_1 - k_2) - k_0 \cdot (k_1 + k_2) + k_1 \cdot k_2}{n_3 \cdot (p_i + k_0 - k_1 - k_2)}. \quad (4)$$

The factor K in (3), arising from the final delta function integration over ω_3 reads $K = 1 + \frac{n_3 \cdot (k_1 + k_2 - k_0 - p_i) + \omega_3}{E_f}$.

The differential cross section (3) depends on eight continuous variables, two angles θ_j , ϕ_j for each emitted photon and the energies ω_1 , ω_2 of two of the emitted photons. The energy ω_3 of the third photon is restricted by the kinematical constraints and is calculated by Eq. (4). In addition, we have six more discrete variables that can take any of two values, namely, the polarization vectors ϵ_j of the photons and the spins $r_{i,f}$ of the electrons.

A couple of general remarks about the expression (3) are the following. Similarly to the double Compton effect [28], the cross section vanishes when all three photons are emitted parallel to the incoming photon. In the rest frame of the electron, this implies $\omega_1 + \omega_2 + \omega_3 = \omega_0$ according to Eq. (4), and therefore corresponds to the incoming photon splitting into three photons without any interaction, which is not allowed. Whenever either of ω_1 , ω_2 , or ω_3 goes to zero, σ_5 diverges as $1/\omega_{1,2,3}$, while the radiated energy remains finite in the infrared. This is the well-known infrared catastrophe of QED, which can be cured by adding radiative corrections [29]. In the current case, the divergences at $\omega_n \rightarrow 0$ would cancel against radiative corrections to the single and double Compton effect. The evaluation of such corrections is beyond the scope of this Letter. We will calculate the differential cross section sufficiently far from the infrared divergences, corresponding to a specific experimental detector threshold. Our calculations therefore have a relative accuracy of the order of the fine-structure constant.

Evaluation of the differential cross section.—The evaluation of σ_5 is performed numerically, by employing an explicit representation of γ^μ and $u_r(p)$. This approach is advantageous [30] if one is interested in polarization-resolved cross sections, in which case the analytic evaluation of σ_5 does not simplify. We calculate the differential cross section (3) for two different experimental setups. The first is the same as considered in Ref. [13], where a measurement of the triple Compton cross section is described. In this setup, a gamma photon of energy $\omega_0 = 0.662$ MeV impinges on an electron at rest, $E_i = m$, and photons are detected in coincidence at $\theta_1 = \theta_2 = \theta_3 = \pi/2$, and $\phi_1 = 2\pi/3$, $\phi_2 = 4\pi/3$, $\phi_3 = 0$. The photon energy threshold was $\omega_n \geq \varepsilon = 13$ keV. What was actually measured was the differential cross section averaged over the solid angles subtended by the detectors,

$$\langle \sigma \rangle = \frac{1}{\Omega^3} \frac{1}{4} \sum_{\text{spin, pol}} \int_{\Omega} d\Omega'_1 d\Omega'_2 d\Omega'_3 \int d\omega_1 d\omega_2 \sigma_5, \quad (5)$$

where the solid angle is $\Omega = 0.378$ sr, and the energy integration is over $\omega_{1,2,3} > \varepsilon$. Performing the integration in (5) over the angular intervals $\Delta\phi_j = \phi_j \pm \sqrt{\Omega}/2$, $\Delta\theta_j = \theta_j \pm \arcsin(\sqrt{\Omega}/2)$ so that $\int_{\Delta\theta_j} d\cos\theta'_j \int_{\Delta\phi_j} d\phi'_j = \Omega$, and over photon energies greater than ε , we obtain

$$\langle\sigma\rangle = 4.1 \times 10^{-9} \text{ b/sr}^3. \quad (6)$$

This value should be compared to the measured value of

$$\langle\sigma\rangle^{\text{exp}} = (8.1 \pm 2.4) \times 10^{-9} \text{ b/sr}^3, \quad (7)$$

which includes an experimental uncertainty of 30% [13]. Although the calculated value lies outside the estimated experimental error bar (by 1.6 standard deviations), we believe that our calculation can be regarded as a confirmation of the measurement in Ref. [13], given the utmost difficult nature of the experiment. We have extended the analysis relevant to the experiment [13] to the energy interval $\omega_0 \in (10^{-2}, 10^2)$ MeV; details will be presented elsewhere.

The second example we consider is a triple Compton backscattering scheme: the head-on collision of a relativistic electron, at an incident energy of $E_i = 5$ GeV and $\mathbf{p}_i = (0, 0, -\sqrt{E_i^2 - m^2})$, with an incoming photon of energy $\omega_0 = 1$ keV. Such parameters are realized in an XFEL, for example at the LCLS in Stanford [31], and would require the reflection of the x-ray beam to collide with the electron beam from the accelerator. In this situation, the photons are backscattered and emitted in a narrow cone around the propagation direction of the electron, $\theta_j \sim \pi - E_i/m$. We estimate the total cross section as $\sigma_{\text{TC}}^{\text{tot}} = 2 \times 10^{-5}$ b, compared to $\sigma_{\text{DC}}^{\text{tot}} = 1 \times 10^{-3}$ b and $\sigma_{\text{SC}}^{\text{tot}} = 3 \times 10^{-2}$ b, where for DC and TC (double Compton and triple Compton scattering, respectively), we have assumed that all photons with energies larger than the assumed photon energy threshold $\varepsilon = 50$ MeV are detected. If we adopt pulse parameters available at the LCLS [31], i.e., 2×10^{13} photons per pulse, 10^9 electrons per bunch, a transverse bunch size of $40 \mu\text{m}$, perfect transverse overlap of the two pulses, and a repetition rate of 120 Hz, we obtain 3 triple photon events per second.

One example of the fully differential cross section in the XFEL setup is shown in Fig. 2. In this figure, we assume a configuration centered near the incoming electron axis, with $\theta_j = \pi - 1.5 \times 10^{-3}$ and $\phi_j = 2j\pi/3$, $j = 1, 2, 3$. We evaluate σ_5 as a function of ω_1 and ω_2 . We sum over the electron spins, but assume a linearly polarized XFEL beam, and also fix the polarization of the emitted photons. We show σ_5 only for $\omega_{1,2,3} > \varepsilon$. For photon energies smaller than ε , the differential cross section is set to zero. It becomes clear from Fig. 2 that high-energy photons in the GeV range can be achieved in this setup, which is interesting since it could provide correlated photon triplet states, with quantifiable entanglement as shown below, in an energy range that is far beyond what can be produced with down-conversion of photons in a nonlinear crystal [8–10]. The rate of 3 events/s obtained above is not small compared to the experimental results in [9], where photon triplets produced from nonlinear down-conversion were detected at an event rate of 5 per hour. The polarization




FIG. 2 (color online). The fivefold differential cross section at $\theta_j = \pi - 1.5 \times 10^{-3}$ and $\phi_j = 2j\pi/3$, $j = 1, 2, 3$ as a function of ω_1 and ω_2 . The color indicates the value of the cross section $\log_{10}(\frac{1}{2} \sum_{\text{spin}} \sigma_5 / \text{b MeV}^{-2} \text{sr}^{-3})$ summed over the electron spins. We have $E_i = 5$ GeV and $\omega_0 = 1$ keV, and the incoming photon beam is linearly polarized in the x direction. The different panels correspond to different polarization of the final photons, we have $(\boldsymbol{\epsilon}_1, \boldsymbol{\epsilon}_2, \boldsymbol{\epsilon}_3) = (\boldsymbol{\epsilon}_1^j, \boldsymbol{\epsilon}_2^k, \boldsymbol{\epsilon}_3^l)$, with (a) $jkl = 111$, (b) 211, (c) 121, (d) 112, (e) 221, (f) 212, (g) 122, and (h) 222. The black dashed line in panel (a) shows the curve along which the energy of photon three is at the assumed detector threshold, $\omega_3 = \varepsilon = 50$ MeV.

of GeV photons can be measured using coherent electron-positron pair production in aligned crystals [32].

Entangled photon triplets.—The three photons in the final state are simultaneously emitted and their polarizations are inevitably entangled due to the quantum nature of the process. In general, the mixed polarization state resulting after the interaction can be described by the density matrix ρ , which has matrix elements

$$\langle\lambda_1 \lambda_2 \lambda_3 | \rho | \lambda'_1 \lambda'_2 \lambda'_3 \rangle = N \sum_{r_i, r_f} M(\lambda_1 \lambda_2 \lambda_3) M^*(\lambda'_1 \lambda'_2 \lambda'_3), \quad (8)$$

where we have written $M(\lambda_1 \lambda_2 \lambda_3) = M(\boldsymbol{\epsilon}_1 = \boldsymbol{\epsilon}_1^{\lambda_1}, \boldsymbol{\epsilon}_2 = \boldsymbol{\epsilon}_2^{\lambda_2}, \boldsymbol{\epsilon}_3 = \boldsymbol{\epsilon}_3^{\lambda_3})$, $\lambda_j \in \{1, 2\}$, and the normalization constant N is chosen so that $\text{Tr}(\rho) = 1$ (we sum over initial and final electron spins r_i and r_f). It is still an open problem and subject of active research how to quantify the entanglement of a given multiparticle state ρ (see Refs. [33–35]). In the current study, we employ the entanglement measure put forward in [35] in order to estimate the degree of polarization entanglement present among the final three photons. Briefly, an entanglement witness W is found by minimizing the trace $\text{Tr}(W\rho)$ with W such that for all subsets s , $W = P_s + Q_s^T$, where T_s denotes the partial transpose with respect to the subset s [36], and the matrices P_s , Q_s , $1 - P_s$, and $1 - Q_s$ should have positive eigenvalues. Then, $\tau(\rho) = -\text{Tr}(W\rho)$ is a measure of the tripartite polarization entanglement present in ρ . In particular, $\tau = 0$ for states that are not genuinely tripartite entangled. As an example, $\tau(\rho_{\text{GHZ}}) = 1/2$ for the Greenberger-Horne-Zeilinger (GHZ) state $\rho_{\text{GHZ}} = |\text{GHZ}\rangle\langle\text{GHZ}|$, with




FIG. 3 (color online). Entanglement measure τ as a function of the photon energies ω_1 and ω_2 , for (a) $\omega_0 = 0.662$ MeV; $E_i = m$; $\theta_j = \pi/2 - 1$; $\phi_j = 2j\pi/3$; $j = 1, 2, 3$; $\epsilon = 13$ keV; and (b) $\omega_0 = 1$ keV; $E_i = 5$ GeV; $\theta_j = \pi - 1.5 \times 10^{-3}$; $\phi_j = 2j\pi/3$; $j = 1, 2, 3$; $\epsilon = 50$ MeV. In both cases, spin was summed over and the initial photon beam was assumed to be linearly polarized in the x direction. The differential cross section, and thus also τ , was set to zero for photon energies below the detector threshold ϵ .

$|\text{GHZ}\rangle = (|111\rangle + |222\rangle)/\sqrt{2}$ [37], so that $\tau = 1/2$ may be regarded as maximum entanglement.

In Fig. 3, we show the value of τ , evaluated as a function of ω_1 and ω_2 at (a) $\omega_0 = 0.662$ MeV, $E_i = m$, $\theta_{1,2,3} = \pi/2 - 1$, and $\phi_j = 2j\pi/3$; and (b) $\omega_0 = 1$ keV, $E_i = 5$ GeV, $\theta_{1,2,3} = \pi - 1.5 \times 10^{-3}$, and $\phi_j = 2j\pi/3$ (the same situation as in Fig. 2). The numerical evaluation of τ was performed with PPTMIXER [35], available at [38]. In both cases, a nonzero value of τ for almost all values of ω_1 , ω_2 where $\sigma_5 > 0$, shows that the three photons are indeed entangled. Almost maximum entanglement $\tau = 1/2$ is achieved at certain places in the $\omega_1\omega_2$ plane.

Conclusions.—We have presented a theoretical study of the triple Compton effect, where a photon is split into three after colliding with an electron. Our formulation of the problem enables us to evaluate the differential cross section at arbitrary angles and energies of both initial and final particles. We verify the 44-year-old experimental result reported in Ref. [13]. Theory and experiment are not in perfect agreement but the discrepancy of roughly 1.6 standard deviations is not significant [see Eqs. (6) and (7)]. Additional measurements are needed to clarify this issue. A straightforward generalization of the formalism to a Compton backscattering geometry then leads to theoretical predictions for the triple Compton effect for typical parameters at modern XFEL facilities, as shown in Fig. 2. Finally, while it is intuitively rather clear what a two-particle entangled quantum state is, the quantification of three-particle entanglement is a much more subtle problem. The entanglement measure τ has been proposed in Ref. [35], and corresponding theoretical predictions in Fig. 3 show that the generation of a multiparticle entangled photon state in the MeV or GeV regime is possible based on the triple Compton effect as a basic three-quanta emission

process. The triplet photon final state is polarization entangled.

We acknowledge support from the National Science Foundation Grant No. PHY-1068547 and from the National Institute of Standards and Technology (NIST precision measurement grant).

*Present address: Laser Technology Laboratory, RIKEN Advanced Science Institute, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan.
loetstedte@riken.jp

- [1] O. Klein and T. Nishina, *Z. Phys.* **52**, 853 (1929).
- [2] F. Mandl and T. H. R. Skyrme, *Proc. R. Soc. A* **215**, 497 (1952).
- [3] C. Harvey, T. Heinzl, and A. Ilderton, *Phys. Rev. A* **79**, 063407 (2009).
- [4] F. Mackenroth, A. Di Piazza, and C. H. Keitel, *Phys. Rev. Lett.* **105**, 063903 (2010).
- [5] E. Lötstedt and U. D. Jentschura, *Phys. Rev. Lett.* **103**, 110404 (2009).
- [6] E. Lötstedt and U. D. Jentschura, *Phys. Rev. A* **80**, 053419 (2009).
- [7] D. Seipt and B. Kämpfer, *Phys. Rev. D* **85**, 101701 (2012).
- [8] J. -W. Pan, D. Bouwmeester, M. Daniell, H. Weinfurter, and A. Zeilinger, *Nature (London)* **403**, 515 (2000).
- [9] H. Hübel, D. R. Hamel, A. Fedrizzi, S. Ramelow, K. J. Resch, and T. Jennewein, *Nature (London)* **466**, 601 (2010).
- [10] D. A. Antonosyan, T. V. Gevorgyan, and G. Y. Kryuchkyan, *Phys. Rev. A* **83**, 043807 (2011).
- [11] S. N. Gupta, *Phys. Rev.* **98**, 1502 (1955).
- [12] F. M. Abel, G. S. Adkins, and T. J. Yoder, *Phys. Rev. A* **83**, 062502 (2011).
- [13] M. R. McGie and F. P. Brady, *Phys. Rev.* **167**, 1186 (1968).
- [14] R. C. Majumdar, V. S. Mathur, and J. Dhar, *Nuovo Cimento* **12**, 97 (1959).
- [15] M. R. McGie, F. P. Brady, and W. J. Knox, *Phys. Rev.* **152**, 1190 (1966).
- [16] B. S. Sandhu, R. Dewan, M. B. Saddi, B. Singh, and B. S. Ghuman, *Nucl. Instrum. Methods Phys. Res., Sect. B* **168**, 329 (2000).
- [17] M. B. Saddi, B. Sing, and B. S. Sandhu, *Nucl. Instrum. Methods Phys. Res., Sect. B* **266**, 3309 (2008).
- [18] M. B. Saddi, B. Sing, and B. S. Sandhu, *Nuclear Technology* **175**, 168 (2011).
- [19] M. Ram and P. Y. Wang, *Phys. Rev. Lett.* **26**, 476 (1971); **26**, 1210(E) (1971).
- [20] R. J. Gould, *Astrophys. J.* **285**, 275 (1984).
- [21] V. N. Baier, V. S. Fadin, V. A. Khoze, and E. A. Kuraev, *Phys. Rep.* **78**, 293 (1981).
- [22] A. A. Krylovetskiĭ, N. L. Manakov, S. I. Marmo, and A. F. Starace, *Zh. Eksp. Teor. Fiz.* **122**, 1168 (2002) [*Sov. Phys. JETP* **95**, 1006 (2002)].
- [23] A. V. Korol and I. A. Solovjev, *Radiat. Phys. Chem.* **75**, 1346 (2006).
- [24] T. Radtke, A. Surzhykov, and S. Fritzsche, *Eur. Phys. J. D* **49**, 7 (2008).

- [25] U.D. Jentschura and A. Surzhykov, *Phys. Rev. A* **77**, 042507 (2008).
- [26] A. Di Piazza, A.I. Milstein, and C.H. Keitel, *Phys. Rev. A* **76**, 032103 (2007).
- [27] C. Itzykson and J.B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980).
- [28] J.M. Jauch and F. Rohrlich, *The Theory of Photons and Electrons* (Springer, Heidelberg, 1980), 2nd ed..
- [29] L.M. Brown and R.P. Feynman, *Phys. Rev.* **85**, 231 (1952).
- [30] S. Schnez, E. Lötstedt, U.D. Jentschura, and C.H. Keitel, *Phys. Rev. A* **75**, 053412 (2007).
- [31] <http://lcls.slac.stanford.edu>.
- [32] A. Apyan *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. B* **234**, 128 (2005).
- [33] G. Tóth, O. Gühne, M. Seevinck, and J. Uffink, *Phys. Rev. A* **72**, 014101 (2005).
- [34] J.-D. Bancal, N. Gisin, Y.-C. Liang, and S. Pironio, *Phys. Rev. Lett.* **106**, 250404 (2011).
- [35] B. Jungnitsch, T. Moroder, and O. Gühne, *Phys. Rev. Lett.* **106**, 190502 (2011).
- [36] M. Lewenstein, B. Kraus, J.I. Cirac, and P. Horodecki, *Phys. Rev. A* **62**, 052310 (2000).
- [37] D.M. Greenberger, M.A. Horne, A. Shimony, and A. Zeilinger, *Am. J. Phys.* **58**, 1131 (1990).
- [38] <http://www.mathworks.com/matlabcentral/fileexchange/30968>.