



Missouri University of Science and Technology
Scholars' Mine

Biological Sciences Faculty Research & Creative Works

Biological Sciences

01 Sep 2016

Conversion of Glycerol to 1,3-propanediol under Haloalkaline Conditions

Melanie R. Mormile

Missouri University of Science and Technology, mmormile@mst.edu

Daniel William Roush

Dwayne A. Elias

Oliver C. Sitton

Missouri University of Science and Technology, ocs@mst.edu

Follow this and additional works at: https://scholarsmine.mst.edu/biosci_facwork



Part of the [Biology Commons](#), and the [Chemical Engineering Commons](#)

Recommended Citation

M. R. Mormile et al., "Conversion of Glycerol to 1,3-propanediol under Haloalkaline Conditions," *U.S. Patents*, Sep 2016.

This Patent is brought to you for free and open access by Scholars' Mine. It has been accepted for inclusion in Biological Sciences Faculty Research & Creative Works by an authorized administrator of Scholars' Mine. This work is protected by U. S. Copyright Law. Unauthorized use including reproduction for redistribution requires the permission of the copyright holder. For more information, please contact scholarsmine@mst.edu.



US009328360B2

(12) **United States Patent**
Mormile et al.

(10) **Patent No.:** **US 9,328,360 B2**
(45) **Date of Patent:** **May 3, 2016**

(54) **CONVERSION OF GLYCEROL TO 1,3-PROPANEDIOL UNDER HALOALKALINE CONDITIONS**

8,034,592 B2 * 10/2011 Elias et al. 435/168
2003/0022323 A1 1/2003 Dunn-Coleman et al.
2010/0028965 A1 2/2010 Liu et al.
2011/0136196 A1 * 6/2011 Elias et al. 435/168
2013/0177956 A1 7/2013 Figge

(71) Applicant: **The Curators of the University of Missouri**, Columbia, MO (US)

OTHER PUBLICATIONS

(72) Inventors: **Melanie Rose Mormile**, Rolla, MO (US); **Daniel William Roush**, Chandler, AZ (US); **Dwayne Alexander Elias**, Knoxville, TN (US); **Oliver Clifford Sitton**, Rolla, MO (US)

Kivisto et al., Journal of Biotechnology 158:242-247, available online Nov. 6, 2011.*
Brown et al., Genbank accession No. CP002304, 2011.*
The 2012 ATCC catalog.*
Van Gerpen, J., Improvement of Crop Plants for Industrial End Uses, Chapter 10, pp. 281-289, P. Ranalli Editor, 2007.*
Cayol et al., Extremophiles 6:131-134, 2002.*
Croft et al., Nature 438:90-93, 2005.*
Ohwada et al., Limnology and Oceanography 17(2):315-320, 1972.*
Kivisto et al., Bioresource Technology 101:8671-8677, 2010.*
Kivisto et al., "Non-sterile process for biohydrogen and 1,3-propanediol production from raw glycerol," International Journal of Hydrogen Energy, Jul. 26, 2013, vol. 38, Issue No. 27, 11749-11755. International Search Report and Written Opinion mailed Dec. 23, 2014 in corresponding PCT/US2014/054434 filed Sep. 6, 2014.
Brown et al., "Complete Genome Sequence of the Haloalkaliphilic, Hydrogen-Producing Bacterium Halanaerobium hydrogeniformans," Journal of Bacteriology, Jul. 2011, vol. 193, No. 14, 3682-3683.
Kivisto et al., "Halophilic anaerobic fermentative bacteria," Journal of Biotechnology 152 (2011) 114-124.
McAdams et al., "Stochastic mechanisms in gene expression", Proc. Natl. Acad. Sci. USA 94 (1997) 814-819, vol. 94.

(73) Assignee: **The Curators of the University of Missouri**, Columbia, MO (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/479,292**

(22) Filed: **Sep. 6, 2014**

(65) **Prior Publication Data**

US 2015/0072388 A1 Mar. 12, 2015

Related U.S. Application Data

(60) Provisional application No. 61/874,752, filed on Sep. 6, 2013.

(51) **Int. Cl.**
C12P 7/18 (2006.01)

(52) **U.S. Cl.**
CPC **C12P 7/18** (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,686,276 A * 11/1997 Laffend et al. 435/158
7,582,457 B2 9/2009 Dunn-Coleman et al.

* cited by examiner

Primary Examiner — Delia Ramirez
(74) *Attorney, Agent, or Firm* — Hovey Williams LLP

(57) **ABSTRACT**

A method of producing 1,3-propanediol. The method comprises fermenting a haloalkaliphilic species of *Halanaerobium* with a source of glycerol into 1,3-propanediol, at a pH of greater than about 10 and at a salt concentration of greater than about 5% w/v. Furthermore, with supplementation of vitamin B₁₂, the yield of 1,3-propanediol to glycerol can be increased.

20 Claims, 6 Drawing Sheets




Fig. 1






Fig. 2



$Peak\ Height = 281 + 536\ Conc$
 $Conc = (Peak\ Height - 281) / 536$
 $R^2 = 99.9\% \text{ Adj.} | R^2 = 99.9\% | P\text{-value} = <0.001$


Fig. 3



$Peak\ Area = 4617 + 9403\ Conc$
 $Conc = (Peak\ Area - 4617) / 9403$
 $R^2 = 99.9\% \text{ Adj.} | R^2 = 99.9\% | P\text{-value} = <0.001$

Fig. 4


1,3-propanediol



Peak Height = 180 + 350 Conc. (mM)
 Conc = (Peak Height - 180) / 350
 R² = 99.9% Adj. | R² =99.8% | P-value = <0.001

Fig. 5

1,3-propanediol



Peak Area = 4492 + 7661 Conc. (mM)
 Conc = (Peak Height - 4492) / 7661
 R² = 99.8% Adj. | R² =99.7% | P-value = <0.001

Fig. 6




Fig. 7

1,3-propanediol Concentration Comparison




Fig. 8




Fig. 9

1

CONVERSION OF GLYCEROL TO 1,3-PROPANEDIOL UNDER HALOALKALINE CONDITIONS

CROSS-REFERENCE TO RELATED APPLICATIONS

The present application claims the priority benefit of U.S. Provisional Patent Application Ser. No. 61/874,752, filed Sep. 6, 2013, entitled CONVERSION OF GLYCEROL TO 1,3-PROPANEDIOL UNDER HALOALKALINE CONDITIONS, incorporated by reference in its entirety herein.

SEQUENCE LISTING

The following application contains a sequence listing in computer readable format (CRF), submitted as a text file in ASCII format entitled "SequenceListing," created on Aug. 28, 2014, as 20 KB. The content of the CRF is hereby incorporated by reference.

BACKGROUND

1. Field of the Invention

This invention will provide a way for biodiesel companies to form a valuable product, (1,3-propanediol) from a waste product (glycerol).

2. Description of Related Art

Chemical waste can be recycled into useful compounds. With the recent surge in biodiesel production, glycerol has gone from a relatively rare commodity to a heavily overproduced waste product. Many major chemical and agriculture companies have been attempting to find high yielding conversions for glycerol. One of the major processes is the conversion of glycerol into 1,3-propanediol by way of microbial metabolism. There has been success in identifying strains of microorganisms that can conduct this reaction; however it may not be commercially feasible as the raw glycerol product needs to be treated. For economic feasibility, the process must be able to convert the raw glycerol product into 1,3-propanediol with limited treatment. For example, 1,3-propanediol, is used frequently in the chemical industry as a building block for many common products, like adhesives, fragrances and perfumes, personal care products, and coatings like paint. Currently, 1,3-propanediol is synthesized from components of crude oil, propylene or ethylene oxide, or glucose derived from corn to synthesize. However, common chemical processes for recycling chemical waste involve making the processing streams more tolerable to bacteria for biological conversion. By adding large amounts of acids or bases, or using large amounts of energy to remove salts and impurities, industries make the conditions suitable for non-extremophilic life. Glycerol is another common waste product of biodiesel production that can be converted into useful compounds. With the recent surge in biodiesel production, glycerol has gone from a relatively rare commodity to a heavily overproduced waste product. Many major chemical and agriculture companies have been attempting to find high yielding conversions for glycerol. One of the major process targets is the conversion of glycerol into 1,3-propanediol by way of microbial metabolism (FIG. 1). There has been success in identifying strains of microorganisms that can conduct this reaction; however it may not be commercially feasible as the raw glycerol product needs to be treated. Glycerol acts very much like salt, in the sense that it increases the pressure put onto the bacteria. For economic feasibility, the process must be able to convert the raw glycerol product into 1,3-

2

propanediol with limited treatment. Accordingly, there remains a need for improved approaches to converting chemical waste into useful compounds and products.

SUMMARY OF THE INVENTION

The invention addresses the problems above by providing a method of producing 1,3-propanediol. The method comprises fermenting a species of *Halanaerobium* with a source of glycerol, whereby 1,3-propanediol is produced. Fermentation can be carried out under high pH and high salt concentrations, and without the removal of impurities from the glycerol feedstock. Fermentative conversion rates can be increased by supplementing the fermentation culture with vitamin B₁₂.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 illustrates the process of converting glycerol to 1,3-propanediol;

FIG. 2 is a scanning electron microscope (SEM) photograph of the haloalkaliphilic species of *Halanaerobium*;

FIG. 3 is a scatterplot graph of glycerol consumption;

FIG. 4 is a scatterplot graph of glycerol consumption;

FIG. 5 is a scatterplot graph of 1,3-propanediol production;

FIG. 6 is a scatterplot graph of 1,3-propanediol production;

FIG. 7 shows scatterplot graphs showing glycerol concentration comparison between the Peek Height and the Peek Area;

FIG. 8 shows scatterplot graphs showing 1,3-propanediol concentration comparison between the Peek Height and the Peek Area; and

FIG. 9 is a graph showing the effect of increasing concentrations of vitamin Bit on 1,3-propanediol production (conversion).

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention provides a process for the fermentative production of 1,3-propanediol without the currently required steps of pretreating, desalination, or the neutralization to decrease the salinity or pH of the feedstock or fermentation media, or remove waste by-products typically present in the crude feedstock. The inventive process is amenable to larger scale, commercial or industrial applications for the production of 1,3-propanediol as a useful precursor material from microbial fermentation of chemical waste, and specifically a chemical waste feedstock comprising glycerol or a source of glycerol in an inexpensive and environmentally friendly manner.

The inventive methods utilize extremophilic microorganisms that can thrive in the presence of untreated chemical waste. Specifically the preferred organism, *Halanaerobium hydrogeniformans* (ATCC Patent Deposit Designation No. PTA-10410, deposited Oct. 13, 2009), grows in conditions with high pH (about pH 11 which is similar to laundry detergent) and high salt concentrations (7% salt, double that of the ocean). These conditions are found in treated biomass used to produce biofuels like ethanol and hydrogen, and crude glycerol generated during biodiesel production.

Fermentation of the chemical waste feedstock is accomplished with a haloalkaliphilic microorganism capable of 1,3-propanediol production under highly alkaline and hypersaline conditions. *H. hydrogeniformans* is able to convert glycerol, a common waste product of biodiesel production, into 1,3-propanediol under extreme conditions, of pH 11 and

7% salt. In preferred embodiments, a conversion rate of about 55% can be achieved with the process. Advantageously, the microorganism can also grow in media containing up to 1M glycerol and can thrive in a solution containing crude waste glycerol.

Unlike other members of the *Halanaerobium* genus, the microorganism is highly alkaliphilic with optimum growth at a pH of from about 10.5 to about 11. Suitable microorganisms for use with the inventive method preferably have a 16S ribosomal DNA (rDNA) sequence comprising (or consisting of) SEQ ID NO: 1, or a 16S rDNA sequence having at least 98% sequence homology with SEQ ID NO: 1, and more preferably at least 99% sequence homology with SEQ ID NO: 1. Suitable microorganisms will preferably have at least one gene encoding for glycerol dehydratase or an enzyme having glycerol dehydratase activity, and preferably an endogenous gene encoding for glycerol dehydratase or an enzyme having glycerol dehydratase activity. In one or more embodiments, the microorganisms comprise an endogenous DNA sequence comprising (or consisting of) SEQ ID NO:2 or a sequence having at least 98% sequence homology with SEQ ID NO: 2, and more preferably at least 99% sequence homology with SEQ ID NO: 2. In one or more embodiments, the microorganisms comprise a gene encoding for an endogenous protein comprising (or consisting of) SEQ ID NO:3, or a sequence having at least 98% sequence homology with SEQ ID NO: 3, and more preferably at least 99% sequence homology with SEQ ID NO: 3.

Suitable microorganisms will preferably have at least one gene encoding for iron-containing alcohol dehydrogenase or an enzyme having alcohol dehydrogenase activity, and preferably an endogenous gene encoding for iron-containing alcohol dehydrogenase or an enzyme having alcohol dehydrogenase activity. In one or more embodiments, the microorganisms comprise an endogenous DNA sequence comprising (or consisting of) SEQ ID NO:4 or a sequence having at least 98% sequence homology with SEQ ID NO: 4, and more preferably at least 99% sequence homology with SEQ ID NO: 4. In one or more embodiments, the microorganisms comprise a gene encoding for an endogenous protein comprising (or consisting of) SEQ ID NO:5, or a sequence having at least 98% sequence homology with SEQ ID NO: 5, and more preferably at least 99% sequence homology with SEQ ID NO: 5. In one or more embodiments, the microorganisms comprise an endogenous DNA sequence comprising (or consisting of) SEQ ID NO:6 or a sequence having at least 98% sequence homology with SEQ ID NO: 6, and more preferably at least 99% sequence homology with SEQ ID NO: 6. In one or more embodiments, the microorganisms comprise a gene encoding for an endogenous protein comprising (or consisting of) SEQ ID NO:7, or a sequence having at least 98% sequence homology with SEQ ID NO: 7, and more preferably at least 99% sequence homology with SEQ ID NO: 7. Suitable microorganisms include mutants and derivatives (progeny) of the microorganism which retain the haloalkaliphilic properties *H. hydrogeniformans*. Mutants (such as by deletion, insertion, and/or substitution of a base in the above-referenced sequences) include those occurring spontaneously in the passage or cultivation of the organism, as well as intentional mutations. In one or more embodiments, haloalkaliphilic microorganisms can also be used, which have been engineered to contain one or more of the genes referenced above or a gene encoding for one or more of the enzymes referenced above.

The chemical waste feedstock comprising glycerol or a source of glycerol is fermented with the microorganism in a culture medium under conditions suitable for 1,3-pro-

panediol production. A preferred culture medium comprises, consists essentially, or even consists of (per liter): 70 g NaCl, 40 g Na₂CO₃, 6.3 g K₂HPO₄, 1 g yeast extract, 0.75 g Na₂S, and 0.6 g cysteine, along with 10 ml of basal medium stock solution and 10 ml of trace mineral solution. The basal medium stock solution preferably comprises 50 mg NH₄NO₃, 8.5 mg MgCl₂·6H₂O, 7.5 mg SiO₂, 4.5 mg MnSO₄·H₂O, 4.2 mg CaCl₂·2H₂O, 4 mg methylene blue, and 1.8 mg FeSO₄·7H₂O. The trace mineral solution preferably comprises (per liter): 3 g MgSO₄·7H₂O, 1.63 g Na₃-NTA, 1 g NaCl, 0.64 g MnCl₂·4H₂O, 0.13 g ZnCl₂, 0.1 g FeSO₄·7H₂O, 0.1 g CaCl₂·2H₂O, CoCl₂·6H₂O, 0.03 g NiSO₄·6H₂O, 0.025 g Na₂MoO₄·2H₂O, 0.025 g Na₂WO₄·2H₂O, 0.01 g AIK (SO₄)₂·12H₂O, 0.01 g H₃BO₃, and 7 mg CuCl₂·2H₂O.

The chemical waste feedstock comprising glycerol or the glycerol source is preferably provided at a glycerol concentration of from about 1 g/L to about 184 g/L, and preferably from about 10 g/L to about 92 g/L. The microorganism can ferment feedstocks with a glycerol concentration as high as 184 g/L. The microorganism ferments the feedstock to generate 1,3-propanediol along with other by-products.

As mentioned, the method is preferably carried out without neutralization of the chemical waste feedstock (i.e., without decreasing the pH to about 7). That is, the pH of the feedstock (and resulting fermenting culture, including the culture media) is preferably greater than or equal to about 10, preferably from about 10 to about 11, and more preferably from about 10.5 to about 11. The salinity (% NaCl content) of the feedstock and fermenting culture (including the culture media) is also preferably greater than or equal to about 5% w/v, more preferably greater than or equal to about 7% w/v, and even more preferably from about 7% to about 7.5% w/v. As used herein, the percentage "weight by volume" of the component in the composition (referred to herein as "% w/v") is calculated based upon the total mass of the component (e.g., salt) in grams per liter of the final solution where 1000 g/L is taken as 100% w/v. These pH and salinity conditions are preferably maintained in the culture medium throughout the fermentation process. That is, the pH of the fermenting culture preferably remains at or above about pH 10, and more preferably from about 10.5 to about 11, while the salinity remains greater than or equal to about 5% w/v, preferably greater than or equal to 7% w/v, and more preferably from about 7% to about 7.5% w/v.

Preferably, the feedstock is not purified or pretreated. In one or more embodiments, the feedstock will also comprise methanol, crude glycerol, sodium hydroxide, water, and mixtures thereof. Methanol is commonly present in the biodiesel waste stream along with glycerol. Advantageously, the microorganism has a tolerance to raw biodiesel waste. Accordingly, in one or more embodiments, such impurities or chemicals are not removed from the feedstock prior to fermentation. The microorganism is expected to form 1,3-propanediol from the glycerol in untreated biodiesel waste, which will save producers from having to remove methanol and other possible contaminants that would typically inhibit less tolerant fermentative bacteria.

In one or more embodiments, the fermentation culture is preferably supplemented with vitamin B₁₂. Preferably, vitamin B₁₂ is present in the fermentation culture at a level of from about 25 µg/L to about 100 µg/L, more preferably from about 25 µg/L to about 75 µg/L, and even more preferably from about 25 µg/L to about 50 µg/L. Advantageously, the endogenous glycerol dehydratase of *H. hydrogeniformans* is not necessarily dependent on B₁₂, and can ferment glycerol to

1,3-propanediol without B₁₂ supplementation. However, it has been shown that B₁₂ can enhance the yield, as much as 0.47 (mol/mol).

In one or more embodiments, fermentation is preferably carried out under substantially anaerobic conditions. As used herein, “substantially anaerobic conditions” refers to conditions where there no free oxygen available (e.g., less than about 0.1 ppm free oxygen, preferably from about 0 to about 0.1 ppm free oxygen), and includes naturally or artificially oxygen-depleted environments. More preferably, for artificial environments (i.e., test tube, fermentation reactor) a gas phase is provided in the headspace above the culture medium, with suitable gases being selected from the group consisting of N₂, CO₂, and mixtures thereof. A particularly preferred gas phase is a combination of about 80% N₂/20% CO₂. In a preferred method, the substantially anaerobic conditions can be maintained by sparging the culture medium with the selected gases.

The culture medium is also preferably agitated during fermentation, preferably at speeds of from about 100 rpm to about 250 rpm, and more preferably from about 100 rpm to about 200 rpm. Agitation can be accomplished via shaking, rotation, impeller, or any combination thereof. Fermentation also preferably proceeds in the absence of light (i.e., the culture is not exposed to any light sources during the fermentation process). Fermentation is preferably carried out at a temperature of from about 6° C. to about 40° C., and preferably from about 25° C. to about 30° C., and for time periods of from about 12 hours to about 72 hours, and preferably for time periods of from about 12 hours to about 24 hours.

The fermentation process preferably results in a percent mole/mole yield of 1,3-propanediol from glycerol of at least about 32% with a glycerol-only medium, and preferably from about 32% to about 60% and a theoretical yield over 90%. With a vitamin B₁₂ amended medium, the yield of 1,3-propanediol from glycerol is preferably greater than about 60%, and preferably from about 60% to about 80% and a theoretical yield over 90%. The yield is calculated by:

$$\frac{(\text{mol 1,3-propanediol produced})}{(\text{mol glycerol initial})} = \text{Yield}$$

or

$$\frac{(\text{mol 1,3-propanediol produced})}{(\text{mol glycerol initial with added } B_{12})} = \text{Yield}$$

In one or more embodiments, the process has a yield of 1,3-propanediol of about 60% at about pH 11 and a media containing about 7% salt and about 0.2% glycerol, with added B₁₂.

Fermentation can be carried out in a fermentation apparatus (fermentation reactor). Suitable fermentation reactors are known in the art. In general, suitable apparatuses will have inlets for the biodiesel waste feedstock, gas for artificial atmosphere, and a fermentation chamber, and outlets for removing the 1,3-propanediol and by-products. The apparatus will contain the microorganism and nutrient culture medium. The apparatus can also be equipped with a stir bar, impeller or other agitation device. The feedstock may be continuously supplied to the fermentation apparatus as needed to keep up with the rate of fermentation of the chemical waste substrate. The fermentation apparatus may be a stand-alone apparatus, or it may be combined with a downstream reactor for receiving and further processing any by-products from the fermentation apparatus.

The process further comprises recovering the produced 1,3-propanediol from the fermentation reaction. The resulting 1,3-propanediol can be separated from the fermentation culture, such as by distillation, extraction, or other separation method.

In yet a further embodiment, the waste stream from the fermentation reactor is recycled and reintroduced into the system. This is feasible in the inventive process because the salt concentration and pH of the waste stream would still be amenable to microbial cultivation using the extremophilic microorganism. The pH or salt concentration may be adjusted (upwards), if necessary. Advantageously, this significantly reduces not only the amount of water required for the process, but the cost of the substrates for the cultivation and thereby the overall cost of the production of 1,3-propanediol.

The benefits and novelty of our process is that the microorganism can convert glycerol to 1,3-propanediol under alkaline conditions without the need to neutralize the raw glycerol to a pH value of 7.0. In addition, the microorganism is halotolerant and can withstand saline conditions. Typically, raw glycerol wastes have a salinity of ~5%. With the microorganism, there is no need to dilute the residual salt in the waste. The use of the microorganism will help to streamline the process of glycerol conversion to 1,3-propanediol. The competitive advantage is that the biodiesel waste stream will not have to be treated to remove the salts or adjust its pH.

Additional advantages of the various embodiments of the invention will be apparent to those skilled in the art upon review of the disclosure herein and the working examples below. It will be appreciated that the various embodiments described herein are not necessarily mutually exclusive unless otherwise indicated herein. For example, a feature described or depicted in one embodiment may also be included in other embodiments, but is not necessarily included. Thus, the present invention encompasses a variety of combinations and/or integrations of the specific embodiments described herein.

The present description also uses numerical ranges to quantify certain parameters relating to various embodiments of the invention. It should be understood that when numerical ranges are provided, such ranges are to be construed as providing literal support for claim limitations that only recite the lower value of the range as well as claim limitations that only recite the upper value of the range. For example, a disclosed numerical range of about 10 to about 100 provides literal support for a claim reciting “greater than about 10” (with no upper bounds) and a claim reciting “less than about 100” (with no lower bounds).

EXAMPLES

The following examples set forth methods in accordance with the invention. It is to be understood, however, that these examples are provided by way of illustration and nothing therein should be taken as a limitation upon the overall scope of the invention.

Example 1

Halanaerobium hydrogeniformans

Halanaerobium hydrogeniformans (formerly *Halanaerobium strain sapolanicus*) was isolated from haloalkaline (pH~10, 15- to 140-g/liter NaCl), anaerobic sediments of Soap Lake, Wash., with extraordinarily high sulfide concentrations of up to 10 g/liter. It is an obligately anaerobic, Gram-negative, nonmotile, nonsporulating, elongated rod

bacterium (FIG. 2). It can utilize a range of C₅ and C₆ sugars with optimal growth at pH 11, 7% (wt/vol) NaCl, and 33° C., producing acetate, formate, and hydrogen as major metabolic end products. The genome sequence for *H. hydrogeniformans* was determined to improve assessment of its metabolic and bioenergy potential, particularly toward improving alkaline or haloalkaline pretreatment regimens for robust hydrogen production by this bacterium. The *H. hydrogeniformans* genome sequence was determined through a combination of Illumina (Bennett, S. 2004. Solexa, Ltd. Pharmacogenomics 5:433-438) and 454 (Margulies, M., et al. 2005. Genome sequencing in microfabricated high density picoliter reactors. Nature 437:376-380.) technologies. The Joint Genome Institute constructed and sequenced an Illumina GAii shotgun library which generated 27,639,916 reads totaling 2,100 Mb, a 454 Titanium standard library generated from 77,351 reads, and a paired-end 454 library with an average insert size of 10.607±2.651 kb that generated 160,293 reads totaling 82.3 Mb of 454 data. A total of 486 additional reactions and 6 shatter libraries were necessary to close gaps and to raise the finished sequence quality. Methods for determining the genome sequence were previously described (Elkins, J. G., et al. 2010. Complete genome sequence of the cellulolytic thermophile *Caldicellulosiruptor obsidiansis* OB47T. J. Bacteriol. doi:10.1128/JB.00950-10), and this is a “finished” genome (Chain, P. S. G., et al. 2009. Genome project standards in a new era of sequencing. Science 326:236-237). The total genome size was 2,613,116 bp, with final assembly based on 52.2 Mb of 454 draft data providing an average 21.5X genome coverage and 463 Mb of Illumina draft data providing an average 178X genome coverage. The genome is 33.1% G+C and contains 2,295 candidate protein-encoding gene models. The genome contains four separate rRNA operons, each containing a 5S, a 16S (SEQ ID NO:1), and a 23S rRNA gene, with 99.9 to 100% identity between 16S rRNA genes. The closest significant 16S rRNA gene matches (GenBank accession number GQ215697) were to other *Haloanaerobium* species. However, all comparative species are physiologically different as they are neutrophilic. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number CP002304.

Example 2

Production of 1,3-propanediol from Glycerol

The culture medium included (per liter): 70 g NaCl, 40 g Na₂CO₃, 6.3 g K₂HPO₄, 1 g yeast extract, 0.75 g Na₂S (as a reductant), 0.6 g cysteine (as a reductant), along with 10 ml of basal medium stock solution and 10 ml of trace mineral solution. The basal medium stock solution included (per liter): 50 mg NH₄NO₃, 8.5 mg MgCl₂·6H₂O, 7.5 mg SiO₂, 4.5 mg MnSO₄·H₂O, 4.2 mg CaCl₂·2H₂O, 4 mg methylene blue (as an oxygen indicator), and 1.8 mg FeSO₄·7H₂O. The trace mineral solution included (per liter): 3 g MgSO₄·7H₂O, 1.63 g Na₃-NTA, 1 g NaCl, 0.64 g MnCl₂·4H₂O, 0.13 g ZnCl₂, 0.1 g FeSO₄·7H₂O, 0.1 g CaCl₂·2H₂O, 0.1 g CoCl₂·6H₂O, 0.03 g NiSO₄·6H₂O, 0.025 g Na₂MoO₄·2H₂O, 0.025 g Na₂WO₄·2H₂O, 0.01 g AlK(SO₄)₂·12H₂O, 0.01 g H₃BO₃, and 7 mg CuCl₂·2H₂O.

The culture bottles were prepared with 50 mL of culture medium and then amended with 2.5 mL of 600 mM Glycerol stock solution (to a final concentration of ~25 mM glycerol). Culture bottles were also amended with 2.5 mL of a 128 µg/mL Vitamin B₁₂ solution (to a final concentration of 53.33 µg/L). The headspace gas was exchanged to contain 100% N₂. The samples were incubated at 30° C. in a shaking incu-

bator at 150 rpm for seven days. The results are shown in Table 1 below, from cultures that were amended with glycerol. Three replicates were amended with glycerol and bacteria. One culture amended with glycerol was not inoculated with the bacterium. Three additional replicates were amended with glycerol and vitamin B₁₂. One culture amended with glycerol and vitamin B₁₂ was not inoculated with the bacterium. These results clearly demonstrate that the bacterium consumed the glycerol amendments. Those cultures that also were amended with vitamin B₁₂ were able to consume a greater amount of glycerol than those that were not amended with this vitamin.

TABLE 1

Treatment	Day 0 Concentration of Glycerol (mM)	Day 7 Concentration of Glycerol (mM)	
Replicate #1	Glycerol	26.706	8.047
Replicate #2	Glycerol	26.565	7.603
Replicate #3	Glycerol	26.446	7.939
Without bacteria	Glycerol	27.552	27.631
Replicate #1	Glycerol + B ₁₂	25.458	0.870
Replicate #2	Glycerol + B ₁₂	25.264	1.170
Replicate #3	Glycerol + B ₁₂	25.484	1.118
Without bacteria	Glycerol + B ₁₂	26.331	26.546

Table 2 shows the results from cultures that were amended with glycerol. Three replicates were amended with glycerol and bacteria. One culture amended with glycerol was not inoculated with the bacterium. Three additional replicates were amended with glycerol and vitamin B₁₂. One culture amended with glycerol and vitamin B₁₂ was not inoculated with the bacterium. These results clearly demonstrate that the bacterium is capable of producing 1,3-propanediol. Those cultures that also were amended with vitamin B₁₂ were able to produce a greater amount of 1,3-propanediol than those that were not amended with this vitamin.

TABLE 2

Treatment	Day 0 Concentration of 1,3-propanediol (mM)	Day 7 Concentration of 1,3-propanediol (mM)	
Replicate #1	Glycerol	Not detected	1.172
Replicate #2	Glycerol	Not detected	1.104
Replicate #3	Glycerol	Not detected	1.203
Without bacteria	Glycerol	Not detected	Not detected
Replicate #1	Glycerol + B ₁₂	Not detected	11.947
Replicate #2	Glycerol + B ₁₂	Not detected	11.619
Replicate #3	Glycerol + B ₁₂	Not detected	11.786
Without bacteria	Glycerol + B ₁₂	Not detected	Not detected

Discussion

Standard Curves are shown in FIGS. 3-6. The differences between day 0 and day 7 glycerol and glycerol+B₁₂ treatment groups were analyzed. Glycerol consumption and 1,3-propanediol production were examined. Acetate production was also examined to determine the activity of the glycerol kinase pathway compared to the glycerol dehydratase pathway. Without protein analysis an exact measure of growth was not available, but acetate production can indicate at the very least that fermentation occurred and an estimate of how much glycerol is being utilized for pyruvate metabolism instead of 1,3-propanediol production.

The scatterplots in FIG. 7 shows that the cultures started with approximately the same concentration of glycerol, how-

ever after 7 days the cultures that were supplemented with B₁₂ (right hand side, 5-8000) utilized more of the total glycerol.

The scatterplots in FIG. 8 show that 1,3-propanediol production from the bacterium was observed under extreme conditions and increased production when B₁₂ is supplemented to the organism. With respect to acetate to examine "growth" roughly along with the activity of normal metabolism, both Peak Area and Peak Height in the B₁₂ supplemented cultures are about half of what is in just glycerol cultures which may help explain the decreased growth. A quick paired T test was performed to make sure the concentration differences were significant in both the Glycerol and Glycerol+B₁₂ cultures. Both p-values were <0.001 indicating a statistically significant production of 1,3-propanediol.

The final glycerol concentrations in the bottles at Day 0 was about 25.3 mM and we were producing about 12 mM 1,3-propanediol, resulting in about a 0.47 mol to mol ratio, however the B₁₂ supplementation was <64 µg/mL due to dilutions from inoculum and carbon source addition.

Example 3

Requirement of B₁₂

Anaerobic cultures were prepared in 160 mL serum bottles. The medium was prepared by boiling to degas under a N₂ blanket. As the medium cooled, reductant stock mix was added to the media that contained 0.75 g Na₂S and 0.6 g cysteine per liter. Once the media was cooled, the flasks were transferred into a Coy anaerobic glove bag where the 50 mL of media was dispensed into 160 mL serum bottles filled and autoclaved (121° C., 20 min). After autoclaving, the headspace gas was exchanged for 80% N₂/20% CO₂ mixture. The bottles then were inoculated with a 10% inoculum from previous stock cultures. 30 mM glycerol was added. Vitamin B₁₂ supplementation from anaerobic, filter-sterilized stocks were added right before inoculation at 0 µg/L, 25 µg/L, 50 µg/L, 75 µg/L, and 100 µg/L.

Samples were taken every 24 hours. 5 mL syringes were degassed with N₂/CO₂ mix, and 1 mL of culture sample was removed for each of the sample periods. The sample was placed in a 1.5 mL Eppendorf tube, and centrifuged for 5 min at 13,000×g. The supernatant was decanted into another 1.5 mL Eppendorf tube, and frozen for HPLC analysis.

For HPLC analysis, filter sterilized samples (0.45 µM PTFE filters) were injected onto a 300×7.8 mm Aminex HPX-87H column (BioRad, Hercules, Calif.) maintained at 50° C. The mobile phase was 2.5 mM H₂SO₄ maintained at a constant flow rate of 0.6 ml/min and at approximately 2.2 MPa. Detection was done with both a UV 231 (at 210 nm) and refractive index monitor.

Results Obtained:

The production capabilities of *H. hydrogeniformans* and the influence of vitamin B₁₂ supplementation were studied. A gradient was prepared to examine the maximum production of 1,3-propanediol from media containing 30 mM glycerol. Approximately 16.5 mM 1,3-propanediol was produced when the culture was amended with 25, 50, 75, or 100 µg/L vitamin B₁₂ and approximately 8.5 mM 1,3-propanediol when no vitamin B₁₂ was provided (FIG. 9). Table 3 indicates the percent mole/mole conversion of glycerol to 1,3-propanediol in *H. hydrogeniformans* cultures when supplemented with vitamin B₁₂.

TABLE 3

Percent mole/mole conversion of glycerol to 1,3-propanediol in <i>H. hydrogeniformans</i> cultures supplemented with vitamin B ₁₂ .	
B ₁₂ Amendment, µg/L	% mol 1,3-propanediol/mol glycerol
0	31.5
25	59.1
50	60.3
75	60.1
100	60.2

Example 4

Tolerance of *H. hydrogeniformans* to Glycerol

The tolerance of *H. hydrogeniformans* to concentrations of glycerol was examined. Anaerobic cultures were prepared in 160 mL serum bottles. The medium was prepared by boiling to degas under a N₂ blanket. As the medium cooled, reductant stock mix was added to the media that contained 0.75 g Na₂S and 0.6 g cysteine per liter. Once the media was cooled, the flasks were transferred into a Coy anaerobic glove bag where the 50 mL of media was dispensed into 160 mL serum bottles filled and autoclaved (121° C., 20 min). After autoclaving, the headspace gas was exchanged for 80% N₂/20% CO₂ mixture. The bottles then were inoculated with a 10% inoculum from previous stock cultures. Sterilized glycerol was added to the serum bottles to achieve 7.5, 15, 30, 60, 120, 240, 480, 960, and 1920 mM. Growth was examined by turbidity readings taken at 600 nM. It was determined that *H. hydrogeniformans* was capable of growth at 7.5, 15, 30, 60, 120, 240, 480, 960, and 1920 mM glycerol. It did not exhibit any growth when glycerol was not present in the medium. The data indicate that *H. hydrogeniformans* can tolerate at least 1M glycerol in addition to 7% (w/v) and pH 11.

TABLE 4

	Glycerol Concentration (mM)									
	0	7.5	15	30	60	120	240	480	960	1920
Growth	X	X	X	X	X	X	X	X	X	—

Example 5

Tolerance of *H. hydrogeniformans* to 1,3-propanediol

The tolerance of *H. hydrogeniformans* to increasing concentrations of 1,3-propanediol was examined. Anaerobic cultures were prepared in 160 mL serum bottles. The medium was prepared by boiling to degas under a N₂ blanket. As the medium cooled, reductant stock mix was added to the media that contained 0.75 g Na₂S and 0.6 g cysteine per liter. Once the media was cooled, the flasks were transferred into a Coy anaerobic glove bag where the 50 mL of media was dispensed into 160 mL serum bottles filled and autoclaved (121° C., 20 min). After autoclaving, the headspace gas was exchanged for 80% N₂/20% CO₂ mixture. The bottles then were inoculated with a 10% inoculum from previous stock cultures and amended with 30 mM glycerol. Sterilized 1,3-propanediol was added to the serum bottles to achieve 10, 30, 60, 120, 380, and 750 mM. Growth was examined by turbidity readings taken at 600 nM. It was determined that *H. hydrogeniformans*

11

was capable of growth when 0, 10, 30, 60, 120, and 380 mM 1,3-propanediol concentrations were present. The data indicates that *H. hydrogeniformans* can tolerate at least 0.38M 1,3-propanediol in addition to 7% (w/v) and pH 11.

TABLE 5

	1,3-propanediol Concentration (mM)						
	0	10	30	60	120	380	750
Growth	X	X	X	X	X	X	—

Example 6

Tolerance of *H. hydrogeniformans* to Crude Glycerol

The tolerance of *H. hydrogeniformans* to crude glycerol was examined. Anaerobic cultures were prepared in 160 mL serum bottles. The medium was prepared by boiling to degas under a N₂ blanket. As the medium cooled, reductant stock mix was added to the media that contained 0.75 g Na₂S and 0.6 g cysteine per liter. Once the media was cooled, the flasks were transferred into a Coy anaerobic glove bag where the 50 mL of media was dispensed into 160 mL serum bottles filled

12

and autoclaved (121° C., 20 min). After autoclaving, the headspace gas was exchanged for 80% N₂/20% CO₂ mixture. The bottles then were inoculated with a 10% inoculum from previous stock cultures. Crude glycerol, obtained from a small biodiesel producer, was added at 0.1% and 0.5% concentrations. No purification steps were applied to the crude glycerol. Growth, after one week, was examined by turbidity readings taken at 600 nM. It was determined that *H. hydrogeniformans* was capable of growth when exposed to crude, unpurified glycerol. *H. hydrogeniformans* can grow in at least 0.5% crude glycerol. Slow growth in 0.1% crude, most likely due to low glycerol concentration. Even slower growth in 0.5% crude glycerol.

DISCUSSION

The work has identified 1,3-propanediol production capability at pH 11 and 7% (w/v) NaCl of *H. hydrogeniformans*. The microorganism is capable of growth in 1M glycerol (along with 7% NaCl and a pH of 11) and 380 mM of 1,3-propanediol. In the absence of B₁₂, the conversion rate is 31%. With B₁₂ supplementation (>25 µg/L B₁₂), the conversion rate is approximately 60% conversion. The microorganism is also capable of growing in at least 0.5% crude glycerol, without treatment.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 7

<210> SEQ ID NO 1

<211> LENGTH: 1191

<212> TYPE: DNA

<213> ORGANISM: Halanaerobium hydrogeniformans

<220> FEATURE:

<221> NAME/KEY: gene

<222> LOCATION: (1)..(1191)

<223> OTHER INFORMATION: 16S rRNA

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (621)..(621)

<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 1

```

ggaagccttc gggcggaaga cgagactaga tagtggcggg cgggtgagta acacgtggat      60
aacctgtcct caagtctggg ataacctggc gaaagtcggg ctaatcccgg gtaagctgag      120
agtgtggcat cacacaatca gaaaagtgcc tattagcatt gtttgaggag gggtcggcgg      180
tagattagct agctgggtgag gtaatggctc accagggcga caatctatag ctggtctgag      240
aggacgatca gtcacactgg aactgagaca cggtcagac tctacggga ggcagcagtg      300
gggaatcttc cacaatgggc gaaagcctga tggagcaacg ccgcgtgagt gaagaaggtc      360
ttaggattgt aaagctctgt ccttagggaa gaaccgtggg tatagaaaat gataccacc      420
tgacggtaac tttggaggaa gcaactggct actacgtgcc agcagccggg gtaatacgtg      480
gagtgcgaag gttgtccgga attattgggc gtaaagggtg cgcaggcggg taatcaagtc      540
aagcgtgaaa ggtgtcggct taaccgacag actgcgtttg aaactgggta tcttgagtg      600
aacagaggag agtggaattc ntagtgtagt ggtgaaatac gtagatatta ggaagaacac      660
cagtggcgaa ggcgactctc tgggtaaca ctgacgctga ggtacgaaag ctgggggagc      720
gaacgggatt agataccccc gtagtcccag ccgtaaacga tggatactag gtgttgagg      780
ttcgaatcct tcagtgccgg agttaacgca ttaagtatcc cgctgggga ttacgatcgc      840
aagattgaaa ctcaaaggaa ttgacggggg ccgcacaag cggtggagca tgtggtttaa      900

```

-continued

```

ttcgaagcaa cgcaagaac cttaccgaga attgacatcc cgtgactacc tgtgaaagca 960
gggtttggca tttatgtcac acggagacag gtggtgcatg gctgtcgta gctcgtgctg 1020
tgagatggtt ggtaagtcc cgcaacgagc gcaaccctg ttcttagttg ccagcgagta 1080
atgtcgggga ctctaagaag actgcggtg aaagtcggag gaaggtgggg atgacgtcaa 1140
gtcctcatgc cctttatc tcgggctaca cacgtgctac aatggttgg a 1191

```

```

<210> SEQ ID NO 2
<211> LENGTH: 1671
<212> TYPE: DNA
<213> ORGANISM: Halanaerobium hydrogeniformans
<220> FEATURE:
<221> NAME/KEY: gene
<222> LOCATION: (1)..(1671)
<223> OTHER INFORMATION: glycerol dehydratase (+)strand

```

```

<400> SEQUENCE: 2

```

```

gtgaaaaggt ccaagcgatt tttagaactt gaaaaaaggc cgattagtaa tgatggattt 60
atcaatgaat ggccagaagc tggctctgtg gctatggaag gacccaatga tccaaaacca 120
agcgttagag ttgaaaatgg taaaatagta gagttagatg gtaaaagaag agaagaattt 180
gacatgttag actcctttat tgcgatcat actcttgatc tagatatagt tgaagatggt 240
atggcccagg attcaaaaga actggctcat aaaatagtag acataaatgt ttgtagagat 300
gaggtaaga gatttggtt aggtatgacc cctgctaaag ttgttgaagt ggttggctcat 360
atgaatggtt ttgaaatgat gatggcggtt cagaagatga gagctagaaa aactccttct 420
aaccagtgcc atgttacaaa tgtaaaagac caccagcat tattagctgc agatgcagca 480
gaagcagctc tgcgtggtt tgatgaaatg gaaacaacag taggtatagt tagatatgcg 540
ccttctaacy ctatttcaat catggtaggt tcacaaactg gccgtggtgg agttttaacc 600
cagtgtgctg ttgaggaagc tatggaatta gaaatgggta tgcgtggatt tacagcctat 660
gctgaaacag tttcagtata tggtagagag caggtatttg ttgatggtga tgatacacca 720
tggctcaaga gtttcttagc atctgcttat gcacccctg gattaaaaat gagatatact 780
tctggtaccg gttctgaagc agaaaaggga tttgctgatg gtaaatctat gctttatctt 840
gaagctcgcg gcttatatat gaccaaaggc gcaggagttc aggtatata gaatggttca 900
atcagttgta ttgggtgtgc cggggcagtt ccatccggag tttagagcaat tctggctgaa 960
aacttaattg ccatgttact tgacttagag tgtgcatctg gtaatgatca gacatttact 1020
cactcaagta caagaagaac tgctagaatg ttaatgcagt ttttgcctgg tactgacttt 1080
gtattctcag gttacagtgc tgtaccaaac tatgataaca tgtttgagg ttcaactcat 1140
gatgttgatg actatgatga ctatctaaca ttacagcgtg acttaaaagt taatggtgga 1200
ttagtacctg tagatgaaga agatgttatt aaagttagaa acaaagggtg tagagcttta 1260
caggcagtat ttaaaagaat cggtatgcct gatattacag atgaagaagt agaggctgca 1320
acatagccc acggtagtaa agatagcct gaccgtgatg taagagaaga tttcagaggt 1380
atagaagaaa tgctcaataa aggaaccaca ggtgttcaaa tcgtacaggc acttgctaaa 1440
catggatttg aagatgtagc tgaaaacttc tttaatctct taaaacagag aattgctgga 1500
gattatctcc atacatcagc gatctttgat gaaaacttcc atgcaataag tgctgttaat 1560
gataaaaatg attatgcagg tcctggaaca ggatagagag tttctgatga actctgggaa 1620
aaacttaaaa atggttagatt cgctaaagat attgacgaaa taggagaata a 1671

```

-continued

<210> SEQ ID NO 3
 <211> LENGTH: 556
 <212> TYPE: PRT
 <213> ORGANISM: Halanaerobium hydrogenoformans
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(556)
 <223> OTHER INFORMATION: glycerol dehydratase

<400> SEQUENCE: 3

Met Lys Arg Ser Lys Arg Phe Leu Glu Leu Glu Lys Arg Pro Ile Ser
 1 5 10 15
 Asn Asp Gly Phe Ile Asn Glu Trp Pro Glu Ala Gly Leu Val Ala Met
 20 25 30
 Glu Gly Pro Asn Asp Pro Lys Pro Ser Val Arg Val Glu Asn Gly Lys
 35 40 45
 Ile Val Glu Leu Asp Gly Lys Arg Arg Glu Glu Phe Asp Met Leu Asp
 50 55 60
 Ser Phe Ile Ala Asp His Thr Leu Asp Leu Asp Ile Val Glu Asp Val
 65 70 75 80
 Met Ala Gln Asp Ser Lys Glu Leu Ala His Lys Ile Val Asp Ile Asn
 85 90 95
 Val Cys Arg Asp Glu Val Lys Arg Phe Gly Leu Gly Met Thr Pro Ala
 100 105 110
 Lys Val Val Glu Val Val Gly His Met Asn Val Val Glu Met Met Met
 115 120 125
 Ala Val Gln Lys Met Arg Ala Arg Lys Thr Pro Ser Asn Gln Cys His
 130 135 140
 Val Thr Asn Val Lys Asp His Pro Ala Leu Leu Ala Ala Asp Ala Ala
 145 150 155 160
 Glu Ala Ala Leu Arg Gly Phe Asp Glu Met Glu Thr Thr Val Gly Ile
 165 170 175
 Val Arg Tyr Ala Pro Ser Asn Ala Ile Ser Ile Met Val Gly Ser Gln
 180 185 190
 Thr Gly Arg Gly Gly Val Leu Thr Gln Cys Ala Val Glu Glu Ala Met
 195 200 205
 Glu Leu Glu Met Gly Met Arg Gly Phe Thr Ala Tyr Ala Glu Thr Val
 210 215 220
 Ser Val Tyr Gly Thr Glu Gln Val Phe Val Asp Gly Asp Asp Thr Pro
 225 230 235 240
 Trp Ser Lys Ser Phe Leu Ala Ser Ala Tyr Ala Ser Arg Gly Leu Lys
 245 250 255
 Met Arg Tyr Thr Ser Gly Thr Gly Ser Glu Ala Glu Met Gly Phe Ala
 260 265 270
 Asp Gly Lys Ser Met Leu Tyr Leu Glu Ala Arg Cys Leu Tyr Met Thr
 275 280 285
 Lys Gly Ala Gly Val Gln Gly Ile Gln Asn Gly Ser Ile Ser Cys Ile
 290 295 300
 Gly Val Pro Gly Ala Val Pro Ser Gly Val Arg Ala Ile Leu Ala Glu
 305 310 315 320
 Asn Leu Ile Ala Met Leu Leu Asp Leu Glu Cys Ala Ser Gly Asn Asp
 325 330 335
 Gln Thr Phe Thr His Ser Ser Thr Arg Arg Thr Ala Arg Met Leu Met
 340 345 350
 Gln Phe Leu Pro Gly Thr Asp Phe Val Phe Ser Gly Tyr Ser Ala Val

-continued

355				360				365							
Pro	Asn	Tyr	Asp	Asn	Met	Phe	Ala	Gly	Ser	Thr	His	Asp	Val	Asp	Asp
	370					375					380				
Tyr	Asp	Asp	Tyr	Leu	Thr	Leu	Gln	Arg	Asp	Leu	Lys	Val	Asn	Gly	Gly
385					390					395					400
Leu	Val	Pro	Val	Asp	Glu	Glu	Asp	Val	Ile	Lys	Val	Arg	Asn	Lys	Gly
				405					410					415	
Val	Arg	Ala	Leu	Gln	Ala	Val	Phe	Lys	Glu	Ile	Gly	Met	Pro	Asp	Ile
		420							425				430		
Thr	Asp	Glu	Glu	Val	Glu	Ala	Ala	Thr	Tyr	Ala	His	Gly	Ser	Lys	Asp
		435					440					445			
Met	Pro	Asp	Arg	Asp	Val	Arg	Glu	Asp	Phe	Arg	Gly	Ile	Glu	Glu	Met
	450					455					460				
Leu	Asn	Lys	Gly	Thr	Thr	Gly	Val	Gln	Ile	Val	Gln	Ala	Leu	Ala	Lys
465					470					475					480
His	Gly	Phe	Glu	Asp	Val	Ala	Glu	Asn	Phe	Phe	Asn	Leu	Leu	Lys	Gln
				485					490					495	
Arg	Ile	Ala	Gly	Asp	Tyr	Leu	His	Thr	Ser	Ala	Ile	Phe	Asp	Glu	Asn
		500							505				510		
Phe	His	Ala	Ile	Ser	Ala	Val	Asn	Asp	Lys	Asn	Asp	Tyr	Ala	Gly	Pro
		515					520					525			
Gly	Thr	Gly	Tyr	Arg	Val	Ser	Asp	Glu	Leu	Trp	Glu	Lys	Leu	Lys	Asn
	530					535					540				
Val	Arg	Phe	Ala	Lys	Asp	Ile	Asp	Glu	Ile	Gly	Glu				
545					550					555					

<210> SEQ ID NO 4

<211> LENGTH: 1167

<212> TYPE: DNA

<213> ORGANISM: Halanaerobium hydrogenoformans

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(1167)

<223> OTHER INFORMATION: iron-containing alcohol dehydrogenase (+)strand

<400> SEQUENCE: 4

```

atgtatgatt atatgttacc aacagtaaat tttatgggag caggttctgt taaagtagta    60
ggagaaaagag caaaaattct ggggtcaaaa aaagttcttt tagtcactga tgacttttta    120
agtaatctag atggtggacc ctttgagacc gtagttaat atattgaaga agcgggttta    180
gcatatgctg tttatgatgg tgtaaaggct aatcccagag atactaatgt ttatgaaggc    240
ttagaaattt acgaaaaatga aaattgtgat atgataatta cagtaggtgg aggaaagtct    300
catgattgtg gtaaggcaat cggggttgct gcaactcatg atggagattt atataaggat    360
tatgcccggag tagaaaaatt agaaaatgaa acacctgcta tgatctgtgt taatacaact    420
gctggaacag ccagtgaggt taccagacat gcagtaataa ctgatacttc tcaaacacct    480
catgtaaagt ttgtaattgt cagctggcgt aatgtgccag atgtgtctat taatgatccg    540
gaactaatgg tagctaaacc tgctgcatta actgcagcta caggaatgga tgctttaact    600
catgctttag aaacctttgt ttcaacaggt gctaattcat taacagatgc agctgccaaa    660
gaagctatgg aactaatagc taagtactta agaagggcag tttataatgg agaagatatt    720
gaagccagag aagagatggc taatgcttca gttttagccg gttttgcctt caacaatgg    780
ggtttaggtt atgttcatgc tatggcccat cagttgggtg gattttatga tatgccacat    840
gggatagcca atgcaatcct cctgccttat gttgaaaaat ttaatttagg agctaaaata    900

```

-continued

```

gataaatttg ccaaggctgc agaaatattt ggagttccta cagctgggct ttctaaaaga    960
gaagctgctg aaaaatcttt agatgcaatt gtacagctgg ctgaagatat cggaatcccc    1020
acttctttaa gtgaatcaga atatgatgtc aaagaagaag attttgagga gatggcaaga    1080
ttagctttag aagatggtaa tgctttaagc aatcctagaa aagcaactca agcagaaatt    1140
gccggaatct ttaaagcagc ttattaa                                     1167

```

```

<210> SEQ ID NO 5
<211> LENGTH: 388
<212> TYPE: PRT
<213> ORGANISM: Halanaerobium hydrogenoformans
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(388)
<223> OTHER INFORMATION: iron-containing alcohol dehydrogenase

```

```

<400> SEQUENCE: 5

```

```

Met Tyr Asp Tyr Met Leu Pro Thr Val Asn Phe Met Gly Ala Gly Ser
1           5           10           15
Val Lys Val Val Gly Glu Arg Ala Lys Ile Leu Gly Ala Lys Lys Val
20          25          30
Leu Leu Val Thr Asp Asp Phe Leu Ser Asn Leu Asp Gly Gly Pro Phe
35          40          45
Glu Thr Val Val Lys Tyr Ile Glu Glu Ala Gly Leu Ala Tyr Ala Val
50          55          60
Tyr Asp Gly Val Lys Ala Asn Pro Arg Asp Thr Asn Val Tyr Glu Gly
65          70          75          80
Leu Glu Ile Tyr Glu Asn Glu Asn Cys Asp Met Ile Ile Thr Val Gly
85          90          95
Gly Gly Ser Pro His Asp Cys Gly Lys Ala Ile Gly Val Ala Ala Thr
100         105         110
His Asp Gly Asp Leu Tyr Lys Asp Tyr Ala Gly Val Glu Lys Leu Glu
115         120         125
Asn Glu Thr Pro Ala Met Ile Cys Val Asn Thr Thr Ala Gly Thr Ala
130         135         140
Ser Glu Val Thr Arg His Ala Val Ile Thr Asp Thr Ser Gln Thr Pro
145         150         155         160
His Val Lys Phe Val Ile Val Ser Trp Arg Asn Val Pro Asp Val Ser
165         170         175
Ile Asn Asp Pro Glu Leu Met Val Ala Lys Pro Ala Ala Leu Thr Ala
180         185         190
Ala Thr Gly Met Asp Ala Leu Thr His Ala Leu Glu Thr Phe Val Ser
195         200         205
Thr Gly Ala Asn Ser Leu Thr Asp Ala Ala Ala Lys Glu Ala Met Glu
210         215         220
Leu Ile Ala Lys Tyr Leu Arg Arg Ala Val Tyr Asn Gly Glu Asp Ile
225         230         235         240
Glu Ala Arg Glu Glu Met Ala Asn Ala Ser Val Leu Ala Gly Phe Ala
245         250         255
Phe Asn Asn Gly Gly Leu Gly Tyr Val His Ala Met Ala His Gln Leu
260         265         270
Gly Gly Phe Tyr Asp Met Pro His Gly Ile Ala Asn Ala Ile Leu Leu
275         280         285
Pro Tyr Val Glu Lys Phe Asn Leu Gly Ala Lys Ile Asp Lys Phe Ala
290         295         300

```

-continued

Lys Val Ala Glu Ile Phe Gly Val Pro Thr Ala Gly Leu Ser Lys Arg
 305 310 315 320
 Glu Ala Ala Glu Lys Ser Leu Asp Ala Ile Val Gln Leu Ala Glu Asp
 325 330 335
 Ile Gly Ile Pro Thr Ser Leu Ser Glu Ser Glu Tyr Asp Val Lys Glu
 340 345 350
 Glu Asp Phe Glu Glu Met Ala Arg Leu Ala Leu Glu Asp Gly Asn Ala
 355 360 365
 Leu Ser Asn Pro Arg Lys Ala Thr Gln Ala Glu Ile Ala Gly Ile Phe
 370 375 380
 Lys Ala Ala Tyr
 385

<210> SEQ ID NO 6
 <211> LENGTH: 1287
 <212> TYPE: DNA
 <213> ORGANISM: Halanaerobium hydrogenoformans
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(1287)
 <223> OTHER INFORMATION: iron-containing alcohol dehydrogenase (+)strand

<400> SEQUENCE: 6
 atgtccgatt attatgatta tatgctgcca actgtaaatt ttatgggacc tggctgtgta 60
 gaggttggtg gagaaaggtg caaaatttta ggtgcaaaaa aagttttaat agtgactgac 120
 agctttttaa gaaatatgga ggggtggacct gtagatcagg ttgttaaata tttaaagaaa 180
 gctaatttga attatgcatt ttatgatgaa gttgaaccta atcctaaaga tgtaaatggt 240
 tatgctgggc ttaagattta cgaaagagaa aattgtgaca tgattgtaac tattggtggt 300
 ggaagtgctc atgattgtgg aaaagcaatt ggagttgcag ctaccatga tggtgattta 360
 tacaagatt atgctgggtat tgaaaaacta gaaaatgaaa ctctcccat ggtctgtgta 420
 aatacaaccg ctggaactgc tagtgaggtt accaggcaca cagttattac tgacacttct 480
 cagactccaa acgttaaatt tgttatagta agttggagga atacaccgga tgtctctatc 540
 aatgatccgg aacttatggt tggtaaacca cctggattaa ctgctgcaac cggtatggat 600
 gctctgacct atgcagtaga aacatatgtc tcaactaatg caaatgcttt aactgatgca 660
 gcagctatta aatcaatcga attggtcgca aataatttaa gaaaagtcgt taaagatggt 720
 caggatatta aagcacgtga aaatatggct aatgcacccg tattatctgg tttgccttc 780
 aacaatggtg gcctggggtta tgttcatgct atggctcatc aactaggtgg tttttatgat 840
 atgccacacg gtatagctaa tgccatttta ctgccttatg tagaaaagtt taatcttggc 900
 acagatgtag agcgtttctc aatatattact gaaatatttg gcaaagaaca aagtaaaata 960
 tctaataatc cagaagctca agaatcaatt aaagctatta aagatgaaat cgataagcta 1020
 aaaagattta aaaaaatcgc tgaagttttt ggtgttgata caagtaatat gtcaacaaga 1080
 gaagcggctg aagcttcttt agacgccatt aaagaactag ctcgagatat tggaaattcca 1140
 agctctctga gcgaatctaa atttgatggt aaaagagacg attttgaaga aatggcaaaa 1200
 ttagctttag aggatggtaa tgctggaact aaccctagaa aaggtagtgt agaagatatt 1260
 gtaagaatat ttgaagatgc ctttttaa 1287

<210> SEQ ID NO 7
 <211> LENGTH: 428
 <212> TYPE: PRT

-continued

```

<213> ORGANISM: Halanaerobium hydrogenoformans
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(428)
<223> OTHER INFORMATION: iron-containing alcohol dehydrogenase

<400> SEQUENCE: 7

Met Ser Asp Tyr Tyr Asp Tyr Met Leu Pro Thr Val Asn Phe Met Gly
1          5          10          15

Pro Gly Cys Val Glu Val Val Gly Glu Arg Cys Lys Ile Leu Gly Ala
20          25          30

Lys Lys Val Leu Ile Val Thr Asp Ser Phe Leu Arg Asn Met Glu Gly
35          40          45

Gly Pro Val Asp Gln Val Val Lys Tyr Leu Lys Lys Ala Asn Leu Asn
50          55          60

Tyr Ala Phe Tyr Asp Glu Val Glu Pro Asn Pro Lys Asp Val Asn Val
65          70          75          80

Tyr Ala Gly Leu Lys Ile Tyr Glu Arg Glu Asn Cys Asp Met Ile Val
85          90          95

Thr Ile Gly Gly Gly Ser Ala His Asp Cys Gly Lys Ala Ile Gly Val
100         105         110

Ala Ala Thr His Asp Gly Asp Leu Tyr Lys Asp Tyr Ala Gly Ile Glu
115         120         125

Lys Leu Glu Asn Glu Thr Pro Pro Met Val Cys Val Asn Thr Thr Ala
130         135         140

Gly Thr Ala Ser Glu Val Thr Arg His Thr Val Ile Thr Asp Thr Ser
145         150         155         160

Gln Thr Pro Asn Val Lys Phe Val Ile Val Ser Trp Arg Asn Thr Pro
165         170         175

Asp Val Ser Ile Asn Asp Pro Glu Leu Met Val Gly Lys Pro Pro Gly
180         185         190

Leu Thr Ala Ala Thr Gly Met Asp Ala Leu Thr His Ala Val Glu Thr
195         200         205

Tyr Val Ser Thr Asn Ala Asn Ala Leu Thr Asp Ala Ala Ala Ile Lys
210         215         220

Ser Ile Glu Leu Val Ala Asn Asn Leu Arg Lys Val Val Lys Asp Gly
225         230         235         240

Gln Asp Ile Lys Ala Arg Glu Asn Met Ala Asn Ala Ser Val Leu Ser
245         250         255

Gly Phe Ala Phe Asn Asn Gly Gly Leu Gly Tyr Val His Ala Met Ala
260         265         270

His Gln Leu Gly Gly Phe Tyr Asp Met Pro His Gly Ile Ala Asn Ala
275         280         285

Ile Leu Leu Pro Tyr Val Glu Lys Phe Asn Leu Gly Thr Asp Val Glu
290         295         300

Arg Phe Ser Asn Ile Thr Glu Ile Phe Gly Lys Glu Gln Ser Lys Ile
305         310         315         320

Ser Asn Asn Pro Glu Ala Gln Glu Ser Ile Lys Ala Ile Lys Asp Glu
325         330         335

Ile Asp Lys Leu Lys Arg Phe Lys Lys Ile Ala Glu Val Phe Gly Val
340         345         350

Asp Thr Ser Asn Met Ser Thr Arg Glu Ala Ala Glu Ala Ser Leu Asp
355         360         365

Ala Ile Lys Glu Leu Ala Arg Asp Ile Gly Ile Pro Ser Ser Leu Ser
370         375         380

```

-continued

Glu Ser Lys Phe Asp Val Lys Arg Asp Asp Phe Glu Glu Met Ala Lys
 385 390 395 400

Leu Ala Leu Glu Asp Gly Asn Ala Gly Thr Asn Pro Arg Lys Gly Ser
 405 410 415

Val Glu Asp Ile Val Arg Ile Phe Glu Asp Ala Phe
 420 425

What is claimed:

1. A method of producing 1,3-propanediol, said method comprising fermenting *Halanaerobium hydrogeniformans* with a source of glycerol in a fermentation reactor, whereby 1,3-propanediol is produced, said reactor comprising an inlet for said source of glycerol, a fermentation chamber containing said *H. hydrogeniformans* in culture media, and an outlet for removing said 1,3-propanediol.

2. The method of claim 1, further comprising recovering said 1,3-propanediol.

3. The method of claim 1, wherein said source of glycerol is chemical waste from biodiesel production comprising glycerol.

4. The method of claim 3, wherein said chemical waste comprises crude glycerol.

5. The method of claim 3, wherein said fermenting is carried out without neutralization of said chemical waste pH prior to said fermenting.

6. The method of claim 3, wherein said fermenting is carried out without diluting the salinity of said chemical waste prior to said fermenting.

7. The method of claim 3, said chemical waste further comprising methanol, wherein said fermenting is carried out without removing said methanol from said chemical waste prior to said fermenting.

8. The method of claim 1, wherein said fermenting comprises culturing said *H. hydrogeniformans* with said source of glycerol in a culture medium to yield a fermentation culture.

9. The method of claim 8, wherein said culture medium comprises Vitamin B₁₂.

10. The method of claim 8, wherein said fermentation culture has a salt content of greater than or equal to 5% w/v.

11. The method of claim 1, wherein said fermenting is carried out at a pH of greater than or equal to 10.

12. The method of claim 1, said fermenting is carried out under substantially anaerobic conditions.

13. The method of claim 1, wherein said *H. hydrogeniformans* is the organism deposited as ATCC designation No. PTA-10410.

14. The method of claim 1, wherein said *H. hydrogeniformans* comprises an endogenous gene coding for a glycerol dehydratase or an enzyme having glycerol dehydratase activity.

15. The method of claim 14, wherein said gene comprises a DNA sequence comprising SEQ ID NO:2 or a sequence having at least 98% sequence identity with SEQ ID NO: 2.

16. The method of claim 14, wherein said glycerol dehydratase comprises SEQ ID NO:3, or a sequence having at least 98% sequence identity with SEQ ID NO: 3.

17. The method of claim 1, wherein said *H. hydrogeniformans* comprises an endogenous gene coding for an iron-containing alcohol dehydrogenase or an enzyme having alcohol dehydrogenase activity.

18. The method of claim 17, wherein said gene comprises a DNA sequence comprising SEQ ID NO:4 or 6, or a sequence having at least 98% sequence identity with SEQ ID NO: 4 or 6.

19. The method of claim 17, wherein said iron-containing alcohol dehydrogenase comprises SEQ ID NO:5 or 7, or a sequence having at least 98% sequence identity with SEQ ID NO: 5 or 7.

20. A method of producing 1,3-propanediol, said method comprising fermenting *Halanaerobium hydrogeniformans* with a source of glycerol, whereby 1,3-propanediol is produced, wherein said source of glycerol is chemical waste from biodiesel production comprising glycerol and methanol, wherein said fermenting is carried out without removing said methanol from said chemical waste prior to said fermenting.

* * * * *