

Anti-hyperlipidaemic effects of fresh and cured *Bhallataka Kshaudra* (*Semecarpus anacardium* L.) in animals

Mohan Krishna Dwivedi*, Mukeshkumar Nariya¹, R Galib and Pradeep Kumar Prajapati

Department of Rasa Shastra and Bhaishajya Kalpana including Drug Research, ¹Pharmacology Laboratory, Institute for Post Graduate Teaching and Research in Ayurveda, Gujarat Ayurved University, Jamnagar

Received 31 December 2016; Revised 29 May 2018

Bhallataka Phala (fruit of *Semecarpus anacardium* L.), is a well-known anti-hyperlipidaemic drug. *Bhallataka* is described in all Ayurvedic classics as *Rasayana* and advocated for various therapeutic purposes such as *Kushtha* (skin diseases), *Arsha* (piles), *Krimi* (worm infestations), *Prameha* (urine disorders), *Medodosha* (lipid disorders), etc. Though its anti-hyperlipidaemic activity has been studied; actual differentiation in their therapeutic efficacy due to storage period has yet not been attempted on experimental animals. This prompted to initiate a comparative anti-hyperlipidemic activity of fixed oil (expressed oil) i.e. non-volatile in nature collected from fresh *Bhallataka* and four months old (cured) *Bhallataka* samples against cholesterol diet induced hyperlipidaemia in rats. Hyperlipidemia was induced by oral administration of cholesterol (20 % suspension in coconut oil, 5 mL/kg) in morning hours and hydrogenated vegetable oil (5 mL/kg) in evening. The effect of drugs was assessed on body weight, serum biochemical and histological parameters. Both drugs produced significant attenuation of the relative weight of liver in cholesterol-fed animals. Cured sample of *Bhallataka* provided better effect in lowering serum cholesterol (21.98 %), triglyceride (60.23 %), VLDL (56.82 %); while fresh sample of *Bhallataka* also found to be effective in lowering serum cholesterol (20.69 %), serum triglyceride (45.59 %), VLDL (46.59 %) in comparison to control group. From the result of the present study, it is concluded that the cured sample of *Bhallataka* has pronounced anti-hyperlipidemic effect than the fresh sample in experimentally-induced hyperlipidemia in rats.

Keywords: Anti-hyperlipidaemia, *Bhallataka*, curing, Polyphenolic compounds, *Semecarpus anacardium* L.

IPC code; Int. cl. (2015.01) - A61K 36/00, 36/22; A61P 3/06

Introduction

Hyperlipidaemia involves abnormally elevated levels of any one of total cholesterol, low-density lipoprotein, triglycerides or all lipids in the serum blood plasma. The metabolic consequences associated with changes in diet and lifestyle has increased the number of hyperlipidemic individuals who could benefit from lipid-lowering therapy¹. Lipoproteins play an essential role in the absorption of dietary cholesterol, long chain fatty acids and fat-soluble vitamins so help in transport of triglycerides, cholesterol and fat-soluble vitamins from the liver to peripheral tissues and vice versa. Multiple epidemiologic studies have demonstrated a strong relationship between serum cholesterol, elevated plasma triglyceride levels and coronary heart disease (CHD). Cardiovascular diseases are the number one cause of death globally². An estimated 7.3 million death were due to coronary heart disease and 6.2

million were due to stroke³. The traditional system of medicines is a better alternative with minimal side effects to combat these types of disorders.

Bhallataka (*Semecarpus anacardium* L.) belonging to family Anacardiaceae is a common tree found throughout the country especially in the Himalayas and hotter parts of Indian in dry and moist deciduous forests⁴. The fruit is useful in treating many diseases like *Kushtha* (skin diseases), *Arsha* (piles), *Krimi* (worm infestations), *Prameha* (urine disorders), *Medodosha* (lipid disorders) etc⁵ and is emphasized as best *Rasayana* (rejuvenator). Fruit of *S. anacardium* is well known for its anti-hyperlipidaemic property (*Medodoshanashaka*)⁶. *Kala-Prakarsha Samskara* (curing) is one of the important *Samskara* (processing and propagation) especially mentioned in *Ayurvedic* texts, which plays a major role in propagating the therapeutic efficacy and safety of medicines⁷. *Kala Prakarsha Samskara* especially emphasizes for *Bhallataka* i.e. storage of *Bhallataka* fruits in heap of barley for a period of four months in rainy season⁸. Preservation time alters the drug properties, mainly its

*Correspondent author
Email: drmkdi@gmail.com

potency. Several pharmacological and clinical trials are reported pertaining to *Bhallataka* fruit⁹⁻¹¹ but experimental studies for comparing the efficacy of fresh and cured samples have not yet been found. Therefore, the present study was planned with an aim to evaluate the comparative anti-hyperlipidaemic efficacy of fresh and cured *Bhallataka* fruits in albino rats.

Materials and Methods

Drug

Fresh *Bhallataka* fruits were collected from Jalna (19.83° N 75.88° E) Maharashtra, India during *Shucho-Shukre* (the month of April and May 2013)¹². The drug was identified and authenticated at Pharmacognosy Laboratory, Institute for Post Graduate Teaching and Research in Ayurveda, Gujarat Ayurved University, Jamnagar and voucher specimen was deposited (IPGT&RA/Phm/6165). *Prashasta* (acceptable) and *Aprashasta* (unacceptable) qualities of fruits were analyzed by immersing them in a vessel containing potable water in a glass jar. The fruits that settled down were collected carefully and used for further experimental study while the floated fruits were discarded as they are unacceptable for the study.

Curing of *Bhallataka*

Acceptable fruits were kept in an aluminium box of 50×35.5×55 cm (L×B×H) filled with a heap of barley to maintain constant temperature and humidity for four months. Before filling barley inside the box, thermocol sheets of 10 mm width were taken and cut off with proper dimension, now these sheets were wrapped with aluminium foil of thickness of 0.009-0.03 mm and width of 250 mm and pasted to the walls of the box. Another cabin was prepared from plain thermocol sheets inside the box with 4.5 cm distanced from previously fitted aluminium foil. *Yava* (Barley - *Hordeum vulgare*) fruits were filled in the created 4.5 cm space in between and lower down the previous sheets. Now, the batches of *Bhallataka* were arranged in the box with alternative barley layers. Finally, barley fruits layers were arranged on the upper part and box were closed for four months. After four months, fruits were collected and processed for further use.

Bhallataka oil extraction

Both samples of *Bhallataka* fruits were subjected to oil extraction by *Patala yantra* (Pit Method)¹³.

Bhallataka fruits were taken inside the vessel after cutting off its pseudocarp and then the mouth of the vessel was wrapped with mesh and another small vessel was fitted upon this large vessel. The whole arrangement was wrapped with mud smeared cloth and then put in a pit. Now, covered with cow-dung cakes and light was ignited. By this process, oil from fruits was extracted and collected into the small vessel.

Bhallataka Kshaudra formation process

To 20 g of *Madhu* (Honey), 160 g oil was added, then, 320 g *Ghrita* (cow ghee) was mixed and triturated for the homogeneous mixture as per Ayurvedic classic¹⁴. The same procedure was followed for the preparation of both test formulations i.e. fresh (FB) and cured (CB) samples of *Bhallataka Kshaudra*.

Animals

Wistar strain albino rats of either sex, weighing 200±20 g were used for the experiment. The animals were maintained under standard conditions of temperature (22±02 °C), humidity (50-60 %) and exposed to 12 hour light and dark cycles. All animals were exposed to the same environmental conditions and were maintained on a standard diet and water *ad libitum*. The experimental protocol was approved by the Institutional Animal Ethical Committee (M.D./IAEC/14/2013/18) as per guidelines of CPCSEA, India.

Dose

The human dose of *Bhallataka Kshaudra* is considered as 500 mg/day¹⁵. Rat dose was calculated on basis of body surface area ratio by referring to the table of Paget and Barnes (1964)¹⁶. Based on that, the rat dose was fixed at 45 mg/kg. The test drug formulations were administered through the oral route by licking.

Experimental design

Protocol designed in the earlier study was followed in the present study with modifications as per experimental need¹⁷. Wistar albino rats of either sex were divided into five groups of six animals each. Group (I) was kept as normal control (NC) which received distilled water (10 mL/kg, po). Group (II) of animals kept as cholesterol control group, received distilled water (10 mL/kg, po) and hyperlipidaemic diet. Group (III) of animals were treated with atorvastatin (7.2 mg/kg, po) along with

hyperlipidaemic diet (SC). Group (IV) and (V) were kept as drug-treated groups which received fresh *Bhallataka Kshaudra* (FB) (45 mg/kg) and cured *Bhallataka Kshaudra* (CB) (45 mg/kg) orally along with hyperlipidemic diet.

Test formulations and vehicle were administered to respective groups in the morning hours daily for 21 days. The hyperlipidaemic diet was concomitantly administered to Group (II) to (V) to induce hyperlipidemia. The hyperlipidemic diet included cholesterol suspension in coconut oil and hydrogenated vegetable oil. Cholesterol extra pure powder (SRL, Mumbai) made into 20 % suspension in coconut oil and administered orally (5 mL/kg) in the morning hours after 1 hour of drug administration while, hydrogenated vegetable oil (5 mL/kg) (Adani Wilmar Ltd., Gujarat) was administered in evening hours.

On the 22nd day, the overnight fasted rats were weighed again and blood was collected from retro-orbital puncture under light ether anaesthesia using capillary tubes, serum was separated and used for estimation of different serum biochemical parameters. An autoanalyzer (BS 200) was used for the estimations of serum total cholesterol¹⁸, triglyceride¹⁹, VLDL-cholesterol, Urea²⁰, creatinine²¹, blood sugar²², total protein²³, albumin, globulin²⁴, SGOT²⁵, SGPT²⁶ and alkaline phosphatase²⁷. The rats were sacrificed and important organs like the liver, kidney, heart and aorta were excised out. The extraneous tissues were cleaned off and weight of organs was noted down except for aorta and stored in 10 % formalin solution for histopathological study²⁸.

Statistical analysis

The data is expressed as mean±standard error of mean for six rats per experimental group. Student's t-test was used to compare the unpaired data to determine the significant difference between groups at $p < 0.05$.

Results

Cholesterol control, standard drug, and cured *Bhallataka* groups showed almost similar values of body weight gain during the experimental period while fresh *Bhallataka* treated group showed a decrease in body weight in comparison to initial values (Table 1). Cholesterol control group showed a non-significant increase in relative weight of liver while significant decrease ($p < 0.05$) in the relative weight of kidney in comparison to normal control group (Table 1). Test drugs produced a non-significant decrease in relative weight of heart, liver and kidney except for a significant decrease in relative weight of kidney in cured *Bhallataka* treated group.

Cholesterol-fed rats showed statistically significant increase in total cholesterol ($p < 0.05$), triglyceride ($p < 0.001$), VLDL-cholesterol ($p < 0.05$) and blood urea ($p < 0.001$) level in comparison to normal control rats. Standard drug and both the fresh and cured *Bhallataka Kshaudra* showed a statistically significant decrease in cholesterol ($p < 0.05$), triglyceride ($p < 0.001$) and VLDL-cholesterol ($p < 0.001$) level in comparison to cholesterol control group. Cured *Bhallataka Kshaudra* produced better effects on lipid profile and blood urea followed by fresh *Bhallataka Kshaudra* (Table 2).

Cured *Bhallataka Kshaudra* showed a statistically significant decrease while fresh *Bhallataka Kshaudra* produced a non-significant decrease in blood glucose level as compared to cholesterol control group whereas both the test drugs showed a significant decrease in total protein, albumin and creatinine ($p < 0.05$) level in comparison to cholesterol control group (Table 2). Cholesterol-fed rats showed a significant increase in SGOT, alkaline phosphatase ($p < 0.001$) and bilirubin ($p < 0.05$) level in comparison to the normal control group. Both the test drugs and atorvastatin-treated groups showed a statistically significant decrease in SGOT level, total and direct

Table 1 — Effect of test drugs on body weight and relative organ weight of albino rats during anti-hyperlipidaemic study

Groups	Body weight		Relative weight (g/100 g body weight)		
	Initial (g)	Final (g)	Liver	Heart	Kidney
NC	177.33±5.90	188.00±4.68	3.377±0.186	0.316±0.017	0.763±0.020
CC	212.00±5.42	222.00±12.22	3.583±0.0804	0.317±0.010	0.713±0.004 [#]
SC	179.67±1.75	189.33±3.17	3.492±0.207	0.328±0.005	0.694±0.029
FB	214.67±7.60	208.33±5.20	3.466±0.103	0.298±0.007	0.677±0.018
CB	206.67±12.56	224.00±12.04	3.481±0.130	0.310±0.010	0.650±0.004 ^{**}

Data: Mean±SEM, [#] $p < 0.05$ when compared to normal control group; ^{**} $p < 0.001$ when compared to cholesterol control group, NC— Normal Control, CC— Cholesterol control, SC— Standard control, FB— Fresh *Bhallataka Kshaudra*, CB— Cured *Bhallataka Kshaudra*

bilirubin levels while a non-significant decrease in alkaline phosphatase level in comparison to cholesterol control group (Table 3).

Histopathological studies have shown that, control group exhibits normal cytoarchitecture in aorta, heart, kidney and liver (Fig. 1a, 2a, 3a and 4a) respectively. Cholesterol administration in rats produced mild increase in thickness of adventitia of aorta (Fig. 1b, c), moderate fatty degenerative changes in heart (Fig. 2b, c), fatty degeneration, edematous changes and intense cell infiltration in kidney (Fig. 3b, c) and liver (Fig. 4b, c). Sections from fresh and cured *Bhallataka Kshaudra* treated groups showed normal cytoarchitecture of aorta (Fig. 1d, e), mild fatty changes in heart (Fig. 2d, e), mild fatty changes with edema in kidney (Fig. 3d, e) and mild fatty changes in liver (Fig. 4d, e) in comparison to cholesterol control group. Standard drug treated rats showed normal cytoarchitecture of aorta (Fig. 1f), moderate fatty degenerative changes and necrosis in heart (Fig. 2f), normal cytoarchitecture of kidney (Fig. 3f) and mild fatty changes in liver (Fig. 4f).

Discussion

Bhallataka is mentioned as *Medonashaka* in Ayurvedic text⁶. The previous study also suggests the

anti-hyperlipidaemic effect of an extract of *Bhallataka*²⁹ but, the difference in the fresh and cured fruits of *Bhallataka* prepared as per the classical method is not yet reported. The present study is carried out to evaluate the comparative anti-hyperlipidaemic efficacy of fresh and cured *Bhallataka* fruits in albino rats. Administration of hyperlipidaemia diet led to a marked increase in body weight, CB showed an increase in body weight but on the other hand, FB showed a decrease in body weight compared to initial values but the effect was non-significant. Cholesterol control group showed a non-significant increase in relative weight of liver and heart in comparison to control group whereas, drug-treated groups showed a non-significant decrease in relative weight of liver and heart in comparison to cholesterol control group. The weight of kidney significantly decreased in the cholesterol control group in comparison to control group. CB treated group showed significant decrease whereas, non-significant decrease in rest of the groups in comparison to cholesterol control group.

Administration of cholesterol and hyperlipidemic diet lead to significant elevation of serum total cholesterol, triglycerides, and VLDL-cholesterol

Table 2 — Effect of test drugs on lipid profile, urea content, blood sugar, total protein, albumin and globulin level in albino rats during anti-hyperlipidaemic study

Parameters	Groups				
	NC	CC	SC	FB	CB
Cholesterol (mg/ dL)	46.00±3.22	77.33±6.36 [#]	56.50±2.63*	61.33±2.93*	60.33±3.461*
Triglyceride (mg/ dL)	96.50±14.05	176.00±9.59 [#]	104.67±11.82**	94.00±5.42**	70.00±10.16**
VLDL (mg/ dL)	19.30±2.81	35.20±1.92 [#]	20.93±2.36**	18.80±1.08**	15.20±1.97**
Urea (mg/dL)	43.50±1.48	61.50±1.65 [#]	41.00±3.20**	41.67±1.26**	34.17±1.17**
Total protein (mg/ dL)	7.28 ±0.16	7.75±0.145	7.62±0.162	6.80±0.115**	6.85±0.118**
Albumin (mg/ dL)	3.88±0.083	3.75±0.062	4.05±0.085*	3.50±0.089*	3.32±0.087*
Globulin (mg/ dL)	3.40±0.13	4.05±0.178 [#]	3.60±0.255	3.30±0.0775*	3.533±0.193
Creatinine (mg/ dL)	0.55 ±0.02	0.55±0.02	0.53±0.03	0.42±0.05*	0.43±0.04*
Sugar (mg/dL)	103.83±5.78	96.00±8.32	96.67±5.03	76.17±4.06	72.83±2.96*

Data: Mean±SEM, [#] $p < 0.05$, [#] $p < 0.001$ when compared to normal control group; * $p < 0.05$, ** $p < 0.001$ when compared to cholesterol control group

Table 3 — Effect of test drugs on SGOT, SGPT alkaline phosphatase and creatinine level in albino rats during anti-hyperlipidaemic study

Groups	SGPT (IU/L)	SGOT (IU/L)	ALP (IU/L)	Bilirubin	
				(T) (mg/dL)	(mg/ dL)
NC	53.00±2.86	145.17±6.48	269.00±46.83	0.65±0.10	0.17±0.02
CC	52.33±3.48	219.83±15.76 [#]	535.00±87.81 [#]	1.07±0.08 [#]	0.28±0.05 [#]
SC	57.60±5.66	154.83±8.77*	426.00±46.81	0.62±0.05**	0.15±0.02*
FB	55.17±4.77	147.67±11.58*	409.83±59.34	0.48±0.03**	0.13±0.02*
CB	52.17±7.28	151.83±7.98*	494.17±79.35	0.52±0.06**	0.15±0.03*

Data: Mean±SEM, [#] $p < 0.05$, [#] $p < 0.001$ when compared to normal control group; * $p < 0.05$, ** $p < 0.001$ when compared to cholesterol control group




Fig. 1 — Photomicrographs of sections of aorta taken at 200x magnification. a) Normal cytoarchitecture (control group) b) and c) Increase in thickness of adventitia layer (CC) (d, e and f) Normal cytoarchitecture (FB, CB and SC respectively)




Fig. 2 — Photomicrographs of sections of heart taken at 200x magnification. a) Normal cytoarchitecture (control group) b) and c) Moderate fatty degenerative changes and necrosis (CC) (d and e) Mild fatty changes (FB and CB) f) Moderate fatty degenerative changes and necrosis (SC).

normal control rats. This establishes the efficacy of the experimental protocol to induce hyperlipidemic

condition. Lipid metabolism to great extent depends upon the formation and turnover of lipoproteins. Almost all lipids in the plasma are transported in the form of complexes with proteins; these proteins are termed as lipoproteins³⁰.

Both test drugs significantly, lowered the cholesterol level, triglyceride and VLDL cholesterol level in rats. The reversal effect was more pronounced in CB treated animals than FB treated animals in comparison to control group. The cholesterol-lowering effect of the drugs might be due to inhibition of dietary cholesterol absorption or esterification. Since two enzymes are involved in these two processes viz. pancreatic cholesterol esterase³¹ and intestinal acyl Co-A-cholesterol acyltransferase enzyme (ACAT)³², thus it could be suggested that the test drugs inhibit the activity of one or both of these enzymes. In both the test drugs ω -7 fatty acids were found which are responsible for lowering cholesterol level.

The decrease in cholesterol may indicate increased oxidation of mobilized fatty acids of inhibition or lipolysis. CB, FB and atorvastatin produced a highly significant decrease in the triglyceride levels and VLDL level compared to the cholesterol-fed group. Reduction in cholesterol, triglycerides and VLDL level may probably be due to the presence of steroids, flavonoids and polyphenolic compounds in *Bhallataka*³³. Flavonoids are a group of ubiquitously distributed plant polyphenols which exhibit a wide range of pharmacological effects. The inhibition of lipid peroxidation is due to the free radical scavenging property of flavonoids. They scavenge free radical by donating their hydrogen groups and prevent the initiation of a chain reaction. They also scavenge singlet O₂, terminating peroxides by their low redox potential³⁴. They also reduce lipid peroxidation by reducing the levels of malondialdehyde and conjugated dienes³⁵. Both the drugs have the presence of steroids, tannins, phenolic compounds, flavonoids and amines. The plant steroids reduce the absorption of cholesterol and thus increase faecal excretion of cholesterol³⁶.

SGOT is a mitochondrial enzyme released from heart, liver, skeletal muscle and kidney. SGPT is a cytosolic enzyme primarily present in the liver³⁷. Serum alkaline phosphatase is produced by many tissues, especially bone, liver, intestine and placenta and excreted in bile. Serum amino-transferase and alkaline phosphatase are elevated in most liver




Fig. 3 — Photomicrographs of sections of kidney taken at 100x magnification. a) Normal cytoarchitecture (control group) b) and c) Fatty degeneration, edematous changes and intense cell infiltration (CC) (d and e) Mild fatty changes with edema (FB and CB) f) Normal cytoarchitecture (SC).




Fig. 4 — Photomicrographs of sections of liver taken at 400x magnification. a) Normal cytoarchitecture (control group) b) and c) Fatty degeneration, edematous changes and cell infiltration (CC) (d, e and f) Mild fatty changes (FB, CB and SC respectively).

disorders. Highest elevations are found in conditions causing extensive hepatic necrosis such as severe

acute viral hepatitis, toxic hepatitis or prolonged circulatory collapse³⁸. Cholesterol-fed rats exhibited an insignificant increase in SGOT and a significant increase in alkaline phosphatase ($p < 0.001$) level suggests the liver, heart and kidney damage in rats. Test drug non-significantly decreased the level of alkaline phosphatase while a significant decrease in SGOT which suggests the protective role of the test drug in high fat diet-induced hyperlipidemia in rats.

A previous study showed that increased levels of total cholesterol, free cholesterol, phospholipids, triglycerides and free fatty acids and decreased levels of ester cholesterol in plasma, liver and kidney found in cancer-suffering animals were reverted back to near normal levels on treatment with the formulation of *Bhallataka*³⁹. Further, it reduces the tissue and serum hyperlipidemia by the inhibition of intestinal cholesterol absorption coupled with peripheral disposal thus possessing anti-atherosclerotic activity²⁹. *S. anacardium* nut. extract oil fraction at a dose of 1 mg/100 g body weight significantly reduced serum cholesterol levels and increased HDL cholesterol levels in the rat fed with atherogenic diet⁴⁰. In this context, a number of other plants have also been reported to have antihyperglycemic, antihyperlipidemic and insulin stimulatory effects^{41,42}.

Conclusion

From the present study, it is concluded that fresh and cured *Bhallataka Kshaudra* prepared as per classical method have anti-hyperlipidaemic activity in cholesterol-fed albino rats. Cured *Bhallataka Kshaudra* has a more pronounced effect than fresh *Bhallataka Kshaudra* in lowering cholesterol and related parameters which may due to the presence of more amount of ω -7 fatty acids, steroids, flavonoids and polyphenolic compounds in cured one as compared to fresh one.

References

- 1 Anthony S F, Dan L, Eugene B, Stephen L H and Larry J, Harrison's principles of internal medicine, 16th edn, edited by L Dennis (Mac Graw-Hill Publication, London) 2005, 2286.
- 2 Sainani G S, *API Textbook of Medicine*, 6th edn (Association of Physicians of India, Mumbai) 1999, 191.
- 3 Alwan A, *Global status report on non-communicable diseases 2010*, (World Health Organization, Geneva), 2011.
- 4 Talbot W A, *Forest flora of the Bombay presidency and Sind*, 1st edn (MS Periodical Experts, Delhi), 1976, 355-6.

- 5 Vriddha Vagbhata, *Astanga Samgraha (Indu-Sashilekhavyakhya)*, Uttar Tantra 49/51, 2nd edn, edited by S P Sharma (Chaukhamba Sanskrit series office, Varanasi), 2008, 917.
- 6 Sushruta and Chandrata, *Sushruta Samhita, Kavyatirtha*, 45/122, edited by J Trikamji, R Acharya (Chaukhamba Orientalia, Varanasi), 2009, 206.
- 7 Agnivesha, Charaka Samhita, in: *Ayurveda Dipika commentary, Vimana Sthana* 1-21/2, 2nd edn, edited by Yadavji T (Chaukhamba Prakashan, Varanasi), 2011, 235.
- 8 Agnivesha, Charaka Samhita, in: *Ayurveda Dipika commentary, Chikitsasthana* 1-2/13, 2nd edn, edited by Yadavji T (Chaukhamba Prakashan, Varanasi), 2011, 382.
- 9 Smit H F, Woerdenbag H J, Singh R H, Meulenbeld G J, Labadie R P, *et al.*, Ayurvedic herbal drugs with possible cytostatic activity, *J Ethnopharmacol*, 1995, **47**, 75-84.
- 10 Ramprasth V R, Shanthi P and Sachdanandam P, Evaluation of antioxidant effect of *Semecarpus anacardium* Linn. nut extract on the components of immune system in adjuvant arthritis, *Vascul Pharmacol*, 2005, **42**, 179-186.
- 11 Arul B, Kothai R and Christina A J, Hypoglycemic and antihyperglycemic effect of *Semecarpus anacardium* Linn. in normal and streptozotocin-induced rats, *Method Find Exp Clin Pharmacol*, 2004, **26**, 759.
- 12 Agnivesha, Charaka Samhita In: *Ayurveda Dipika commentary, Chikitsasthana* 1-2/13, 2nd edn, edited by Trikamji Y (Chaukhamba Prakashan, Varanasi), 2011, 382.
- 13 Yogi B, *Rasarnava*, 12/21-22, 4th edn, edited by Tripathi I (Chaukhamba Sanskrit Series office, Varanasi), 2001, 174.
- 14 Agnivesha, Charaka Samhita, in: *Ayurveda Dipika commentary, Chikitsasthana* 1-2/13, 2nd edn, edited by Yadavji T (Chaukhamba Prakashan, Varanasi), 2011, 382.
- 15 Sharma S, *Rasatarangini*, 24/482, 11th edn, edited by Kashinathshastri (Motilal Banarasidas, New Delhi), 2004, 737.
- 16 Paget G E and Barnes J M, Evaluation of drug activities, in *Pharmacometrics*, vol I, edited by D R Laurence & A L Bacharach (Academic Press, London), 1964, 50.
- 17 Nadkarni M, *Clinico experimental study of hyperlipidemia and its management by mustadighanavati*, M.D. Ayu dissertation, Institute of Post Graduate Teaching and Research in Ayurveda, Gujarat Ayurved University, Jamnagar, 2009.
- 18 Roeschlau P, Bernt E and Gruber W A, Enzymatic determination of total cholesterol in serum, *Clin Biochem*, 1974, **12**, 226-228.
- 19 Fossati P and Prencipe L, Serum triglycerides determined colorimetrically with an enzyme that produces hydrogen peroxide, *Clin Chem*, 1982, **28**, 2077-2080.
- 20 Talke H and Schubert G E, Enzymatic urea determination in the blood and serum in Warburg optical test, *Klin Wochenschr*, 1965, **42**, 174-5.
- 21 Slot C, Plasma creatinine determination: A new and specific Jaffe reaction method, *Scand J Clin Lab Invest*, 1965, **17**, 381-7.
- 22 Pennock C A, Murphy D, Sellers J and Longdon K J, A comparison auto analyzer method for the estimation of glucose in blood, *Clin Chim Acta*, 1973, **48**, 193-201.
- 23 Tietz N W, *Text book of Clinical Chemistry* (WB Saunders, Philadelphia, PA), 1986, 579.
- 24 Doumas B T, Arends R L and Pinto P C, in *Standard methods of Clinical Chemistry* (Academic Press, Chicago), 1972, 175-189.
- 25 Tietz N W, *Clinical guide to laboratory tests*, 3rd edn (WB Saunders, Philadelphia, PA), 1995, 76.
- 26 Burtis C A, Ashwood E R, and Burns D E, *Tietz textbook of Clinical Chemistry*, 3rd edn (WB Saunders, Philadelphia, PA), 1999, 652.
- 27 Wilkinson J H, Boutwell J H and Winsten S, Evaluation of a new system for kinetic measurement of serum alkaline phosphatase, *Clin Chem*, 1969, **15**, 487-95.
- 28 Raghramulu N, Nair M K and Kalyansundaram S, *A Manual of Laboratory technique* (National Institute of Nutrition, Jamai Osmania, Hyderabad), 1983, 92.
- 29 Sharma A, Mathur R and Dixit V P, Hypocholesterolemic activity of nut shell extract of *Semecarpus anacardium* (Bhilawa) in cholesterol fed rabbits, *Indian J Exp Biol*, 1995, **33**, 444-8.
- 30 Barter P, The role of HDL cholesterol in preventing atherosclerotic disease, *Eur Heart J Suppl*, 2005, **7**, 4-8.
- 31 Gallo L, Benett C, Myers S and Vanouny G, Cholesterol absorption in rat intestine- Role of cholesterol esterase and ACAT, *J Lipid Res*, 1984, **25**, 604-612.
- 32 Park Y B, Jeon S M, Byun S J, Kim H and Choi M S, Absorption of intestinal free cholesterol is lowered by supplementation of *Areca catechu* L. extracts in rats, *Life Sci*, 2002, **70**, 1849-1859.
- 33 Girija K, Anti-hyperlipidemic activity of methanol extracts of three plants of *Amaranthus* in triton-WR 1339 induced hyperlipidemic rats, *Asian Pac J Trop Biomed*, 2011, S65.
- 34 Klopman G and Dimayuga M L, Computer-automated structure evaluation of flavonoids and other structurally related compounds as glyoxalase I enzyme inhibitors, *Mol Pharmacol*, 1988, **34**, 218-222.
- 35 Sigers C P and Younes M, Effect of bioflavonoids on lipid peroxidation induced by GSH depletion in Proceedings of the International Bioflavonoid Symposium, *Kuwch FRG*, 1981, 403-409.

- 36 Delbas J M, Fernandez-Larrea J, Blay M, Ardevol A, Arola M J, *et al.*, Grape seed procyanidinins improve atherosclerotic risk index and induce liver CYP7A1 and STIP expression in healthy rats, *FASEB J*, 2005, **19**, 479-481.
- 37 Mohan H, *Text book of Pathology*, 5th edn (Jaypee Brothers Medical Publishers (P) Ltd, New Delhi), 2005, 611.
- 38 Gupta A, Kapoor N K and Nityanand S, Mechanism of hypolipidemic action of standardized extract, *Indian J Pharmacol*, 1982, **14**(1), 14.
- 39 Veena K, Shanthi P and Sachdanandam P, The biochemical alterations following administration of Kalpaamruthaa and *Semecarpus anacardium* in mammary carcinoma, *Chem Biol Interact*, 2006, **161**, 69-78.
- 40 Tripathi Y B and Pandey R S, *Semecarpus anacardium* L, nuts inhibit lipopoly- saccharide induced NO production in rat macro- phages along with its hypolipidemic property, *Indian J Exp Biol*, 2004, **42**, 432-436.
- 41 Yadav J P, Saini S, Kalia A N and Dangi A S, Hypoglycemic and hypolipidemic activity of ethanolic extract of *Salvadora oleoides* in normal and alloxan-induced diabetic rats, *Indian J Pharmacol*, 2008, **40**(1), 23-27.
- 42 Saini S and Yadav J P, Antidiabetic and antihyperlipidemic effects of ethanolic extract of *Salvadora persica* L. on alloxan-induced diabetic rats, *Der Pharmacia Sinica*, 2013, **4**(3), 178-182.