

Waterlogging tolerance in black gram [*Vigna mungo* (L.) Hepper] is associated with chlorophyll content and membrane integrity

Ruchi Bansal*, Shivani Sharma, Kuldeep Tripathi, Gayacharan & Ashok Kumar

ICAR-National Bureau of Plant Genetic Resources, Pusa Campus, New Delhi -110 012, Delhi, India

Received 28 November 2018; revised 05 December 2018

Black gram (*Vigna mungo* L.) is waterlogging sensitive legume crop. We studied the effect of waterlogging stress on membrane stability index (MSI), lipid peroxidation, superoxide dismutase (SOD) activity, chlorophyll content and chlorophyll fluorescence in four *Vigna* genotypes namely (Uttara, T-44, IC530491, IC519330). Stress was imposed for 10 days at vegetative stage (30 days after sowing). Thereafter, excess water was drained to allow recovery in stressed plants. Waterlogging treatment significantly increased lipid peroxidation and SOD activity in all the genotypes, which showed the oxidative injury posed by stress conditions. Chlorophyll content and fluorescence reduced under stress conditions. SOD activity, MSI and chlorophyll content was more in IC530491 and IC519330, T44 as compared to Uttara. Lipid peroxidation was high in Uttara. Though chlorophyll fluorescence reduced in all the genotypes under waterlogging, genotypic differences were non-significant. More efficient antioxidative scavenging to maintain membrane stability and chlorophyll content in black gram was found to be associated with tolerance to waterlogging.

Keywords: Antioxidants, Black gram, Chlorophyll, Membrane stability, Waterlogging

Water logging is a major limitation to crop production worldwide. It may result due to erratic rainfall, undulated land or poor drainage due to heavy soil texture. Oxygen diffusion is very much restricted (10000 times slower) in waterlogged conditions compared to air. In addition, respiration by plant roots and microorganisms exaggerate the oxygen deficiency in the rhizosphere¹. Short term waterlogging may result in hypoxia (oxygen deficiency) and if prolonged, it may lead to anoxia (absence of oxygen). Therefore, oxidative phosphorylation is hampered and due to low energy supply, plants adapt to anaerobic metabolism.

Due to unavailability of oxygen, the electron carriers in electron transport chain become reduced, affecting redox state of the cell². Saturated electron transport carriers, altered redox potential of intracellular environment and energy deficit lead to oxidative stress and reactive oxygen species (ROS) are produced. ROS may damage the plant cell by causing lipid peroxidation, enzyme inactivation and oxidative damage to DNA³. Changes in ROS concentration, antioxidant enzyme activities, cell membrane permeability, lipid peroxidation, and

hydrogen peroxide generations are well documented in legumes under waterlogged conditions^{4,5}. Enzymatic as well as non-enzymatic sources generate ROS in plant cells under low O₂ concentrations⁵⁻⁷. An interplay between antioxidant activity and ROS production plays an important role in determining the response of plant to waterlogging⁸. The activities of antioxidant enzymes increased under waterlogging in pigeonpea⁴, mungbean⁵ and citrus⁹.

India is the world's largest producer and consumer of pulses predominantly tropical and sub-tropical legumes such as chickpea, black gram, red gram, green gram and lentil. To ensure self-sufficiency, the requirement for pulses in the country is projected at 39 million tons by the year 2050 at an annual growth rate of 2.2%¹⁰. Black gram (*Vigna mungo* L.) is an important legume crop of rainfed agriculture. Crop is generally sown during *kharif* season. Being a legume, black gram is highly sensitive to waterlogged conditions. Present study was undertaken to study the effect of waterlogging stress on physiological and biochemical responses in selected black gram genotypes at early growth stage.

Materials and Methods

Plant material and waterlogging treatments

A pot experiment was conducted with four black gram lines namely Uttara, T-44, IC530491,

*Correspondence:

Phone (Mob): +91-9599276680

E-mail: Ruchi.Bansal@icar.gov.in

IC519330) in *kharif* 2018. The experimental lines (Uttara, IC530491, IC519330) were selected from a set of 290 lines, which was screened in a preliminary screening in augmented block design during *kharif* 2016 and 2017 for waterlogging tolerance. During previous screening, Uttara did flowering but no pod formation occurred, while, IC530491 and IC519330 produced seeds. Since, we could not find any study on waterlogging tolerance in *Vigna mungo* L., we used T-44 (an already reported waterlogging tolerant *Vigna radiata* genotype) as check. A pot experiment was conducted in complete randomized design with three replications. Plants were grown in plastic pots (diameter 15 cm and height 15 cm) filled with 1 kg sand. At the time of sowing, five seeds were sown in each pot. After germination, they were thinned to one plant per pot. The plants were supplemented with half strength Hoagland solution on each alternate day¹¹. Waterlogging stress was implemented 30 days after sowing by keeping the pots in water filled containers. Water level was maintained 2-3 cm above the soil surface. Control plants were watered as per the requirement.

Physiological and biochemical traits

Chlorophyll and fluorescence were measured at the last day of stress on the first fully matured leaf from the top. Membrane stability index, lipid peroxidation and SOD activity was measured in root tissues. After recording physiological traits, the plants were uprooted carefully to avoid damage to the roots. Samples were stored in deep freezer (-20°C) and lipid peroxidation and SOD activity were measured as soon as possible.

Lipid peroxidation was determined by measuring MDA¹². Root tissue (100 mg) was homogenized in 5 mL 5% (w/v) Trichloroacetic acid solution and centrifuged at $10000 \times g$ for 20 min. 0.5 mL supernatant was added to 1 mL 0.5% (w/v) thiobarbituric acid in 20% TCA. The mixture was heated at 95°C for 20 min and immediately cooled in ice bath. The samples were again centrifuged at $10000 \times g$ for 5 min. The absorbance was recorded at 532 nm. The value for non-specific absorption at 600 nm was subtracted. The amount of MDA-TBA complex (red pigment) was calculated from the extinction coefficient as $155 \text{ mM}^{-1} \text{ cm}^{-1}$.

Superoxide dismutase (EC1.15.1.1) assay was measured spectrophotometrically¹³. Root sample (100 mg) was homogenized in 10 mL extraction buffer (0.1 M phosphate buffer, pH 7.5 containing

0.5 mM EDTA). Enzyme extract was centrifuged at $10000 \times g$ for 10 min at 4°C and supernatant was collected. Reaction mixture (3.0 mL) consisted of 0.1 mL 1.5 M sodium carbonate, 0.2 mL 200 mM methionine, 0.1 mL 2.25 mM NBT, 0.1 mL 3 mM EDTA, 1.5 mL 100 mM potassium phosphate buffer, 1.0 mL distilled water and 0.1 mL enzyme extract. The reaction was started by adding 0.1 mL riboflavin ($60 \mu\text{M}$) and placing the tubes below a light source of two 15W florescent lamps for 15 min. Absorbance was recorded at 560 nm in spectrophotometer. One unit of SOD activity was defined as 50% inhibition of the basic rate of the reaction. Total soluble protein was determined by Bradford method¹⁴.

To measure MSI, samples (0.5 g) with 10 mL double distilled water in glass vial were incubated in a water bath at 40°C for 30 min. After cooling, electrical conductivity (C1) was recorded with a conductivity meter (Sanco, India). Samples were again incubated in a boiling (100°C) water bath for 10 min and conductivity (C2) was measured. Membrane stability index (MSI) was calculated as follows¹⁵.

$$\text{MSI} = [1 - (C1/C2)] \times 100$$

Chlorophyll content was recorded with a self-calibrating chlorophyll meter (Opti-science, USA) on the fully expanded leaves and expressed as SPAD units. Data was taken with three replications each from control and drought treatment.

Chlorophyll fluorescence (Fv/Fm) was measured on first fully expanded leaves using a handheld chlorophyll fluorometer (Opti-science Inc., Hudson, NH). Data was recorded after full dark adaptation for 30 min. Fluorescence is the ratio of variable fluorescence (difference between maximum and minimum fluorescence, Fv) and maximum fluorescence (Fm).

Data were subjected to analysis of variance for completely randomized design factorial. Differences at $P < 0.01$ were considered statistically significant¹⁶.

Results and Discussion

It is well established that energy deficit induced under waterlogging leads to formation of reactive oxygen species (ROS) viz, free radicals (O_2 , OH), hydrogen peroxide (H_2O_2) and phenols. In response plants activates its ROS scavenging system involving enzymatic [peroxidase (POX), catalase (CAT), superoxide dismutase (SOD), ascorbate peroxidase (APX), glutathione reductase (GR)] and non enzymatic antioxidants (ascorbate, glutathione)¹⁷.

Waterlogging caused significant increase in MDA content after imposing stress in all the genotypes (Fig. 1A). MDA content was the maximum in Uttara (0.89 TBARS content) followed by T-44 (0.58 TBARS content). Percent increase ranged from 43.7-161.7 across the genotypes in comparison to control condition.

MDA content was the lowest in genotype IC519330 (0.46 TBARS content) under stress.

Waterlogging caused increase in activity of SOD in all studied black gram genotypes (Fig. 1B). Under waterlogged conditions, enzyme activities were higher in waterlogging tolerant genotypes (IC519330, T-44 and IC530491) compared to waterlogging




Fig. 1 — Effect of waterlogging stress on (A) Malondialdehyde content; (B) Superoxide dismutase activity; (C) Membrane stability index; (D) Chlorophyll content; and (E) Chlorophyll fluorescence in different black gram genotypes

susceptible genotype. Stress increased SOD activity from 2.6 to 42.7% compared to control in studied genotypes. SOD activity was 26.71 units mg^{-1} protein min^{-1} in IC519330 and 26.3 units mg^{-1} protein min^{-1} in T-44 but was least in Uttara (20.45 units mg^{-1} protein min^{-1}) at final stage of observation.

Membrane stability index decreased significantly as a consequence of waterlogging stress (Fig. 1C). In Uttara, cell membrane injury was drastic and stability reduced more than two folds after imposing the stress. Study showed that extent of injury to cell membrane was less in T-44 (35.24), IC519330 (36.56) and IC530491 (34.76) compared to susceptible genotype (17.32).

In the present study, lipid peroxidation and cell membrane injury increased immediately in black gram genotypes after imposing stress. It reflects reduced membrane stability and damage to biomolecules caused by ROS generated during waterlogging. Previous study suggested that free radical-induced peroxidation of membrane lipids indicates stress-induced damage at the cellular level¹⁸. Waterlogging caused significant increase in MDA content in stress susceptible genotypes. Present results were consistent with prior studies conducted in pigeonpea^{4,19}, mungbean²⁰, Malus²¹, and cotton²². High membrane stability in IC519330 and IC530491 accompanied with less lipid peroxidation shows efficient ROS detoxification in these genotypes. Membrane stability was reduced due to enhanced electrolytic leakage. Maintenance of cell membrane stability in tolerant genotypes revealed efficient scavenging of ROS in these genotypes compared to others.

SOD is the enzyme, which dismutates superoxide radical to H_2O_2 and is usually considered the first line of defense against oxidative stress. Increased SOD activity was associated with increased protection from damage associated with oxidative stress²³. In this study, SOD activity increased on account of waterlogging stress and was higher in T-44, IC519330 and IC530491. Increase in SOD activity was also demonstrated in citrus⁹, pigeonpea⁴ and maize²⁴ during waterlogging. Induction of SOD expression was recorded during waterlogging in pigeonpea⁵. In this study also, high SOD activity increased the ROS scavenging efficiency in waterlogging tolerant genotypes.

As duration of waterlogging stress increases, leaf yellowing starts followed by wilting of the plant.

Hence, chlorophyll content reduced significantly in all the genotypes in response to stress (Fig. 1D). Percent reduction varied 21.0-51.0 among different genotypes. Chlorophyll content was 18.7 SPAD units in IC519330, followed by 17.8 SPAD units in T-44. Chlorophyll content reduced by 51% in Uttara (11.6 SPAD units) under stress.

Chlorophyll fluorescence shows the quantum efficiency of photosystem II. Fluorescence reduced under waterlogging, but reduction was non-significant with respect to genotype as well as treatment (Fig. 1E). Fluorescence values ranged 0.70 to 0.79 across different genotypes under normal and waterlogged conditions, which reflect normal functioning of PSII.

Chlorophyll is the most important pigment required for photosynthesis. Its loss under flooding is well documented and is visible by increased yellowing of leaves^{25,26}. Chlorophyll content reduced in all the genotypes, but tolerant one sustained the stress by relatively maintaining the chlorophyll content. Similar results were reported in mungbean²⁶ and pigeon pea²⁷. Chlorophyll fluorescence, which shows the maximum photosynthetic efficiency of photosystem II has been used in phenotyping studies against different abiotic stress. Damage to light-harvesting complex was reported in tomato²⁸ and mung bean under waterlogging²⁹. In the present study, we could not observe significant differences in fluorescence under control and stress conditions. During waterlogging, photosynthetic parameters are affected adversely and stomatal as well as non stomatal components posed limitation to photosynthesis in pigeonpea²⁷. In present study, reduction in chlorophyll was found to be associated with waterlogging susceptibility, while fluorescence remained unaffected during the stress.

Conclusion

Present study showed that membrane stability as evident by SOD activity and MDA content was directly related to waterlogging induced oxidative injury. Furthermore, more chlorophyll content sustained photosynthetic capacity in black gram genotypes under waterlogging stress.

References

- 1 Jackson MB, Ethylene and responses of plants to soil waterlogging and submergence. *Annu Rev Plant Physiol*, 36 (1985) 145.
- 2 Chirkova TV, Zhukova TM & Bugrova MP, Redox reactions of plant cells in response to short term anaerobiosis. *Vestnik SPBGU3*, (1992) 82.

- 3 Shewfelt RL & Purvis AC, Toward a comprehensive model for lipid peroxidation in plant tissue disorders. *Hort Sci*, 30 (1995) 213.
- 4 Bansal R & Srivastava JP, Antioxidative defense system in pigeonpea roots under water logging stress. *Acta Physiol Plant*, 34 (2012) 515.
- 5 Kumutha D, Ezhilmathi K, Sairam RK, Srivastava GC, Deshmukh PS & Meena RC, Waterlogging induced oxidative stress and antioxidant activity in pigeonpea genotypes. *Biol Plant*, 53 (2009) 75.
- 6 Blokhina OB, Virolainen E & Fagerstedt KV, Antioxidants, oxidative damage and oxygen deprivation stress: a review. *Ann Bot*, 91 (2003) 179.
- 7 Bolwell GP & Wojtaszek P, Mechanisms for generation of reactive oxygen species in plant defence – a broad perspective. *Physiol Mol Plant Pathol*, 51(1997) 347.
- 8 Kato C, Ohshima N, Kamada H & Satoh S, Enhancement of the inhibitory activity for greening in xylem sap of squash root with waterlogging. *Plant Physiol Biochem*, 39 (2001) 513.
- 9 Hossain Z, Lopez-Climent MF, Arbona V, Perez-Clemente RM & Gomez-Cadenas A, Modulation of the antioxidant system in citrus under waterlogging and subsequent drainage. *J Plant Physiol*, 166 (2009) 1391.
- 10 Singh P, Shahi B & Singh KM, Trends of pulses production, consumption and import in India: current scenario and strategies. (2007). Doi: <https://mpr.ub.uni-muenchen.de/81589>.
- 11 Hoagland DR & Arnon DI, The water-culture for growing plants without soil. *Calif Agric Exp Stat Circ*, 347 (1950) 32.
- 12 Heath RL & Packer L, Photoperoxidation in isolated chloroplasts. I. Kinetics and stoichiometry of fatty acid peroxidation. *Arch Biochem Biophys*, 125 (1968) 189.
- 13 Dhindsa RS & Matowe W, Drought tolerance in two mosses: correlated with enzymatic defense against lipid peroxidation. *J Exp Bot*, 32 (1981) 79.
- 14 Bradford MM, A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem*, 72 (1976) 248.
- 15 Sairam RK, Effect of moisture stress on physiological activities of two contrasting wheat genotypes. *Ind J Exp Biol*, 32 (1994) 594.
- 16 Gomez KA & Gomez AA, Statistical procedures for agricultural research. (John Wiley and Sons Inc., New York), 1984.
- 17 Noctor G & Foyer C, Ascorbate and glutathione: keeping active oxygen under control. *Annu Rev Plant Physiol Mol Biol*, 49 (1998) 249.
- 18 Jain M, Mathur G, Koul S & Sarin NB, Ameliorative effects of proline on salt stress-induced lipid peroxidation in cell lines of groundnut (*Arachis hypogea* L). *Plant Cell Rep*, 20 (2001) 463.
- 19 Singh VP, Srivastava JP & Bansal R, Biochemical responses as stress indicator to waterlogging in pigeon pea (*Cajanus cajan* L.). *Indian J Biochem Biophys*, 54 (2017) 300.
- 20 Sairam RK, Dharmar K, Lekshmy S & C Viswanathan, Expression of antioxidant defense genes in mung bean (*Vigna radiata* L.) roots under water-logging is associated with hypoxia tolerance. *Acta Physiol Plant*, 33 (2011) 735.
- 21 Bai T, Li C, Ma F, Feng F & Shu F, Response of growth and antioxidant system to root-zone hypoxia stress in two *Malus* species. *Plant Soil*, 325 (2010) 95.
- 22 Guo WQ, Chen BL, Liu RX & Zhou ZG, Effects of nitrogen application rate on cotton leaf antioxidant enzyme activities and endogenous hormone contents under short-term waterlogging at flowering and boll-forming stage. *Ying Yong Sheng Tai Xue Bao*, 21 (2010) 53.
- 23 Asada K, The water-water cycle in chloroplasts: scavenging of active oxygens and dissipation of excess photons. *Annu Rev Plant Physiol Plant Mol Biol*, 50 (1999) 601.
- 24 Jaiswal A & Srivastava JP, Changes in reactive oxygen species scavenging systems and protein profiles in maize roots in response to nitric oxide under waterlogging stress. *Indian J Biochem Biophys*, 55 (2018) 26.
- 25 Li C, Jianga D, Wollenweber B, Li Y, Dai T & Cao W, Waterlogging pretreatment during vegetative growth improves tolerance to waterlogging after anthesis in wheat. *Plant Sci*, 180 (2011) 672.
- 26 Kumar P, Pal M, Joshi R & Sairam RK, Yield, growth and physiological responses of mung bean [*Vigna radiata* (L.) Wilczek] genotypes to waterlogging at vegetative stage. *Physiol Mol Biol Plants*, 19 (2012) 209.
- 27 Bansal R & Srivastava JP, Effect of water logging on photosynthetic and biochemical parameters in pigeon pea. *Russ J Plant Physiol*, 62 (2015) 322.
- 28 Janowiak F, Else MA & Jackson MB, A loss of photosynthetic efficiency does not explain stomatal closure in flooded tomato plants. *Adv Agric Sci Probl*, 481 (2002) 229.
- 29 Ahmed S, Nawata E & Sakuratani T, Effects of waterlogging at vegetative and reproductive growth stages on photosynthesis, leaf water potential and yield in mungbean. *Plant Prod Sci*, 5 (2002) 117.