



Length-weight relationship, condition factor and gonado-somatic index of *Labeo calbasu* (Hamilton, 1822)

E Vahneichong¹ • Sudhir Kumar Das¹ • Anandamoy Mondal² • Dibakar Bhakta^{1,3}


¹ Department of Fishery Resource Management, Department of Fishery Economics and Statistics; Faculty of Fishery Sciences, WBUAFS, Chakgaria, Kolkata 700 094, West Bengal, India.

² Department of Fishery Economics and Statistics; Faculty of Fishery Sciences, WBUAFS, Chakgaria, Kolkata 700 094, West Bengal, India.

³ ICAR-Central Inland Fisheries Research Institute, Regional Centre, B-12, Hans Society, Harney Road, Vadodara 390 022, Gujarat, India.

Correspondence

Prof Sudhir Kumar Das; Department of Fishery Resource Management, Department of Fishery Economics and Statistics; Faculty of Fishery Sciences, WBUAFS, Chakgaria, Kolkata 700 094, West Bengal, India.

 dassudhirkumar1@gmail.com

Manuscript history

Received 21 July 2017 | Revised 24 March 2018 | Accepted 28 March 2018 | Published online 08 September 2018

Citation

Vahneichong E, Das SK, Mondol A and Bhakta D (2018) Length-weight relationship, condition factor and gonado-somatic index of *Labeo calbasu* (Hamilton, 1822). Journal of Fisheries 6(3): 645–649. DOI: 10.17017/jfish.v6i3.2018.269

Abstract

An attempt was made to study dynamics of length-weight relationship (LWR), relative condition factor (K_n) and gonado-somatic index (GSI) of *Labeo calbasu* from the wetlands of South 24 Parganas district of West Bengal, India. A total of 275 fish specimens (130 to 430 mm) were examined over nine months from December 2006 to August 2007. Specimens were categorized into two groups based on the size at first maturity of 260 mm. The LWR was recorded $W = 0.007269 L^{3.181}$ for Group A (<260 mm) and $W = 0.007367 L^{3.031}$ for Group B (≥ 260 mm). The individuals exhibited isometric growth. The monthly mean K_n values varied from 0.924 to 1.141 with an average value of 1.059. The mean monthly values of GSI varied from 0.544 to 3.301 for male and 0.560 to 9.649 for female. There was significant variation ($P < 0.05$) in the mean values of GSI in different months between sexes.

Keywords: *Labeo calbasu*; length-weight relationship; relative condition factor; gonado-somatic index

1 | INTRODUCTION

Labeo calbasu (Hamilton 1822) commonly called as Kalbasu or Black rohu supports an important commercial fishery in ponds, lakes, rivers, and reservoirs of India. It occurs frequently in the commercial catch in rivers of Narmada, Godavari, Yamuna and Ganga rivers. It is also an important candidate species in culture ponds due to its compatibility with other major carps (Borah *et al.* 2014). This species is distributed throughout India, Pakistan, Bangladesh, Burma and Nepal (Chondar 1999) and com-

mon in natural waters (e.g. Mohsin *et al.* 2013; Chaki *et al.* 2014; Galib *et al.* 2009–2018). Kalbasu was once dominant species in Indian waters till the 1980s but now it is considered a locally threatened species (Ramasamy and Rajangam 2016) and categorized as Vulnerable in West Bengal (Das *et al.* 2006). A range of studies have been carried out on *L. calbasu* (Alam *et al.* 2000; Haroon *et al.* 2002; Khan *et al.* 2007) but these lack representation of individuals from wetlands that has been described in this paper. The condition of a fish is influenced by seasonal changes of gonads and also by the feeding intensity

(Sarkar *et al.* 2017a and 2017b). Studies on the reproductive biology of any species are essential in evaluating the commercial potential of the stocks. Gonado-somatic index (GSI) value gives a clear idea regarding spawning season of any species which enable to set seasonal closures during the reproductive season as per need. In majority of the fishes the number of eggs does not change significantly as the season progress but the gonad weight increases due to an increase in water content or organic matter derived from food or organic matter transferred from somatic tissues (Bagenal 1978). Sporadic works have been done on the length-weight relationship, condition factor and GSI of *L. calbasu* (Alam *et al.* 2000) but work on *L. calbasu* from wetlands is scant. So, an attempt was made to study different biological indices of *L. calbasu* in the wetlands of South 24 Parganas district of West Bengal to bridge the gap.

2 | METHODOLOGY

The sample constituted 275 specimens of *L. calbasu* with size ranged from 130 to 430 mm in total length collected from various wetlands (Bhara Haripota Bheri, Trees Bigha and Bees Bigha) of South 24 Parganas district of West Bengal during the period of investigation of nine months from December 2006 to August 2007 (Figure 1). About 30 specimens were collected each month during morning hours (0600 – 0700 hours) from wetlands. The specimens were caught by cast net (mesh 25–30 mm) and drag net (mesh 5–30 mm). The specimens were brought to the laboratory of Department of Fisheries Resource Management, Faculty of Fishery Sciences, WBUAFS, Chakgaria, Kolkata, West Bengal.



FIGURE 1 Study area of wetlands of South 24 Parganas district of West Bengal (source: Google)

The total length of fish was measured with a millimetre scale to the nearest millimetre and body weights were measured by mono-pan balance to the nearest gram. The fishes were categorized into two groups *i.e.* Group A (<260 mm) and Group B (≥ 260 mm) for convenience of interpretation. Le Cren (1951) proposed a non-linear equation in the form of $W = aL^b$, which explains the rela-

tionship between length (L) and weight (W) of fish. Isometry in growth was tested by employing student t-test. The relative condition factor (K_n) was calculated by using the Froese (2006) formula $K_n = W_o/\hat{W}$, where W_o is observed weight and \hat{W} is calculated weight. The GSI was calculated according to Strum (1978) as follows: $GSI = \text{weight of gonad}/\text{weight of fish} \times 100$.

3 | RESULTS AND DISCUSSION

A total sample of 275 fish with size ranged from 130 to 430 mm in total length was examined for their bioindices during nine months of study. It was found that the length at first maturity was 260 mm. The numbers of sample specimens were 119 and 156 for Group A (<260 mm) and Group B respectively. The length-weight relationship recorded for corresponding groups were $W = 0.007269 L^{3.181}$ ($\text{Log } W = -2.138 + 3.181 \text{ Log } L$) and $W = 0.007367 L^{3.031}$ ($\text{Log } W = -2.133 + 3.031 \text{ Log } L$). The b value did not vary significantly from 3 indicating isometric growth in the present study. It bears resemblance with earlier works on *L. calbasu* which was being 3.105 in Bhavanisagar reservoir (Natarajan 1971), 3.00 in Loni Reservoir of Madhya Pradesh (Pathak 1975), 3.105 in immature males in Nagarjunasagar Reservoir (Vinci and Sugunan 1981), 3.109 in fish of size 16 to 46 cm in Sylhet basin, Bangladesh (Alam *et al.* 2000). Sani *et al.* (2010) reported b value of 2.94 of *L. calbasu* from the Betwa (Yamuna River tributary) and Gomti (Ganga River tributary) rivers. Froese (2006) mentioned the value of regression coefficients b of family Cyprinidae is ranged from 2.5 to 3.5. In the present study, the b value resemble with earlier works and the smaller size group (Group A) exhibited relatively higher b value compared to the larger fishes (Group B) indicating more growth rate which bears similarity with earlier works in *Liza tade* (Das 2004) and in *Chelon planiceps* (Pramanick *et al.* 2017). Several researchers (Chatterji *et al.* 1980; Haroon *et al.* 2002; Rizvi *et al.* 2012) have reported allometric growth but Pathak (1975) and Khan (1988) have reported isometric growth pattern in *L. calbasu*.

The monthly mean pooled relative condition factor (K_n) values varied from 0.924 (August) to 1.141 (June) (Figure 2). The condition factor provides ideas on important aspects of feeding, spawning and other aspects related to the well-being of the fish. A high K_n value from April to June corresponds to the maturity of gonads and fall of the value from July to August might be due to spawning.

Rao and Rao (1972) and Choudhary *et al.* (1982) observed high K_n values of *L. calbasu* from the Godavari River and Brahmaputra River which might be due to high feeding tendency of the fishes. The condition factor (K) for *L. calbasu* varied from 0.79 – 1.81 ± 0.150 from the River Chenab of Punjab in Pakistan (Naeem *et al.* 2012). In the pre-

sent study the K_n value found to be higher in smaller size fishes which might be due to relatively more growth and it is in agreement with the earlier workers (Das 2004; Pramanick *et al.* 2017)). Condition factor in fishes generally influenced by several factors like food abundance and physicochemical characters of the environment (Ranganathan and Natarajan 1970), age and sex of fish (Everhart *et al.* 1975), spawning (De Silva and Silva 1979), sex and maturity (Gowda *et al.* 1987), environmental condition, breeding, feeding (Dhanze and Dhanze 1997) and pollution (Sandhya and Shameem 2003; Rao *et al.* 2005).




FIGURE 2 Monthly mean variations of pooled relative condition factor (K_n)




FIGURE 3 Monthly mean variations in gonado-somatic index of *Labeo calbasu*

The monthly mean values of GSI for the male varied from a minimum of 0.544 during the month of December to maximum of 3.301 during June. In case of females, the corresponding values were 0.560 and 9.649 (Figure 3). An increasing trend of GSI values was noticed from December towards June in both sexes. There was significant variation ($P < 0.05$) in the mean values of GSI in different months as well as in both sexes which were due to gonadal maturity. Rao and Rao (1972) found that the peak breeding season of *L. calbasu* was observed in the month of June in Godavari River. But Pathak (1975) and Vinci and Sugunan (1981) reported July as the peak breeding season of *L. calbasu* in the Loni and Nagarjunasagar reservoirs respectively. In the present study, the peak spawning season was observed during the month of June and July which is in agreement with earlier studies. However, the dynamics of increasing gonad weight in proportion to body weight may change from year to year depending on environmental factors such as food availability and temperature (Delahunty and de Vlaming 1980).

4 | CONCLUSION

The length-weight relationship was found to be isometric and K_n value indicated that *L. calbasu* exhibits healthy and robust conditions, showing good compatibility with the studied wetlands. Knowing of stock assessment of certain species is prerequisite for sustainable management of the fishery in a region. This data will provide some important information for the carp species that are present in the wetlands to improve the management strategies in natural habitats.

ACKNOWLEDGEMENTS

The authors are thankful to the Dean, Faculty of Fishery Sciences, Panchasayar, Chakgaria, Kolkata, West Bengal for providing necessary facilities to conduct the study. Thanks are also due to the reviewers for their valuable comments to make the manuscript in presentable form.

REFERENCES

- Alam M, Amin SMN and Haroon AKY (2000) Population dynamics of *Labeo calbasu* (Hamilton) in the Sylhet basin, Bangladesh. *Indian Journal of Fisheries* 47(1): 1–6.
- Bagenal TB (1978) Fish fecundity and its relations with stock and recruitment. *Rapp. P-V. Rapports et procès-verbaux des réunions* 164: 186–98.
- Borah BC, Gogoi R and Rahman A (2014) Performance of orangefin labeo, *Labeo calbasu* (Hamilton 1822) as a component of polyculture system. *Journal of Fisheries* 2(1): 86–89.
- Chaki N, Jahan S, Fahad MFH, Galib SM and Mohsin ABM (2014) Environment and fish fauna of the Atrai River: global and local conservation perspective. *Journal of Fisheries* 2(3): 163–172.
- Chatterji A, Ali M and Mumtaz A (1980) A note on the length-weight relationship of kalbasu, *Labeo calbasu* (Ham.). *Indian Journal of Fisheries* 27(1–2): 261–263.
- Chondar SL (1999) *Biology of fin and shell fish*. SCSC Publishers (India), Howrah, India. 514 pp.
- Choudhary M, Kolekar V and Chandra R (1982) Length-weight relationship and relative condition factor of four Indian major carps of river Brahmaputra, Assam. *Journal of the Inland Fisheries Society of India* 14(2): 42–48.
- Das P, Mahanta PC and Pandey AK (2006) Fish genetic resources and their conservations. In: *Handbook of Fisheries and Aquaculture*, ICAR, New Delhi, India. pp. 31–55.
- Das SK (2004) Biology of length weight and condition in a brackish water fish *Liza tade* (Forsk.). *Indian Journal of Environment and Ecoplanning* 8(1): 209–211.
- De Silva SS and Silva EIL (1979) Biology of young grey mullet, *Mugil cephalus* L., populations of a coastal lagoon in Sri

- Lanka. Journal of Fish Biology 15: 9–20.
- Delahunty G and de Vlaming VL (1980) Seasonal relationship of ovary weight, liver weight and fat stores with body weight in the goldfish, *Carrasius auratus* (L.). Journal of Fish Biology 16: 5–13.
- Dhanze R and Dhanze JR (1997) Biology of scale carp and grass carp, length-weight relationship and growth performance under the agro-climatic zone of Himachal Pradesh. Indian Journal of Fisheries 44(3): 255–263.
- Everhart WH, Eiper AW and Youngs WD (1975) Principles of Fishery Science. Cornell University Press, New York. 228 pp.
- Froese R (2006) Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. Journal of Applied Ichthyology 22: 241–253.
- Galib SM, Lucas MC, Chaki N, Fahad FH and Mohsin ABM (2018) Is current floodplain management a cause for concern for fish and bird conservation in Bangladesh's largest wetland? Aquatic Conservation: Marine and Freshwater Ecosystems 28(1): 98–114.
- Galib SM, Naser SMA, Mohsin ABM, Chaki N and Fahad MFH (2013) Fish diversity of the River Choto Jamuna, Bangladesh: Present status and conservation needs. International Journal of Biodiversity and Conservation 5(6): 389–395.
- Galib SM, Rashid MA, Chaki N, Mohsin ABM and Joadder MAR (2016) Seasonal variation and community structure of fishes in the Mahananda River with special reference to conservation issues. Journal of Fisheries 4(1): 325–334.
- Galib SM, Samad MA, Mohsin ABM, Flowra FA and Alam MT (2009) Present status of fishes in the Chalan Beel- the largest Beel (wetland) of Bangladesh. International Journal of Animal and Fisheries Science 2(3): 214–218.
- Gowda G, Shanbhogue SL and Udupa KS (1987) Length-weight relationship and relative condition of grey mullet, *Valamugil seheli* (Forsk.), from Mangalore waters. Indian Journal of Fisheries 34(4): 340–342.
- Haroon AKY, Alam M, Amin SMN, Dewan S and Islam S (2002) Population dynamics of Gangetic major carps from the Sylhet basin, Bangladesh. Indian Journal of Fisheries 49(2): 161–168.
- Khan MA (1988) Biology of *Labeo calbasu* (Ham.) from Tilaiya Reservoir, Bihar length-weight relationship, condition index and feeding habits. Proceeding of National Academy of Sciences, India 56B: 41–70.
- Khan MA, Rashid H, Miah MI and Shahjahan M (2007) Length-weight relationship and condition factor of some selected fish species from 'Haor' area of Kishoregong in Bangladesh. Progressive Agriculture 18(1): 173–178.
- Le Cren ED (1951) The length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca flaviatilis*). Journal of Animal Ecology 20: 201–219.
- Mohsin ABM, Haque SMM, Galib SM, Fahad MFH, Chaki N, Islam MN and Rahman MM (2013) Seasonal abundance of fin fishes in the Padma River at Rajshahi district, Bangladesh. World Journal of Fish and Marine Sciences 5(6): 680–685.
- Naeem M, Bhatti AH and Nouman MF (2012) External morphological study of wild *Labeo calbasu* with reference to body weight, total length and condition factor from the River Chenab, Punjab, Pakistan. International Journal of Biological, Biomolecular, Agricultural, Food and Biotechnological Engineering 6(7): 429–432.
- Natarajan V (1971) Biology and fishery of *Labeo calbasu* (Ham.) in Bhavanisagar Reservoir. Madras Journal of Fisheries 6: 14–56.
- Pathak SC (1975) Length-weight relationship, condition factor and food study of *Labeo calbasu* (Ham.) from Loni reservoir (M.P.). Journal of the Inland Fisheries Society of India 8: 58–64.
- Pramanick S, Das SK, Bhakta D and Johnson C (2017) Length-weight relationship and condition factor of Tade gray mullet, *Chelon planiceps* (Valenciennes, 1836) from Hooghly-Matlah Estuary, West Bengal, India. Journal of Fisheries 5(1): 469–472.
- Ramasamy M and Rajangam S (2016) Threatened species of IUCN red list: *Labeo calbasu* (Hamilton, 1822) with requirement of imperative conservational management from Lower Anicut, Tamil Nadu, India. International Journal of Fisheries and Aquatic Studies 4(1): 294–303.
- Ranganathan V and Natarajan V (1970) Studies on the occurrence and biology of *Rhinomugil corsula* Hamilton in Krishnagiri and Sathanur reservoirs, Tamil Nadu. Journal of Bombay Natural History Society 66: 478–592.
- Rao GRM and Rao LH (1972) On the biology of *Labeo calbasu* (Ham.-Buch.) from the river Godavari. Journal of the Inland Fisheries Society of India 4: 74–86.
- Rao LM, Lakshmi BB and Bangaramma Y (2005) Length-weight relationship and condition factor of *Liza parsia* (Hamilton-Buchanan) in relation to industrial pollution. Indian Journal of Fisheries 52(3): 345–349.
- Rizvi AF, Amitabh CD and Singh KP (2012) Growth pattern of *Labeo calbasu* (Ham) affected by Sociogeographical condition of the riverine habit. Current Research Journal of Biological Sciences 4(3): 250–257.
- Sandhya V and Shameem U (2003) Length-weight relationship and condition factor of *Liza macrolepis* (Smith) from polluted and unpolluted waters of Visakhapat-

nam. Indian Journal of Fisheries 50(4): 543–546.

Sani R, Gupta BK, Sarkar UK, Pandey A, Dubey VK and Lakra WS (2010) Length-weight relationship of 14 Indian freshwater fish species from the Betwa (Yamuna River tributary) and Gomti (Ganga River tributary) rivers. Journal of Applied Ichthyology 26(3): 456–459.

Sarkar M, Das SK, Mondal A and Bhakta D (2017a) Length-weight relationship and relative condition factor of carps *Labeo bata* and *Labeo rohita* from Kulia beel (wetland) of Nadia district of West Bengal. Journal of Entomology and Zoology Studies 5(5): 1033–1036.

Sarkar S, Das SK and Bhakta D (2017b) Length weight relationship and relative condition factor of Indian shad, *Tenualosa ilisha* from Hooghly estuary system, West Bengal. Journal of Inland Fisheries Society of India 49(1): 22–26.


Strum LMG (1978) Aspects of the biology of *Scombemorus maculatus* (Mitchill) in Trinidad. Journal of Fish Biology 13: 155–172.


Vinci GK and Sugunan VV (1981) Biology of *Labeo calbasu* (Ham.) of the Nagarjunasagar Reservoir (A.P.). Journal of the Inland Fisheries Society of India 13(2): 22–39.

CONTRIBUTION OF THE AUTHORS

EV & AM primary data collection; **DB** data analysis and manuscript preparation; **SKD** research supervision



Sudhir Kumar Das  <https://orcid.org/0000-0003-2229-2887>

Dibakar Bhakta  <https://orcid.org/0000-0002-3871-3028>