

Biosystems Diversity

ISSN 2519-8513 (Print)
ISSN 2520-2529 (Online)
Biosyst. Divers., 26(4), 292–302
doi: 10.15421/011844

Influence of flow velocity, river size, a dam, and an urbanized area on biodiversity of lowland rivers

K. N. Ivicheva*, N. N. Makarenkova*, V. L. Zaytseva*, D. A. Philippov**

*L. S. Berg State Research Institute on Lake and River Fisheries, Vologda, Russia

**Papanin Institute for Biology of Inland Waters of Russian Academy of Sciences, Borok, Russia

Article info

Received 17.09.2018

Received in revised form
25.10.2018

Accepted 29.10.2018

L. S. Berg State Research
Institute on Lake and River
Fisheries, Levicheva st., 5,
Vologda, 160012, Russia.

Papanin Institute for Biology
of Inland Waters of Russian
Academy of Sciences,
Borok, 152742,
Yaroslavl oblast, Russia.
Tel.: +7-485-472-40-42.
E-mail:
philippov_d@mail.ru

Ivicheva, K. N., Makarenkova, N. N., Zaytseva, V. L., & Philippov, D. A. (2018). Influence of flow velocity, river size, a dam, and an urbanized area on biodiversity of lowland rivers. *Biosystems Diversity*, 26(4), 292–302. doi:10.15421/011844

Biodiversity of aquatic organisms is formed under the influence of not only natural, but also anthropogenic factors. In this work, the influence of the flow velocity, river size, flow regulation and urbanization on various groups of aquatic organisms was studied in several lowland rivers. The study was conducted in 2013 on six tributaries of the Upper Sukhona River. Five sampling sites were in different parts of the Vologda River and five sites on small rivers, Losta, Lukhta, Komya, Chernyj Shingar, and Belyj Shingar (one site per river). Phytoplankton, zooplankton and zoobenthos were sampled six times, from April to October, and macrophytes were studied in August. In total, 469 species of aquatic organisms were found in the tributaries of the Upper Sukhona River, belonging to the following phyla: Cyanophyta (5 species), Chrysophyta (8), Bacillariophyta (62), Xanthophyta (1), Cryptophyta (10), Dinophyta (4), Euglenophyta (12), Chlorophyta (17), Streptophyta (1), Bryophyta (2), Marchantiophyta (3), Equisetophyta (1), Magnoliophyta (63), Rotifera (22), Cnidaria (1), Platyhelminthes (1), Annelida (29), Mollusca (33), Arthropoda (194). The maximum number of species was found in the Vologda River, the largest of all the tributaries. The number of zoobenthos species was similar at different sites in the Vologda River and in the small rivers. The number of species of other groups of aquatic organisms in the small rivers was lower than those registered in the Vologda River. The greatest number of macrophyte and zoobenthos species was recorded in the Upper Vologda River and Belyj Shingar River, where the flow is strong all the year round. The greatest number of phyto- and zooplankton species was found at the extra-city sites where current is almost absent. In the dam backwater, species richness was higher than that registered downstream of the dam. At the same time, the species richness of macrophytes and zoobenthos in the dam backwater was lower. The smallest number of species was found in the Vologda River, downstream of the city of Vologda. Decreases in the species richness and Shannon's biodiversity index were witnessed in the Vologda River city site and in the small rivers, as they get closer to the city. Cluster analyses performed for the studied groups of aquatic organisms showed dissimilar results; however, the studied sites on the Vologda River having the highest anthropogenic load formed a cluster. Aquatic organisms of the Upper Sukhona tributaries experience both natural (flow velocity and size of the watercourse) and anthropogenic factors (proximity to the city and flow regulation).

Keywords: macrophytes; phytoplankton; zooplankton; zoobenthos; anthropogenic load; species richness

Introduction

The biological diversity of rivers is a well established indicator of their ecological status. With environmental degradation, the greatest decrease in species diversity is observed in freshwaters rather than in terrestrial ecosystems (Sala et al., 2000). The species richness in rivers varies under the influence of both natural and anthropogenic factors. Natural factors include changes in the hydrological characteristics of the flow in different parts of the river and extreme natural events (Alimov et al., 2013). It is believed that biodiversity also depends on the size of the watercourse (Alimov et al., 2013; Vorste et al., 2017). Anthropogenic load leads to changes in the habitat conditions of aquatic organisms and, as a consequence, the loss of the most sensitive species. There are four groups of factors threatening the biodiversity of rivers, namely: water management, pollution, drainage basin disturbance and biotic factors (Vörösmarty et al., 2010). According to another approach, there are five categories of threats to freshwater biodiversity: over-exploitation, pollution, water level fluctuations, habitat disturbance, and invasions (Dudgeon et al., 2006).

Alteration of the hydrological regime of a watercourse with a dam causes changes in habitat conditions. Consequently, various communities of aquatic organisms are formed upstream and downstream of the dam. It is believed that the species richness of plankton is significantly higher in the reservoirs than in the downstream areas (Alhassan et al., 2015; Fan

et al., 2015). Upstream of the dam, the accumulation of pollutants by sediments can lead to a decrease in the diversity of the bottom dwelling aquatic organisms, compared with the downstream areas. However, the species richness in the backwater areas may either be lower and higher, depending on the specific habitat conditions (Mbaka & Mwaniki, 2015). The channels in the banks affect the species richness and diversity of zoobenthos negatively due to the habitat disturbance (Horsak et al., 2009).

There are different types of pollution, including heat pollution, eutrophication, toxification. In natural ecosystems, pollution is usually complex. With increasing pollution, species diversity decreases due to the elimination of the most sensitive organisms (Barinova et al., 2008; Lock et al., 2011; Karpova & Klepets, 2014; Wright et al., 2017). Disturbance of drainage basins includes various types of agriculture as well as areas impermeable to runoff, including urban areas. Thus, negative impacts of agriculture on aquatic species diversity was reported by Carvalho et al. (2011), Opiso et al. (2014), and Kim et al. (2016); the influence of urban areas was characterized by Beixin et al. (2012), Lakew & Moog (2015). Urban areas exert multiple pressures on river ecosystems, undermining their biodiversity (Rusanov & Khromov, 2016; Grizzetti et al., 2017; Kuzmanovic et al., 2017). Human population density can cause a negative impact on biodiversity (Luck, 2007). Thus, it seems highly relevant to assess the influence of natural and anthropogenic factors on the biodiversity of river ecosystems as an indicator of the ecological status of both a water body and its basin.

The purpose of this study was to determine the species diversity of several groups of aquatic organisms, namely, higher aquatic plants, phytoplankton, zooplankton, and zoobenthos, in lowland rivers, with an emphasis on the changes in species diversity caused by disturbance of the hydrological regime and the complex impact of urban areas.

Material and Methods

The study area is located in the southern taiga, in the south of the central part of the Vologda Region, northwest Russia (Fig. 1). All rivers studied are the tributaries of the Upper Sukhona River. The drainage basins of these rivers are located within three landscapes with different morphological features; in the south-west, the largely paludified Priskhonskaya lowland is adjoined by the terraced Vologda-Gryazovets upland, and in the south-east, by the dome-shaped Avniga upland (Vorobyov, 2007). The flat terrain of the Priskhonskaya lowland and backwaters of the Sukhona River support the hydrological characteristics of the rivers located in the area; during the baseflow period, the almost complete absence of current causes intensive sedimentation. The Vologda-Gryazovets upland is characterized by a long history of development and a high degree of urbanization. In the territory of the Priskhonskaya lowland, economic activity is low on account of the significant degree of paludification.

Field studies were conducted on six water bodies: five small rivers located at different distances from the city, and one medium river, the Vologda River. In the Vologda river, sampling was carried out at five sites, namely: in the upper part (1a, Fig. 1, Table 1), in the reservoir (1b), one in the reservoir downstream of the dam, upstream of the city of

Vologda (1c), within the city (1d), and downstream of the city (1e). Only in the Upper Vologda River (1a), is the current observed during the whole vegetation period. Sampling site 1b is located in the backwater of a dam and is characterized by the greatest depths. Sampling sites 1c, 1d, 1e are located downstream of the dam and have hydrological conditions similar to each other. The Vologda River passing through the city receives industrial discharges and rainwater sewage. In four studied small rivers, the Lukhta, Komya, Chernyj Shingar, Belyj Shingar, sampling sites were located in lower courses, the Losta River had a sampling site in the middle course. In these rivers, except the Belyj Shingar River, the flow velocity is very low, almost absent; only the Belyj Shingar River due to passing through the steep slope of the Avniga Upland has a pronounced flow throughout the growing season. The drainage basins of the small rivers are characterized by varying degrees of disturbance (Ivicheva & Filonenko, 2017): with the proximity to the city, the share of forests decreases and the share of open territories increases. The least affected by human activity is the drainage basin of the Chernyj Shingar River (it is located in the heavily paludified Priskhonskaya lowland), the most affected is the drainage basin of the Lower Vologda River (within and downstream of the city).

Hydrochemical analysis was performed in the Accredited Testing Laboratory of the Federal State Institution of the State Agrochemical Service Center "Vologodskiy" (accreditation certificate No POCC RU.0001.21ПЧ08). The territory genesis determines the high content of iron, copper and zinc in the river water. The waters in the studied rivers belong to the hydrocarbonate class, calcium group, with high TDS levels. Hydrocarbonates predominate in the anionic complex, which is a characteristic feature of water in the entire Vologda Region (Vorobyov, 2007).




Fig. 1. Sampling sites on the tributaries of the Upper Sukhona River: rivers and sampling sites are listed in Table 1.

Table 1

Characteristics of the sampling sites

Site Index	River, sampling site	Coordinates	D, km	W, m	d, m	V, m/s	Bottom substrate	pH	TDS, mg/l
Vologda									
1a	– upstream	59°24'18" N, 39°14'54" E	101.2	13	0.3–1.0	0.4	coarse sand, gravel, boulders	7.1	558
1b	– reservoir	59°16'47" N, 39°42'21" E	57.8	42	0.5–5.0	0.01	silted sand	7.3	726
1c	– downstream of the dam	59°14'25" N, 39°47'56" E	46.0	36	0.5–3.5	0.01	silted sand	7.6	686
1d	– city of Vologda	59°13'28" N, 39°53'23" E	32.4	70	0.5–4.0	0.01	sand, detrital matter	7.2	681
1e	– downstream of the city	59°13'59" N, 40°01'28" E	16.3	90	0.5–3.0	0.01	silt, detrital matter	7.1	588
2	Losta	59°09'20" N, 40°01'25" E	23.4	3–4	0.5–1.2	0.01	silt, detrital matter	6.9	490
3	Lukhta	59°01'56" N, 40°15'51" E	10.0	3–5	0.2–1.0	0.01	sand, detrital matter	6.9	683
4	Komya	59°02'49" N, 40°18'28" E	7.8	4–5	0.3–1.0	0.01	sand, detrital matter	6.9	586
5	Chernyj Shingar	59°10'13" N, 40°38'47" E	3.8	3–5	1.0–1.5	0.01	silt, detrital matter	7.0	507
6	Belyj Shingar	59°10'42" N, 40°39'53" E	3.2	2.5–3.5	0.2–0.6	0.4	sand	7.0	522

Note: D – distance to the mouth of the river, W – river width, d – depth, V – flow velocity during the baseflow period, TDS – total dissolved solids.

Ranunculaceae (5 each), Cyperaceae, Polygonaceae, and Scrophulariaceae (4 each), Asteraceae, Brassicaceae, Lamiaceae, and Lemnaceae (3 each). These top 10 families account for 56.3% of the entire river flora (40 species). The foremost genus was *Potamogeton* (6 species). Another 9 genera have 2 species each (*Bidens*, *Eleocharis*, *Epilobium*, *Glyceria*, *Juncus*, *Persicaria*, *Rorippa*) or 3 species (*Ranunculus*, *Veronica*). The core of the aquatic flora represented 54.9% or 39 species: 18 species of hydrophytes (Lemnaceae, Potamogetonaceae, *Batrachium*, *Elodea*, *Nuphar*, *Myriophyllum*, *Chara*), 10 species of helophytes (*Alisma*, *Butomus*, *Equisetum*, *Sagittaria*, *Schoenoplectus*, *Phragmites*, *Typha*, *Sparganium*), 11 species of hygrophilophytes (*Eleocharis*, *Glyceria*, *Rorippa*, *Veronica*). The shoreline-aquatic plant complex was not rich (partly because only those plants that inhabit the narrow shoreline area were included in the list). It has 32 species, 25 of which are – hygrophytes, 3 – hygromesophytes, 4 – mesophytes.

The composition of macrophytes, in general, was typical for water bodies in this part of the Vologda Region (Lobunicheva et al., 2013; Philippov & Bobrov, 2016). Interesting findings includes some relatively rare macrophytes: *Potamogeton* × *sparganifolius* (Upper Vologda River), *P. crispus* (near the dam), *Riccia cavernosa* (sandbanks near the dam), and *Chara vulgaris* (in the reservoir and near the dam). The last two

findings were published in separate papers earlier (Sofronova et al., 2015; Vishnyakov & Philippov, 2018). As for the adventive plants, *Elodea canadensis* and *Echinocystis lobata* were found.

The size of the flora in the studied rivers varied greatly: from 9–12 species in small rivers – the Losta, Lukhta, Komya, Chernyj Shingar, Belyj Shingar, to 65 species in the Vologda River. The richness of the flora in the Vologda River can be explained not only by greater number of studied sites, but also by the more diverse conditions, which include both naturally preserved areas (upper course) and anthropogenically transformed (reservoir, dam, city). As the degree of anthropogenic impact on the river and its drainage basin increases, the number of macrophyte species decreases (from 33 in the upper course to 10 in the center of the city of Vologda). Downstream of the city, the species richness is also small (14 species), with most of the species found in the shoreline-aquatic ecotone area.

Cluster analysis (Fig. 2a) showed that all studied sites can be divided into two groups. The first group includes the most urbanized sites of the Vologda River, center of the city and downstream of the city. All other sites made up the second group. All the small rivers united in one subgroup, and the Vologda River – into another; the Belyj Shingar River site was the most specific.




Fig. 2. Similarity of macrophyte flora (a), algae flora (b), zooplankton fauna (c) and zoobenthos fauna (d) based on the Bray-Curtis Index: rivers and sampling sites are listed in Table 1

Phytoplankton. In the studied rivers, phytoplankton was represented by 118 species, varieties and forms of algae from 8 divisions, 11 classes, 30 families, 49 genera were registered (Table 2). Bacillariophyta had the greatest species richness – 62 species (52.5% of the total amount). Chlorophyta and Euglenophyta had 16 and 12 species (13.6 and 10.2%), respectively. Less represented were Cryptophyta – 10 species (8.5%), Chrysophyta – 8 species (6.8%), Cyanophyta – 5 species (4.2%), Dinophyta – 4 species (3.4%), Xanthophyta – 1 species (0.8%). Families and genera represented by a single species (or intraspecific taxon) were dominant – 40.0% of all families and 61.2% of all genera. Most of the species found belong to the families Naviculaceae (17 species), Euglenaceae (12), Cryptomonadaceae (10), Fragilariaceae (8), Scenedesma-ceae (8), Nitzschaceae (6), Aulacosiraceae (5), Dinobryaceae (5), Surirellaceae (5), Gomphonemataceae (4). The most species-rich genera are

Navicula (11 species), *Cryptomonas* (7), *Aulacoseira* (5), *Nitzschia* (5), *Scenedesmus* (5), *Dinobryon* (4), *Euglena* (4), *Fragilaria* (4), *Gomphonema* (4), *Phacus* (4). In Cryptophyta and Euglenophyta, on average, 5.0 and 3.0 species per genus, respectively, were registered. Phytoplankton in the Vologda River, upstream of the city, were characterized by genus index 3.2. In the small rivers, genus indices were smaller and varied from 1.5 in the Lukhta River to 2.5 in the Chernyj Shingar River. Genus indices in the Lukhta, Losta (2.0) and Belyj Shingar Rivers (1.9) indicate less favourable environmental conditions that impede the existence of closely related species in the same community. For the first time in the Vologda Region, *Geitlerinema acutissimum*, *Characiopsis spinifer* var. *robusta*, *Crucigeniella truncata*, *Scenedesmus aldavei*, *S. dimorphus* and *Westellopsis linearis* were registered. In the phytoplankton of the rivers, among species with known geographical distribution,

Taxon	1a	1b	1c	1d	1e	2	3	4	5	6
<i>Cladopelma lateralis</i> (Goetgh.)	–	1	–	–	–	–	–	–	–	–
<i>Cladopelma viridula</i> (L.)	–	–	–	1	–	–	–	–	–	–
<i>Cladotanytarsus gr. mancus</i>	1	3	3	–	–	1	1	–	1	3
<i>Constempellina brevicosta</i> (Edw.)*	–	–	1	–	–	–	–	–	–	–
<i>Cryptochironomus gr. defectus</i>	–	1	1	2	–	–	–	–	–	1
<i>Cryptotendipes nigronitens</i> (Edw.)	–	–	–	–	–	–	1	–	–	–
<i>Demicrocryptochironomus vulneratus</i> (Zett.)	1	1	–	–	–	–	–	–	–	–
<i>Dicrotendipes nervosus</i> (Staeg.)	–	1	–	2	–	1	3	1	1	–
<i>Endochironomus albipennis</i> (Meig.)	1	3	2	2	–	–	–	1	1	–
<i>Endochironomus impar</i> (Walk.)	–	1	1	1	–	–	1	1	–	–
<i>Endochironomus tendens</i> (Fabr.)	–	1	–	–	–	–	1	1	–	–
<i>Glyptotendipes gripekoveni</i> (Kief.)	–	2	1	2	+	–	–	–	1	–
<i>Glyptotendipes paripes</i> (Edw.)	–	2	–	1	–	–	–	–	–	–
<i>Harnischia curtilamellata</i> (Mall.)	2	1	1	1	1	–	–	–	–	1
<i>Microchironomus tener</i> (Kief.)	–	1	–	1	1	–	–	–	–	–
<i>Micropsectra gr. praecox</i>	1	–	1	–	–	–	–	–	–	–
<i>Microtendipes pedellus</i> De Geer	2	2	1	1	–	2	1	3	2	1
<i>Parachironomus arcuatus</i> (Goetg.)	–	1	–	1	–	–	–	–	–	1
<i>Parachironomus vitiosus</i> (Goetgh.)	–	–	–	1	–	–	–	–	–	–
<i>Paracladopelma camptolabis</i> (Kief.)	–	–	1	1	1	–	–	–	–	1
<i>Paralauterborniella nigrohalteralis</i> Mall.	–	1	1	1	–	–	–	–	–	1
<i>Paratanytarsus</i> spp.	3	1	1	1	–	2	3	3	1	2
<i>Paratendipes albinus</i> (Meig.)	1	2	–	2	–	2	3	–	1	1
<i>Polypedilum bicrenatum</i> Kief.	–	1	1	–	–	–	3	–	1	1
<i>Polypedilum convictum</i> (Walk.)	2	1	–	1	–	–	2	1	1	1
<i>Polypedilum exectum</i> (Kief.)	–	–	1	–	–	1	1	–	1	1
<i>Polypedilum scalaeum</i> (Schrank)	1	1	1	1	–	1	1	2	2	1
<i>Polypedilum sordens</i> V. d. Wulp	–	–	–	1	–	–	–	–	–	–
<i>Polypedilum nubeculosum</i> (Meig.)	–	2	1	1	1	–	–	–	–	–
<i>Polypedilum pedestre</i> (Meig.)	1	1	–	–	–	–	–	–	–	–
<i>Stempellina bausei</i> (Kief.)	–	–	–	–	–	–	–	–	–	2
<i>Stempellina minor</i> (Edw.)	–	–	–	–	–	–	–	–	–	2
<i>Stictochironomus gr. histrio</i>	–	1	2	–	1	1	2	1	–	2
<i>Tanytarsus</i> spp.	2	2	2	1	–	3	1	2	1	3
<i>Xenochironomus xenolabis</i> Kief.	1	–	–	–	–	–	–	–	–	–
<i>Zavrelia pentatoma</i> Kief.	1	–	–	–	–	–	1	–	1	–
<i>Monodiamesa bathyphila</i> (Kief.)	–	–	–	–	–	–	1	–	–	2
<i>Odontomesa fulva</i> Kief.	–	–	–	–	–	–	–	–	–	2
<i>Pothastia gaedii</i> (Meig.)	1	1	–	–	–	–	–	–	–	2
Total number of invertebrates species	113	98	105	96	38	59	93	78	79	108

Note: rivers and sampling sites are listed in Table 1; asterisk (*) marks new species reported for the territory of the Vologda Region; numbers denote the abundance of species: 1 – sporadic, 2 – common, 3 – dominant.

Based on the results of a cluster analysis performed using phytoplankton abundance, three groups can be distinguished: 1) Vologda River sites; 2) Chernyj Shingar and Komya Rivers; 3) Losta, Lukhta and Belyj Shingar Rivers. Among the Vologda River sites, the similarity with the Upper Vologda River site and the city site (Fig. 2b) was noted. In the Upper Vologda River, a higher flow velocity affects the phytoplankton community; in the city center, phytoplankton experiences anthropogenic stress. These sites have similarity with the site downstream of the dam, where the pronounced flow is also present. The reservoir site is distinguished among other sites due to the most favourable conditions for phytoplankton. Among the small rivers, the first group includes the Chernyj Shingar and Komya Rivers with a relatively low level of anthropogenic load. The second group combines rivers with the most disturbed drainage basins (Losta and Lukhta Rivers), as well as the Belyj Shingar River with relatively high flow velocity.

Zooplankton. In total, 65 zooplankton species were found in the studied tributaries of the Upper Sukhona River (Table 3), the major part of which belongs to Cladocera (34 species); a smaller amount belongs to Rotifera (22 species), and 9 species belong to Copepoda.

The largest number of species (56) were registered in the Vologda River (20 – Rotifera, 27 – Cladocera, 9 – Copepoda). The smallest number of species (19) was recorded in the upper course, which has a swift flow. Only at this site, rotifers made up a little more than a quarter of the total species richness. Other sites are characterized by very slow flows, and as a result, the species composition is significantly richer there. The largest number of species (33) was recorded in the reservoir. Species richness in the reservoir was greater than that at the site downstream of the dam. At the city site, the species composition did not differ significantly, and no reduction in the number of species was observed.

In the small rivers, the number of species varied from 14 to 19. The lowest species richness was observed in the Shernyj Shingar River, characterized by the least anthropogenically disturbed drainage basin. The greatest share of rotifers was witnessed in the Losta River, closest to the city of Vologda. The number of species in the Belyj Shingar River, having a rapid flow, was close to those found in rivers with low flow velocities.

Species diversity (Table 4) was higher at all sites of the Vologda River compared to those in the small rivers. The maximum values of the Shannon index were observed at the reservoir site.

Table 4
Number of species and values of Shannon Index calculated using abundance

Aquatic organisms	Rivers and sites									
	1a	1b	1c	1d	1e	2	3	4	5	6
Number of species										
Macrophyte	33	25	32	10	14	10	10	9	11	12
Phytoplankton	32	46	16	18	11	8	9	15	25	19
Zooplankton	19	33	27	28	25	15	15	19	14	19
Zoobenthos	94	65	78	68	13	44	78	59	65	89
Total number	178	169	153	124	63	77	112	102	115	139
Shannon index										
Phytoplankton	1.86±2.67±1.95±1.98±1.91±1.54±1.74±0.99±1.47±2.90±0.68	0.57	0.63	0.63	0.53	0.34	0.39	0.54	0.45	0.41
Zooplankton	1.09±1.56±1.37±1.11±1.28±1.04±0.90±1.30±0.96±1.08±0.17	0.47	0.23	0.16	0.21	0.16	0.37	0.26	0.32	0.25
Zoobenthos	2.14±1.64±1.78±1.41±0.26±1.62±1.91±1.52±1.85±1.80±0.09	0.12	0.10	0.25	0.09	0.12	0.16	0.17	0.13	0.22

Note: rivers and sampling sites are listed in Table 1; values of Shannon Index are given as mean ± standard error.

Cluster analysis (Fig. 2c) showed that all sites could be divided into two groups. All sites of the Vologda River, except for the upper course, made up one group; the second group included all the small rivers and the Upper Vologda River site. The greatest species similarity was seen between stations 1d and 1e experiencing the greatest anthropogenic load.

Zoobenthos. The fauna of benthic macroinvertebrates was represented by 215 species and taxa of supraspecific rank from 10 classes, 20 orders, and 70 families (Table 3). The most common were insects, representing 69% or 149 species belonging to Ephemeroptera (9.5% or 21 species), Trichoptera (14.9% or 32 species), and Diptera (35.8% or 77 species), among the latter, the most numerous were Chironomidae (29.3% or 64 species). The rest of the fauna was represented by Mollusca (15.8% or 33 species), Oligochaeta (9.9% or 21 species), and Hirudinea (3.6% or 8 species). For the first time in the Vologda Region, 24 species of benthic macroinvertebrates were registered: 10 species of molluscs (*Acroloxus rossicus*, *Anisus charteus*, *Euglesa likharevi*, *Henslowiana infirmicostata*, *H. ostroumovi*, *H. polonica*, *Hiberneuglesa normalis*, *H. parvula*, *Pulchelleuglesa acuticostata*, *Sphaerium westerlundii*) and 14 species of insects (*Ephemerella mucronata*, *E. notata*, *Eurylophella karelica*, *Habrophlebia fusca*, *Libellula fulva*, *Beraeodes minutus*, *Goera pilosa*, *Lepidostoma hirtum*, *Lype phaeopa*, *Notidobia ciliaris*, *Plectrocnemia conspersa*, *Constempellina brevicosta*, *Epoicocladius flavens*, *Prosilocerus danubialis*). Also, in the Belyj Shingar River, in August 2013, we found specimens of *Semblis phalaenoides* – a rare vulnerable species in the Vologda Region (category 3/VU) (Bolotova et al., 2010).

Zoogeographic analysis showed that the basis of the fauna was comprised by the transpalearctic species (40%). Molluscs and caddisflies represented European species (21%); chironomids mostly represented Holarctic species (23%); cosmopolite species (4%) were completely represented by annelids. Species with narrow natural habitat was not found.

The largest number of species (168) was recorded in the Vologda River. From upper course site to the site downstream of the city, a decrease in the number of species was witnessed. In the upper course of the Vologda River, where the most species were found (94 species), rheophilic species of zoobenthos were dominant. Downstream, where the flow rate decreases and sedimentation occurs, the number of species decreases. At the same time, in the reservoir, the species richness was lower than that registered at the site downstream of the dam (65 and 78 species, respectively). At these two sites, the invasive *Dreissena poly-*

morpha was found, forming druses in the reservoir. A decrease in the species composition was observed further down the river; 68 species were registered at the city site and 13 species downstream from the city, where *Limnodrilus hoffmeisteri* was an absolute dominant species with the most occurrence (in 85% of samples); other species were found sporadically.

Among the small rivers, the maximum species richness of zoobenthos was registered in the Belyj Shingar River (89 species), the only river where the flow is evident throughout the growing season. Of all the rivers analyzed, it was here that the greatest species richness of Insecta was observed (71% of all species), Diptera in particular (Chironomidae – 40%, other groups – 11%). In the Chernyj Shingar River, 65 species of benthic macroinvertebrates were found. In the Lukhta River, 78 species were registered, including the greatest number of Trichoptera (10 species). The smallest number of species (44) was found in the Losta River, closest to the city of Vologda.

The highest values of the Shannon Index showed samples from the Upper Vologda River (Table 4). In the middle course and downstream of the city, the indices of species diversity were lower. At the city site, a sharp decrease in the Shannon Index was observed. Among the small rivers, the lowest value of the species diversity index was registered in the Losta River, closest to the city.

Cluster analysis (Fig. 2d) showed significant differences between sites. The first group included all sites on the Vologda River except the upper course. Sites 1b, 1c, 1d, and 1e are located at a short distance from each other, and are affected by the dam. Within this group, the sites make up two sub-groups: upstream of the city (sites 1b, 1c), and within and downstream of the city (sites 1d, 1e). These differences are probably caused by an increasing anthropogenic load. The most specific fauna was observed in the Belyj Shingar River (sites 6). By the richness of fauna, Losta, Lukhta, Komya, and Chernyj Shingar Rivers (sites 2–5) as well as the Upper Vologda River (site 1a) made up the second group. The greatest similarity was witnessed between the faunas of the Losta River (site 2) and the Lukhta River (site 3). The drainage basins of these rivers are the most disturbed anthropogenically.

Discussion

In our study, the cluster analysis of fauna and flora showed slightly different results for different groups of aquatic organisms. In all studied groups, the sites on the Vologda River, within the city and downstream of the city, were distinguished among others. The closest results of cluster analysis were obtained for the zooplankton and zoobenthos which distinguished the fauna of the Vologda River.

The influence of river size and flow velocity. The largest number of species in all studied groups was registered in the Vologda River. At individual sites, the Vologda River generally has a greater number of species than on small rivers, even under similar hydrological conditions. Our results are in agreement with an observation on the dependence of the species richness and diversity (Shannon Index) on the size of the water object (Alimov et al., 2013; Vorste et al., 2017). Thus, an increase in the species richness and diversity of mosses and macroinvertebrates commensurate with an increase in the sizes of rivers was studied in Finland (Heino et al., 2005). The association of drainage basin size with the number of algae species was shown in the Russian Far East (Bogatov & Nikulina, 2010). An increase of the diatoms' diversity and species richness along with river size was witnessed in Central Europe (Stenger-Kovács et al., 2014). According to the river continuum concept, from headwaters to mouth, the physical variables within a river system present a continuous gradient of physical conditions (Vannote et al., 1980). In headwaters, where the flow rate is significantly higher, benthic organisms play the main role; in the lower course, planktonic and benthic communities are equally developed (Alimov et al., 2013). In general, our results confirm this pattern. In our case, with the similarity of most abiotic conditions, the main difference between the sites is the flow velocity. Due to the lowland position of the river drainage basins, most sites are characterized by a very low flow velocity. Only in the upper course of the Vologda River (site 1a) and the Belyj Shingar River (site 6) is the flow pronounced, and, as a consequence, the large species richness of the rheophilic zoobenthos was evident. The work of Breuer et al.

(2016) showed the negative relationships between the flow velocity and the species composition of diatoms. However, in our study, the association between the plankton species richness and the flow velocity was not witnessed.

The influence of a dam. The influence of the dam on different ecologic groups of aquatic organisms was not manifested equally. The species richness and diversity of zooplankton and phytoplankton in the reservoir was much higher, which is in agreement with the results reported by Alhassan et al. (2015). Here, in backwater of the dam, the largest species richness and diversity of phytoplankton in our study was registered. An increase of phytoplankton and zooplankton species richness in backwater of the dam after its construction was seen in the Lancang-Mekong River (Fan et al., 2015). At the same time, organisms associated with the bottom (zoobenthos and macrophytes) were characterized by lower species richness in the reservoir in our study. This might be caused by the accumulation of pollutants in bottom sediments in the backwater area. However, the negative impact of dams on zoobenthos communities is not always evident (Mbaka & Mwaniki, 2015). Due to dam construction, changes in the species structure and an increase in the species richness can occur in homotopic zoobenthos. In our study, homotopic species also prevailed in the backwater area.

The influence of the city. In the Vologda River, downstream of the dam, with a gradual increase in the anthropogenic load, a decrease in the number of species of macrophytes and zoobenthos was witnessed, which corresponds with the results reported by Krylova (2010), Fominykh (2014), Karpova & Klepets (2014), and Philippov & Bobrov (2016). A decrease in the number of macrophyte species, especially typical aquatic plants, and simplification of their coenoses within the city and downstream of the city (sites 1c, 1d and 1e) may indicate an increasing anthropogenic load on the river ecosystem. This pattern can be caused by both the quality of the aquatic environment and the anthropogenic impact on the riverbed, banks and the adjacent areas of drainage basin. At the same time, the species richness and diversity of zooplankton, as well as the species diversity of phytoplankton within and downstream of the city remained at the same level. Apparently, the absence of a pronounced flow and high content of natural organic matter create favourable conditions for the development of these groups of aquatic organisms. For example, Breuer et al. (2016) showed that nutrients are the second limiting factor for phytoplankton development after temperature. Zoobenthos species diversity also decreased at the city site and reached its minimum values downstream of the city. Here, absolute dominance of Oligochaeta was witnessed, which indicates heavy organic pollution of the bottom substrate. A decrease in the species richness and diversity of zoobenthos with increasing organic pollution was also observed by Lock et al. (2011), Arimoro et al. (2015), and Wright et al. (2017).

In the small rivers, the lowest species richness and diversity was registered in the Losta River, which experiences the greatest anthropogenic load. A decrease (compared with the other small rivers) in the number of phytoplankton and zoobenthos species was evident there, apparently as a response to an increase in eutrophication (Mangadze et al., 2016). The drainage basins of the Lukhta and Komya Rivers are largely ploughed while the drainage basin of the Chernyj Shingar River is almost intact. Nevertheless, the species richness and species diversity in these rivers were close.

Aquatic organisms associated with substrate are considered well established indicators of the water quality in rivers (Semenchenko, 2004); which is in agreement with our results. The species richness of phytoplankton decreased with increasing anthropogenic load, while the species diversity remained at the same level. Thus, phytoplankton may also be considered as an indicator of water quality, as it was shown by Mangadze et al. (2016). Zooplankton is more dependent on physiographic factors, which was shown by Kurashov et al. (2017). The agricultural use of the drainage basins had a lesser effect on the species richness and diversity than the degree of urbanization.

Conclusion

The study of biodiversity in the tributaries of the Upper Sukhona River showed that the species richness of rivers with similar hydrolo-

gical conditions was determined by the size of river. In rivers with a pronounced flow, the number of bottom associated species (zoobenthos and macrophytes) was greater than that in rivers where the flow velocity is low most of the year. The dam had a varying influence on the studied groups of aquatic organisms. A burst in the species richness of phytoplankton and zooplankton was witnessed upstream of the dam, in the reservoir; and by contrast, the bottom associated organisms, zoobenthos and macrophytes, showed lower species richness at this part of the river. Increasing anthropogenic load leads to a decrease in the number of macrophyte species, phytoplankton and zoobenthos and the dominance of certain species. In the small rivers, the lowest species richness was observed in the Losta River, closest to the city of Vologda. In the rivers with ploughed and almost intact drainage basins, the species richness remained at the same level. Aquatic organisms inhabiting the tributaries of the Upper Sukhona River are affected by both natural (flow velocity and size of a watercourse) and anthropogenic (proximity to a city and regulation of the river) factors.

Research of D. A. Philippov was performed in the framework of the state assignments of FASO Russia to IBIW RAS (theme No. AAAA-A18-118012690099-2).

References

- Alhassan, E. H., Ofori-Danson, P. K., & Samman, J. (2015). Ecological impact of river impoundment on zooplankton. *Zoology and Ecology*, 25(2), 136–142.
- Alimov, A. F., Bogatov, V. V., & Golubkov, S. M. (2013). *Produkcionnaya gidrobiologiya* [Production Hydrobiology]. Nauka, Saint-Petersburg (in Russian).
- Arimoro, F. O., Odume, O. N., Uhunoma, S. I., & Edegbene, A. O. (2015). Anthropogenic impact on water chemistry and benthic macroinvertebrate associated changes in a southern Nigeria stream. *Environmental Monitoring and Assessment*, 187(2), 14.
- Barinova, S., Medvedeva, L., & Nevo, E. (2008). Regional influences on algal biodiversity in two polluted rivers of Eurasia (Rudnaya River, Russia, and Qishon River, Israel) by bioindication and canonical correspondence analysis. *Applied Ecology and Environmental Research*, 6(4), 29–59.
- Beixin, W., Dongxiao, L., Shuru, L., Yong, Z., Dongqi, L., & Lizhu, W. (2012). Impacts of urbanization on stream habitats and macroinvertebrate communities in the tributaries of Qiantang River, China. *Hydrobiologia*, 680(1), 39–51.
- Bogatov, V. V., & Nikulina, T. V. (2010). Relationship between algal communities species diversity and catchment of rivers in the southern Russian Far East. *Inland Water Biology*, 3(3), 249–253.
- Bolotova, N. L., Ivanter, E. V., & Krivokhatskiy, V. A. (Eds.) (2010). *Krasnaya kniga Vologodskoy oblasti. Tom 3. Zhivotnye* [Red Data Book of the Vologda Region. Vol. 3. Animals]. Vologda (in Russian).
- Breuer, F., Janz, P., Farrelly, E., & Ebke, K.-P. (2016). Environmental and structural factors influencing algal communities in small streams and ditches in central Germany. *Journal of Freshwater Ecology*, 32(1), 65–83.
- Carvalho, L., Cortes, R., & Bordalo, A. A. (2011). Evaluation of the ecological status of an impaired watershed by using a multi-index approach. *Environmental Monitoring and Assessment*, 174, 493–508.
- Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z.-I., Knowler, D. J., Leveque, C., Naiman, R. J., Prieur-Richard, A.-H., Soto, D., Stiassny, M. L. J., & Sullivan, C. A. (2006). Freshwater biodiversity: Importance, threats, status and conservation challenges. *Biological Reviews*, 81(2), 163–182.
- Fan, H., He, D., & Wang, H. (2015). Environmental consequences of damming the mainstream Lancang-Mekong River: A review. *Earth-Science Reviews*, 146, 77–91.
- Fominykh, A. S. (2014). *Vysshaya vodnaya rastitel'nost' basseyna reki Beloy v rayone goroda Sterlitamak* [Higher aquatic vegetation of the Belaya River Basin in the vicinity of Sterlitamak]. *Vodnoe Khozyajstvo Rossii: Problemy, Tekhnologii, Upravlenie*, 4, 17–30 (in Russian).
- Grizzetti, B., Pistocchi, A., Lique, C., Udias, A., Bouraoui, F., & van de Bund, W. (2017). Human pressures and ecological status of European rivers. *Scientific Reports*, 7, 205.
- Heino, J., Paavola, R., Virtanen, R., & Muotka, T. (2005). Searching for biodiversity indicators in running waters: Do bryophytes, macroinvertebrates, and fish show congruent diversity patterns? *Biodiversity and Conservation*, 14(2), 415–428.
- Horsak, M., Bojkova, J., Zahradkova, S., Omesova, M., & Helesic, J. (2009). Impact of reservoirs and channelization on lowland river macroinvertebrates: A case study from Central Europe. *Limnologia*, 39(2), 140–151.
- Ignatov, M. S., Afonina, O. M., Ignatova, E. A., Abolina, A. A., Akatova, T. V., Baisheva, E. Z., Bardunov, L. V., Baryakina, E. A., Belkina, O. A., Bezgodov, A. G., Boychuk, M. A., Cherdantseva, V. Y., Czemyadjeva, I. V., Doroshina, G. Y., Dyachenko, A. P., Fedosov, V. E., Goldberg, I. L., Ivanova, E. I., Jukoniene, I., Kanukene, L., Kazanovsky, S. G., Kharzinov, Z. K., Kurbatova, L. E., Maksimov, A. I., Mamatkulov, U. K., Manakyan, V. A., Maslovsky, O. M., Napreenko, M. G., Omyukova, T. N., Partya, L. Y., Pisarenko, O. Y., Popova, N. N., Rykovsky, G. F., Tubanova, D. Y., Zheleznova, G. V., & Zolotov, V. I. (2006). Check-list of mosses of East Europe and North Asia. *Arctoa*, 15, 1–130.
- Ivicheva, K. N., & Filonenko, I. V. (2017). O vliyani osvoennosti vodosbora reki Verhnej Suhony (Vologodskaja oblast') na himicheskij sostav vod ee pritokov [On the impact of the development of the Verkhnyaya Sukhona river catchment area (the Vologda Region) on the chemical composition of the waters in its tributaries]. *Principy Ekologii*, 6(3), 81–92 (in Russian).
- Karpova, G. A., & Klepets, Y. V. (2014). Influence of urban landscapes on the structural indices of macrophytes in the Vorskla river. *Hydrobiological Journal*, 50(6), 3–16.
- Kim, J.-A., Lee, S.-W., Hwang, G.-S., Hwang, S.-J., Kim, C., & An, K.-J. (2016). Effects of streamline complexity on the relationships between urban land use and ecological communities in streams. *Paddy and Water Environment*, 14(2), 299–312.
- Komárek, J. (2013). Cyanoprocaryota 3. Teil: Heterocytous genera. In: Büdel, B., Gärtner, G., Krienitz, L., & Schlager, M. (eds.) *Stiwwasserflora von Mitteleuropa 19/3*. Springer Spektrum, Berlin, Heidelberg.
- Komárek, J., & Anagnostidis, K. (2005). Cyanoprocaryota 2. Teil: Oscillatoriales. In: Büdel, B., Gärtner, G., Krienitz, L., & Schlager, M. (eds.) *Süswasserflora von Mitteleuropa 19/2*. Elsevier/Spektrum, Heidelberg.
- Komárek, J., & Fott, B. (1983). Chlorophyceae (Grünalgen), Ordnung: Chlorococcales. In: Huber-Pestalozzi, G. (ed.) *Das Phytoplankton des Süswassers; Systematik und Biologie*. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart (in German).
- Konstantinova, N. A., Bakalin, V. A., Andrejeva, E. N., Bezgodov, A. G., Borovichev, E. A., Dulin, M. V., & Mamontov, Y. S. (2009). Checklist of liverworts (Marchantiophyta) of Russia. *Arctoa*, 18, 1–64.
- Krylova, E. G. (2010). Structure of hydrophilic vegetation of a small river in an urban environment. *Inland Water Biology*, 3(2), 119–125.
- Kurashov, E. A., Aleshina, D. G., Guseva, M. A., Petrova, T. N., & Krylova, J. V. (2007). The role of environmental factors in the formation of zooplankton in tributaries of Lake Ladoga (Russia). *Applied Ecology and Environmental Research*, 15(4), 1511–1540.
- Kuzmanovic, M., Dolédec, S., de Castro-Catala, N., Ginebreda, A., Sabater, S., Muñoz, I., & Barceló, D. (2017). Environmental stressors as a driver of the trait composition of benthic macroinvertebrate assemblages in polluted Iberian rivers. *Environmental Research*, 156, 485–493.
- Lakew, A., & Moog, O. (2015). A multimetric index based on benthic macroinvertebrates for assessing the ecological status of streams and rivers in central and southeast highlands of Ethiopia. *Hydrobiologia*, 751(1), 229–242.
- Lobunicheva, E. V., Borisov, M. Y., Filonenko, I. V., & Philippov, D. A. (2013). *Ocenka ekologicheskogo sostojanija malyh vodojmov* [Environmental assessment of small water bodies]. *Izdatel'stvo Kopemik35*, Vologda (in Russian).
- Lock, K., Asenova, M., & Goethals, P. L. M. (2011). Benthic macroinvertebrates as indicators of the water quality in Bulgaria: A case-study in the Iskar river basin. *Limnologia*, 41(4), 334–338.
- Luck, G. W. (2007). A review of the relationships between human population density and biodiversity. *Biological Reviews*, 82(4), 607–645.
- Mangadze, T., Bere, T., & Mwedzi, T. (2016). Choice of biota in stream assessment and monitoring programs in tropical streams: A comparison of diatoms, macroinvertebrates and fish. *Ecological Indicators*, 63, 128–143.
- Martinez, A., Larranaga, A., Basaguren, A., Perez, J., Mendoza-Lera, C., & Pozo, J. (2013). Stream regulation by small dams affects benthic macroinvertebrate communities: From structural changes to functional implications. *Hydrobiologia*, 711(1), 31–42.
- Mbaka, J. G., & Mwaniki, M. W. (2015). A global review of the downstream effects of small impoundments on stream habitat conditions and macroinvertebrates. *Environmental Reviews*, 23(3), 257–262.
- Opiso, E. M., Quimpong, V. T., Leano, E. P., Galan, G. L., Acma, F. M., Coritico, F. P., Labadan, A., Forten, R. R., Coquilla, L., Bruno, A. G., & Amoroso, V. B. (2014). Assessment of biodiversity and water quality in association with land use in the Alanib River, Mt. Kitanglad Range Park, Philippines. *Asian Journal of Biodiversity*, 5, 54–77.
- Philippov, D. A., & Bobrov, Y. A. (2016). *Flora malyh vodojmov goroda Vologdy i analiz ee struktury* [Flora of small reservoirs of Vologda city and its structure]. *Vestnik of Orenburg State Pedagogical University, Electronic Scientific Journal*, 4, 32–44 (in Russian).
- Plantae in GBIF Secretariat (2017). GBIF Backbone Taxonomy. Checklist Dataset (searched on 2018-01-10).
- Rusanov, A. G., & Khromov, V. M. (2016). Longitudinal distribution of periphyton algae in the Moskva river under eutrophication. *Water Resources*, 43(3), 513–521.

- Sala, O. E., Chapin, F. S., Armestro, J. J., Berlow, R., Bloomfield, J., Dirzo, R., Huber-Sanwald, E., Huenneke, L. F., Jackson, R. B., Kinzig, A., Leemans, R., Lpdge, D., Mooney, H. A., Oesterheld, M., Poff, N. L., Sykes, M. T., Walker, B. H., Walker, M., & Wall, D. H. (2000). Global biodiversity scenarios for the year 2100. *Science*, 287(5459), 1770–1774.
- Semenchenko, V. P. (2004). Principy i sistemy bioindikacii tekushchih vod [Principles and systems of bioindication of flowing waters]. Oreh, Minsk (in Russian).
- Sofronova, E. V., Abdurachmanova, Z. I., Afonina, O. M., Akatova, T. V., Andrejeva, E. N., Bakalin, V. A., Bezgodov, A. G., Borovichev, E. A., Czemyadjeva, I. V., Doroshina, G. Y., Dulin, M. V., Fedosov, V. E., Golovina, E. O., Ignatov, M. S., Ignatova, E. A., Kotkova, V. M., Kozhin, M. N., Kučera, J., Kurbatova, L. E., Kushnevskaya, E. V., Leushina, E. G., Makarova, M. A., Maksimova, A. Y., Nikolajev, I. A., Philippov, D. A., Popova, N. N., Potemkin, A. D., Prelovskaya, E. S., Teleganova, V. V., Vilnet, A. A., Volkova, E. M., & Zolotukhin, N. I. (2015). New bryophyte records. 5. *Arctoa*, 24(2), 584–609.
- Stenger-Kovács, C., Tóth, L., Tóth, F., Hajnal, É., & Padišák, J. (2014). Stream order-dependent diversity metrics of epilithic diatom assemblages. *Hydrobiologia*, 721(1), 67–75.
- The Angiosperm Phylogeny Group (2016). An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of the Linnean Society*, 181(1), 1–20.
- Tomwall, B., Sokol, E., Skelton, J., & Brown, B. L. (2015). Trends in stream biodiversity research since the river continuum concept. *Diversity*, 7(1), 16–35.
- Vannote, R. L., Minshall, G. W., Cummins, K. W., Sedell, J. R., & Cushing, C. E. (1980). The river continuum concept. *Canadian Journal of Fisheries and Aquatic Sciences*, 37(1), 130–137.
- Vasser, S. P. (ed.) (1989). *Vodorosli: Spravochnik [Algae: Reference book]*. Naukova Dumka, Kiev (in Russian).
- Vishnyakov, V. S., & Philippov, D. A. (2018). Novye nahodki harovyh vodorosley (Charales) na Evropeyskom Severe Rossii [New records of charophytes (Charales) from the Northern European Russia]. *Botanicheskii Zhurnal*, 103(8), 1016–1031 (in Russian).
- Vorobyov, G. A. (ed.) (2007). *Priroda Vologodskoj oblasti [Nature of the Vologda Region]*. Izdatelskij Dom Vologzhanin, Vologda (in Russian).
- Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., Glidden, S., Bunn, S. E., Sullivan, C. A., Liermann, C. R., & Davies, P. M. (2010). Global threats to human water security and river biodiversity. *Nature*, 467, 555–561.
- Vorste, V. R., McElmurray, P., Bell, S., Eliason, K. M., & Brown, B. L. (2017). Does stream size really explain biodiversity patterns in lotic systems? A call for mechanistic explanations. *Diversity*, 9(3), 26.
- Wright, I. A., Belmer, N., & Davies, P. J. (2017). Coal mine water pollution and ecological impairment of one of Australia's most 'protected' high conservation-value rivers. *Water, Air and Soil Pollution*, 228, 90.