

Article under Review for the International Journal of Spatial Data Infrastructures Research
submitted 2019-09-10

Lost in space and time? A conceptual approach to harmonise data for marine spatial planning*

Wanda Holzhüter¹, Hanna Luhtala², Henning Sten Hansen³
Kerstin Schiele⁴

¹Institute for Baltic Sea Research Warnemünde, Germany; wanda.holzhueter@io-warnemuende.de

²University of Turku, Finland; hanna.luhtala@utu.fi

³Aalborg University, Denmark; hsh@plan.aau.dk

⁴Institute for Baltic Sea Research Warnemünde, Germany; kerstin.schiele@io-warnemuende.de

Abstract

Despite a list of national and international efforts to harmonise data management procedures, the categorisation of space and time within datasets in marine spatial planning (MSP) has not been addressed so far. This paper proposes a conceptual approach to categorise the spatial and temporal dimensions of data used in MSP and introduces a method to jointly manage non-spatial information and spatial data in the same geographic information system (GIS). The presented categorisation provides easy and intuitive classifications for a more detailed and transparent data description of spatial and temporal data properties, which can be applied both in attribute tables and in metadata. It allows the differentiation of the vertical and the horizontal dimensions, enabling users to focus on operations taking place at specific parts of the marine environment. The categorisation with predefined attribute domains allows space and time based automatic analyses. The inclusion of non-spatial data within GIS repositories ensures the availability of all relevant data in one database minimising the risk of incomplete data. Overall, the concept provides effective steps towards a more coherent data management and subsequently may foster better use of information in MSP processes.

Keywords: marine spatial planning, data management, non-spatial data, spatial dimension, temporal dimension, spatial data infrastructure, data harmonisation

*This work is licensed under the Creative Commons Attribution-Non commercial Works 3.0 License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/> or send a letter to Creative Commons, 543 Howard Street, 5th Floor, San Francisco, California, 94105, USA.

DOI: *to be assigned if accepted*

1. INTRODUCTION

Marine spatial planning (MSP) is a complex, data intensive, and evidence-based process (MSP Data Study, 2016). The success of a MSP process largely depends on the quality and availability of pertinent data and the capacity for their analysis (Stamoulis & Delevaux, 2015). Consequently, in a world of data, where it is possible to gather an unlimited amount of datasets, data collection, processing, management and storage need to be handled with great care.

Challenges arise especially through different concepts of data management, e.g. during the transition from local or national planning to international, cross-border operations (Stamoulis & Delevaux, 2015) when different administrative structures, languages and procedures, different stages of planning, and respective data requirements and standards need to be coordinated.

Several international efforts to establish coherence among datasets that are collected at various geographical scales and institutional domains exist. Already in the early 1960s, it was recognised that international efforts are needed to coordinate oceanographic data exchange, which led to the establishment of the International Oceanographic Data and Information Exchange programme (IODE). Since 2009 the European Marine Observation and Data Network (EMODnet¹), a network of more than 150 organisations provides and processes data of the European marine environment. Data, metadata and information are available via the EMODnet portal following international standards and supported by the EU integrated maritime policy².

A major general development in data management is the Directive 2007/2/EC establishing an Infrastructure for Spatial Information in the European Community (INSPIRE) which came into force in the member states of the European Union in 2007 (Bartha & Kocsis, 2011). It aims to improve consistency, availability, and reuse of spatial information to support environmental policies (European Union, 2007). Worldwide, the FAIR principles support a similar agenda to improve the findability, accessibility, interoperability, and reuse of digital assets (Wilkinson et al. 2016). However, neither INSPIRE nor the FAIR principles do consider all thematic aspects related to data needs in MSP (MSP Data Study, 2016). In the Baltic Sea region, the HELCOM-VASAB MSP Working Group (HELCOM – Baltic Marine Environment Protection Commission and VASAB – Vision and Strategy around the

¹ <http://www.emodnet.eu/> (last visit 23.07.2019)

² https://ec.europa.eu/maritimeaffairs/policy_en (last visit 23.07.2019)

Baltic Sea³) is promoting advances in MSP data management (HELCOM-VASAB, 2018).

The importance of introducing data harmonisation measures to more efficiently communicate space and time as the spatial and temporal dimension in data is repeatedly discussed in the literature as a pressing need in MSP (Ehler, 2008; Hattam et al., 2015; MSP Data Study, 2016; Stamoulis & Delevaux, 2015). Spatial information is recognised as one of the most critical elements in decision making as it provides geographic context to planning and management (Strain et al., 2014). Marine ecosystems may change over time and thus the basis for compatibility of the sea uses. Therefore information on the temporal dimension is likewise important for the management of human activities at sea (Schaefer & Barale, 2011). The place-based and dynamic nature of ecosystems and the spatial and temporal nature of human activities raise a demand to manage marine areas in a way that includes both the three-dimensional aspects as well as the time-dependent perspective of data in the marine environment (Crowder & Norse, 2008; Ehler, 2008; Shucksmith & Kelly, 2014). Besides the two dimensional horizontal dimensions, it is increasingly relevant to take the vertical and temporal dimensions into account as MSP aims to avoid potential conflicts and foster synergies (HELCOM, 2010, 2016, 2017; Uusitalo et al., 2016).

Despite the afore mentioned directives and initiatives, the actual implementation of spatial and temporal aspects in datasets remains challenging (Shucksmith & Kelly, 2014) and coherent solutions are still lacking. Furthermore, even though it is commonly mentioned that MSP requires both spatial and non-spatial information (Shucksmith & Kelly, 2014), the data discussion in the literature is highly focused on spatial data, maps, and geospatial analysis. Non-spatial data may serve as important sources of evidence for spatial planning, but since the spatial representation is missing, these datasets remain unaccounted in MSP data repositories. Consequently, non-spatial data are often excluded also from discussions on data harmonisation.

For planners, evidence needs to be of a certain quality and reliability irrespective of the data source. Since often, data and information are not produced by the planning authority itself, it is crucial that metadata provide a clear and transparent description (MSP Data Study, 2016).

The concept presented here was developed in the BONUS BASMATI⁴ project. Data management within the project and preparing project results as input to MSP

³ <http://www.helcom.fi/helcom-at-work/groups/helcom-vasab-maritime-spatial-planning-workinggroup>
(last visit 23.07.2019)

⁴ <https://bonusbasmati.eu>

in the Baltic Sea region posed the following challenges: (I) harmonisation and integration of different dimensions of space and time to datasets in a meaningful way, (II) managing spatial data (GIS data) and other types of information that may have indirect spatial implications (e.g. socio-economic data and policies), and (III) ensuring data quality in a transnational environment (e.g. common language).

The aim of the current paper is to provide a conceptual approach to describe spatial and temporal data properties in MSP datasets and associated metadata, using a systematic categorisation and common wording. Second, the objective is to show how spatial and non-spatial information can be jointly managed in the same geographic data repository.

2. HETEROGENEITY OF MARINE DATA

2.1 Data types in marine spatial planning

Data in MSP cover a variety of topics as well as different phases of the planning cycle. Evidence is needed about the current state (stocktaking), future scenarios and visions, as well as policies and planning decisions (Ehler & Douvere, 2009).

The European Commission's technical study on 'Evaluation of data and knowledge gaps to implement MSP' (MSP Data Study 2016) identifies four broad types of data that have been used in existing plans and corresponding planning processes: (I) administrative boundaries, (II) data on the geophysical environment and biological/ecological features, (III) data relating to relevant human activities and sectors, and (IV) socio-economic and policy-related data. Most data available and accessible belong to data types (I) and (II). The amount of data belonging to type (IV) is currently increasing. Especially policy related data including data on human pressures for impact assessments will become more important in the future (HELCOM, 2016, 2017; Uusitalo et al., 2016). The majority of available data within all data types is descriptive and of applied evidence, meaning data gathered by measurements, sample analysis or models. Strategic evidence describing future scenarios or visions is still rare.

2.2. Diversity of spatial and temporal information

2.2.1 Spatial dimensions

The characteristics of spatial information in datasets can be described and documented in several ways. The geographic positions can be given as coordinates. The features can be presented as points, lines, or polygons. The spatial coverage can be addressed by the spatial scale or the resolution of a dataset (Lam & Quattrochi, 2018).

Spatial information for MSP purposes is in general embedded in the three-dimensional nature of the marine environment (Gilliland & Laffoley, 2008). Hence, the horizontal as well as the vertical dimension of a dataset need to be documented. The horizontal dimension is sometimes addressed by text attributes referring to the spatial coverage of the dataset (e.g. local, regional, or national). However, the vertical dimension is often not documented even though the data can be strongly associated with a specific water layer, such as the sea surface (e.g. ferry lines), or the seafloor (e.g. sea cables).

2.2.2 Temporal dimension

Time plays a central role in the compatibility of sea uses. The same area can host multiple uses either if they are compatible with each other or if they are occurring at different times. Whether activities take place once a year or are ongoing several hours each day will influence planning decisions.

Three aspects of 'time' may be differentiated within a dataset. (I) time of data collection or sampling; referring to a point in time and commonly expressed as date in a dataset. (II) time as a process step; referring to a time line (e.g. past, present, future). For instance, objects can be under construction or operational, permits can be active or already expired and datasets can include historical data as well as future scenarios. Similarly, marine spatial plans have a different status from preparation phase to full legal force. (III) time as a feature within the data itself referring to the occurrence and temporal frequency of activities. Regularly operating ferry lines are periodically occurring activities. Bird migration is a seasonal phenomenon and installation of new infrastructures is a single period.

2.2.3 Spatial and non-spatial data types

The majority of data used in MSP are spatial data as such. Still, there are many datasets which are not presented in spatial format but actually include spatial information. Many statistics useful for planning purposes are presented at a national or regional level without a reference to individual locations. Strategic documents are non-spatial evidence useful in spatial planning. Siting decisions related to marine activities may need information from economic baseline studies or impact studies (MSP Data Study, 2016). National laws and regulations, as well as international agreements and policies, are important sources of background information (Cornu et al., 2014) even though they may lack direct references to space.

As a forward-looking process, MSP manages activities and guides future development in a sea area (Schaefer & Barale, 2011). Consequently, future-oriented information is needed besides the stocktaking of data on current activities. Still, future scenarios such as climate change related information, trends

or forecasts cannot be mapped precisely. Information on future economic developments or impacts of technological and knowledge advances may be presented without spatial reference at all. Even many important MSP stakeholder groups do not have a clear vision about their future uses of marine space (Zaucha, 2012).

The social dimension (e.g. stakeholder values, cultural services) is essential in planning and management of public assets (Strickland-Munro et al. 2016; Chan et al. 2012). Socio-economic datasets can include, among others, statistics about economic indicators, traditional knowledge of local residents, or evidence on the willingness of communities to contribute towards conservation efforts. The great majority of the social indicators used in planning are not of spatial character (Cornu et al., 2014). Cultural values may have direct linkages to specific places, but often linkages may be indirect or even completely inappropriate (Gee et al., 2017).

3. METHODOLOGY AND RESULTS

We propose a concept to document information on the spatial and temporal dimensions of datasets adequately for MSP purposes. It is applicable to both the attribute tables or the metadata. The categorisation includes the vertical and the horizontal dimensions as well as the complexity of the temporal dimension. Furthermore, we propose a straightforward method to include and manage nonspatial datasets within a geographic information system (GIS).

3.1 Basis for the data concept

The concept builds on the data specification scheme by the HELCOM-VASAB MSP Working Group (HELCOM-VASAB, 2018) and develops it further. The data scheme introduces a set of technical requirements (data specifications) based on INSPIRE land use data specifications to facilitate the interoperability and harmonisation of spatial datasets. Predefined code list values and an inclusive list of attribute codes along with descriptions structure information on data features in detail (HELCOM-VASAB, 2018). (Figure 1).




Figure 1: Relation of HELCOM-VASAB MSP output data specification (grey) and the proposed conceptual approach for a categorisation of spatial and temporal dimensions (blue). Code list values with respective attribute domains (named 'attribute codes' in HELCOM-VASAB, 2018) and descriptions are selective (only for illustration purposes).

Our approach drives this scheme further by including categories for the spatial and temporal dimensions of datasets. The categories are applied to code list values, attribute domains and additional descriptions. (Figure 1).

3.2 Categories for spatial and temporal information

We introduce two categories to describe the spatial dimension: the 'vertical dimension' and the 'horizontal dimension' (Table 1). The main goal of adding a category for the vertical dimension is to provide approximate background information on whether the objects of interest occur in the surface water layer, at the seafloor, or somewhere in between (Figure 2). The vertical dimension includes additional attribute domains to cover data themes not directly in contact with the

sea itself, but strongly related to it. This comprises topics related to the air column above the sea surface as well as activities along the coast or in the coastal region (Figure 1, 2; Table 1). The horizontal dimensions represent different spatial scales from local to international. We adopt a Baltic Sea perspective and therefore suggest attribute domains such as 'BalticWide' and 'EU' (Figure 1, 2; Table 2). However, attribute domains can be adjusted for other sea areas and regions around the world.

Table 1: Categorisation of the vertical and horizontal spatial dimensions with predefined attribute domains

spatial dimension	attribute domain	description
vertical	airColumn	air column above the sea, e.g. bird migration routes, scenery values
	surfaceWater	upper water column including water surface, no defined depth description
	waterColumn	whole water column, e.g. aquaculture
	bottomWater	near water body above the seafloor, no defined depth description
	seafloor	solid ground and sediment of the marine environment, e.g. seagrass, cables
	entireColumn	seafloor, water column, and the air above them, e.g. offshore wind power, bridges, other infrastructure
	coastalRegion	near shore, coastline and coastal area, where maritime induced activities and infrastructure dominate, e.g. lighthouses, hotels, diving schools
	unknown	the information is not available
horizontal	point	Point like features with specific coordinate information, e.g. ship wreck
	local	small spatial extent, a few km or km ² , e.g. protected area, dredging plume
	regional	spatial scale reflecting ecological, historical, political, climatic or morphological zone, e.g. estuarine
	national	administrative boundary, e.g. state borders, exclusive economic zone
	basinWide	spatial scale following morphological characteristics of sea basins, e.g. Bornholm Basin
	BalticWide	spatial scale reflecting the whole geographic extent of the Baltic Sea
	EU	European Union wide information, e.g. EU policies and agreements

Information on the temporal dimension is divided into three categories: ‘temporal occurrence’, ‘temporal frequency’, and ‘time line’ (Table 2). ‘Temporal occurrence’ reveals whether e.g. activities are completely absent, occur only once or more often, or are present all the time. ‘Temporal frequency’ provides further information on the frequency of the activities that have more or less regular occurrence patterns (Figure 3). ‘Time line’ gives an indication whether the datasets represent past, present, or future activities, scenarios or conditions.




Figure 2: Illustration of the vertical (black) and horizontal (red italic) dimensions of marine environments.

Table 2: Categorisation of the temporal dimension with predefined attribute domains

temporal dimension	attribute domain	description
temporal occurrence	absent	e.g. for species that are not present anymore in a region, or for not existing data on marine spatial plans
	oneTimeEvent	can be applied for disastrous events, e.g. oil leak
	irregular	e.g. activities that follow no pattern, such as dredging events, which depend on weather conditions, supply and demand and administrative authorisations
	regular	e.g. cargo shipping, dive sites

	static	permanent and ongoing status e.g. maritime infrastructures
	notApplicable	no characteristic temporal occurrence can be applied to the data in a meaningful way
	unknown	the information is not available
temporal frequency	daily	event or activity that occurs (almost) every day, e.g. ferry
	monthly	event or activity that occurs (approximately) once a month, e.g. service trips to wind park
	seasonal	event or activity that occurs during specific seasons, e.g. algae bloom, bird migration, blockade of a shipping route with floating ice in winter
	annual	event or activity that occurs once a year, annual updates of socioe-conomic statistics
	decadal	reoccurring events on a long, at times irregular frequency, e.g. fresh water inflows from the North Sea to the Baltic Sea, extreme weather events
	otherPeriod	to describe unique frequency patterns that differ strongly from the listed classes above
	notApplicable	no characteristic temporal frequency can be applied to the data in a meaningful way
	unknown	the information is not available
time line	past	e.g. historical data, data of outdated marine spatial plans, historic coastlines
	recent	data that are collected in the past but describe the most current situation, e.g. environmental data, model derived data
	present	data on existing uses or present state, e.g. legally adopted documents (INSPIRE guideline), mariculture facilities
	future	applicable for planned infrastructures, future climate predictions, e.g. planned wind park, sea level rise
	notApplicable	no characteristic time line can be applied to the data in a meaningful way




Figure 3: Illustration of 'temporal occurrence' and 'temporal frequency', representing information on time in a dataset.

3.3 Technical design

The technical design of the conceptual approach can be implemented in two ways: in the attribute table or in the corresponding metadata. Advantages, challenges and limits of transferring information either as attribute encoded in the data or as description in the metadata are given below (Table 3). In addition, an example for the use of an auxiliary file to include non-spatial data in a GIS repository is presented.

3.3.1 Attribute table

Information in the attribute table is linked directly to the spatial feature (line, point, or polygon) and appropriate columns are created for each spatial and temporal category. Respective attribute domains can be selected according to the description. The risk of losing information during data exchange is diminished to a minimum.

The attribute table of datasets is directly accessible for GIS applications and other automated, multifunctional tools (Table 3). The user can identify spatial and temporal overlap by queries. Analysis can focus, for example, on seasonal events such as bird migration or algae blooms. The 3-dimensionality of the marine realm can be addressed more efficiently when information on the vertical and temporal dimension are directly accessible for data analysis.

Implementation in the attribute table raises two technical challenges. First, the file size of a dataset increases, depending on the amount of information included. Even though automated tools and systems can process a large amount of data, file size tends to increase the processing time in general. It can further hamper data

exchange between users due to the volume limits of exchange-tools. Second, for each spatial and temporal category only a single attribute domain can be selected. There might be datasets which are difficult to categorise using a single attribute domain. However, a combination of multiple choices in the attribute table hinders proper work flow with the data. Moreover, it may lead to conflicts and misinterpretation between users when data are exchanged. Automated analysing algorithms may not be able to handle multiple combinations of attribute domains, leading to false results. A more detailed description within the metadata is hereby mandatory.

3.3.2 *Metadata*

Similar to the attribute table, information can be stored in the metadata using the same categorisation. Contrary to the limited options of expressing spatial and temporal dimensions in the attribute table, metadata descriptions allow running text to categorise, describe, and define data in more detail (Table 3). Besides choosing from the same predefined attribute domains, additional text phrases, tags, and keywords can be noted. Multiple attribute domains can be applied to describe more complex data types, where the selection of a single one is not appropriate. When information is placed in the metadata, the file size of the original dataset does not increase. Therefore, processing time for data analysis is not affected by the amount of metadata.

Data queries and automated processing with analysis tools or GIS applications is not yet possible with information stored in the metadata. Metadata is stored in a separate file alongside with the dataset and needs to be transferred in parallel with each data exchange. The risk of losing this information during data exchange is not neglectable even for advanced practitioners. During data exchange, both systems (consignor and recipient) need to have the same technical and software standards to process data and metadata at the same high-quality level. Institutions with different technical capacities, both transnationally and nationally, need to overcome such obstacles.

Table 3: Overview on advantages (+) and limits (-) of transferring information (I) as attribute encoded in the data and (II) as description in the metadata.

Attribute encoded	Metadata
<ul style="list-style-type: none"> + Information is included in dataset (risk of loss during data exchange diminishes) + Accessible for data analysis + Access with multifunctional tools 	<ul style="list-style-type: none"> + File size does not increase (process time not affected) + Multiple attribute domains can be applied + All additional information for the dataset in the metadata (additional text phrases, tags and keywords)
<ul style="list-style-type: none"> - Increasing file size (processing time might be affected) - No multiple choices in attribute domains - Additional information (detailed description, tags, keywords, etc.) needs to be placed separately in the metadata 	<ul style="list-style-type: none"> - No automated access for multifunctional tools - Separate file (risk of loss during data exchange) - Same technical system and software standards necessary to share and process data

3.4 Inclusion of non-spatial data

Non-spatial data can be given a suggestive spatial frame. By referring to an auxiliary spatial data layer, such as a shapefile in vector format, non-spatial data can be included in a geographic database. The shapefile is delineated by the target area of the information in the non-spatial dataset. The actual dataset can be uploaded to the database in its original format and attached to the auxiliary file. Alternatively, the information can be linked to its original source if it is available online. The method allows including further information on the dataset the same way as for spatial datasets, for example on the spatial and temporal dimensions of the data.

The status of MSP processes in the Baltic Sea countries is used as an illustrative example for including non-spatial data in a GIS-application. Here, national waters are chosen as target areas for the spatial auxiliary file (Figure 4). Information about the planning area and the status of planning may as such be included in the metadata or in the attribute table (Table 4). Links to the official planning documents and further information not directly relevant for data queries may be included in the metadata alone (Table 4, grey background). Additional columns can be implemented to specify more detailed information about the dataset.




Figure 4: The status of marine spatial planning in the Baltic Sea EU countries as an example of implementing non-spatial data in a GIS-application using an auxiliary file.

Table 4: Information for an attribute table and/ or metadata (grey background) of an auxiliary file that visualises non-spatial data. The example corresponds to Figure 4, representing the status of marine spatial plans in the EU.

Planning area	Status	Link to original data	Further information
Denmark	Not available		www.msp-platform.eu/countries/denmark
Estonia Hiiumaa Island	Pilot Plan		https://www.msp-platform.eu/countries/estonia
Estonia Parnu Bay area	Pilot plan		https://www.msp-platform.eu/countries/estonia

Estonia	In progress		https://www.msp-platform.eu/countries/estonia
Finland	In progress	https://www.merialuesuunnittelu.fi/en/	https://www.msp-platform.eu/countries/finland
Finland Kymenlaakso Region	Existing plan		https://www.msp-platform.eu/countries/finland
Germany EEZ	Existing Plan	https://www.bsh.de/EN/TOPICS/Offshore/Maritime_spatial_planning/maritime_spatial_planning_node.html	www.msp-platform.eu/countries/germany
Germany MV	Existing Plan	https://www.bsh.de/EN/TOPICS/Offshore/Maritime_spatial_planning/maritime_spatial_planning_node.html	www.msp-platform.eu/countries/germany
Germany SH	Existing Plan	https://www.bsh.de/EN/TOPICS/Offshore/Maritime_spatial_planning/maritime_spatial_planning_node.html	www.msp-platform.eu/countries/germany
Latvia	In progress		https://www.msp-platform.eu/countries/latvia
Lithuania	Existing plan	https://www.mspplatform.eu/practices/supplement-lithuanian-masterplan-marine-spatialolutions	https://www.msp-platform.eu/countries/lithuania
Poland	In progress		https://www.msp-platform.eu/countries/poland
Sweden	In Progress	https://www.havochvatten.se/en/swam/eu-international/marinespatial-planning/consultation.html	www.msp-platform.eu/countries/sweden

4. DISCUSSION

Applying the data harmonisation concept holds several benefits for different user groups. The accessibility, understandability and transparency of different data types in MSP will improve in different aspects (Table 5).

4.1 Potential user benefits

Besides the complexity and heterogeneity of datasets utilised in MSP, various user groups (e.g. planners, stakeholders, technology) have different requirements and needs for datasets related to the planning processes.

Planners need information, not only to understand the object of planning and the developments there, but also to ensure proper communication and stakeholder involvement throughout the planning process (Zaucha, 2012). For stakeholder integration, it is important to pay attention to the language and terminology used in presenting the information. The presentation format affects how easily spatial and temporal information is understood by users (Shucksmith & Kelly, 2014). Stakeholders with varying backgrounds may perceive maps, visualisation methods, and choices of words differently, which in turn may cause unnecessary misconceptions or biases interpreting the evidence. Here, it is crucial to understand both the spatial and temporal coverage of the information.

The proposed concept provides a basis to document and compare information using a common language. Simple wording and intuitive categories may reduce the potential of misinterpretation and enable easy access and understanding for planners, stakeholders or other users. Maximal benefit from the categorisation is gained when all datasets are structured the same way. However, adapting old datasets to the proposed model may be challenging.

Providing information on the vertical and temporal dimension besides the horizontal extent, helps to decide whether activities can occur simultaneously (without affecting each other). It is highly relevant for both the planners and the stakeholders to fully consider the influence that activities are causing on the environment and on other sea use sectors (HELCOM, 2016, 2017). Highlighting the versatility in dimensions in human uses may also help to identify opportunities for co-location. Also here, planners and stakeholders could benefit from more detailed information on spatial and temporal dimensions in datasets.

While in general, planners are interested in incorporating social data in planning evidence, data availability and integration of social and biophysical data can be challenging (Cornu et al., 2014). The visibility and availability of inherently nonspatial information in traditionally spatial domains of MSP data can be enhanced by storing the knowledge in a common database. The suggested auxiliary file method presented in this paper is straightforward and easily adaptable. Comparing different data types is additionally improved by the use of a common categorisation with predefined attribute domains.

The application of Spatial Decision Support Systems (SDSS), multi-functional tools, interactive mapping, and automated analysing instruments are standard procedures within MSP processes (Fiduccia et al., 2016; Dapuzo et al., 2015; Greco et al., 2018; Janssen et al., 2015; Pinarbaşı et al., 2017). In general, applied technical devices do not set requirements on data e.g. categorisation, relations, or other specifications. Nonetheless, for robust analyses, all data need to be compatible. The proposed categorisation including the predefined attribute domains meets the need for harmonious wording and facilitates the access with automated tools. The spatial and temporal categorisation can support transparent visualisation at the user interface, facilitate navigation and data queries, as well as enhance data organisation and management. Data on the temporal dimension can potentially enable data exploration methods such as the animation of seasonal variations in data.

The proposed concept meets the fundamental FAIR data principles: Findability Accessibility, Interoperability and Reusability (Wilkinson et al. 2016) for good data management for users as well as automated instruments by harmonious wording and terminology, transparent and intuitively structured categorisation and widespread descriptions of information.

4.2 Applicability of spatial and temporal categorisation to MSP data types

4.2.1 Administrative boundaries

In general, 'administrative boundaries' represent a rather static type of data, although there are a general trend towards larger administrative units. Additional information on the spatial and temporal dimension of the datasets provides little practical benefits (Table 5).

Administrative boundaries do not have vertical limitations, they usually cover the whole column of the marine environment, from the depths of the seafloor to the sea surface and the air column above it. Therefore, a more detailed description of the vertical dimension is rarely applicable. The horizontal dimension on the contrary may add useful information, e.g. when searching for information on a specific administrative level, such as municipal boundaries on a local scale.

Information on the temporal dimension can be useful but also redundant within 'administrative boundaries' datasets. State borders or economic zones represent static information where temporal occurrence and temporal frequency are not applicable. Changes may occur in the time line as information represents either past, present, or future situations.

4.2.2 Environmental data

'Environmental data' (including model data) is usually based on field measurements or observations. This means that data are derived at specific coordinates, describing a temporary situation, such as current direction or intensity at a certain location.

For some environmental data, the vertical component is specified by providing the depth of measurement: e.g. methane concentration in 35 metres depth below sea surface. However, without information on the total water depth at the sampling location, such data does not imply whether the sample originates from close to the seafloor or rather from the topmost water layer (in relative terms). Thus, a categorisation of the vertical dimension can help users to assess whether the data is suitable for their purposes.

Information about the temporal dimension is beneficial with environmental datasets. Recurring events such as bird migration, algae blooms, tidal activities, and weather conditions can be described more accurately when adding information on occurrence and frequency. Datasets of modelled environmental conditions may reconstruct the past or predict the future which can be referred to in the category 'time line'.

4.2.3 Human activities

The data type 'human activities' seems to be categorised most naturally into the proposed spatial and temporal attribute domains. Human activities have pronounced spatial and temporal aspects.

Leisure boating or surfing affects merely the surface water layer, whereas installed pipelines or cables have an impact on the seafloor. Wind energy infrastructure, sediment dredging, or oil ricks, on the contrary, are examples of activities that have an effect on the entire water column. Maritime transport activities can include onetime voyages or operate on a regular basis. There are many activities that are strongly seasonal, such as recreational activities along the coastline. The time line can include information on past activities as well as predictions and scenarios on future developments.

Overall, a detailed description of space and time for human activity data can facilitate the identification of synergies and conflicts of these. The detailed categorisation meets the demands of more transparency and better description of spatial and temporal aspects for impact and pressure assessments (HELCOM, 2010, 2016, 2017; Uusitalo et al., 2016).

4.2.4 Socio-economic and policy-related data

With spatial auxiliary files, non-spatial (socio-economic and policy-related) information can be handled as spatial data, which allows all information to be accessible in one place. Moreover, it allows using a coherent categorisation on space and time in the same way as for naturally spatial datasets.

In regard to the spatial dimension, e.g. national laws or sector development plans can be linked to an auxiliary file (shapefile) with the national borders as target areas. Socio-economic data, such as the willingness of residents to contribute to a nature protection scheme (e.g. acquired via interviews or workshops), may address very local settings. In these cases, a shapefile can include the area to be protected as a polygon feature or locations of individual events as point features.

Considering the vertical dimension, the attribute domain 'coastalRegion' can be selected for activities that are not operated directly in water. Such activities may relate to statistics of overnight stays in the area. Evidence on cultural values can be assigned to the above sea surface environment (attributes domains 'airColumn', 'coastalRegion') if it considers scenery values or to 'seafloor' if it lists underwater cultural heritage sites. Offshore wind park developments have multiple vertical impact layers. The above sea surface parts of the constructions can have an effect on migratory birds or landscape values, and at the same time, the submarine parts affect seafloor habitats or influence local currents. Here, the attribute domain 'entireColumn' is appropriate.

Similar to 'administrative data', 'socio-economic and policy data' can be rather static in nature and a detailed description of temporal occurrence and frequency is not applicable. However, it can be distinguished whether the data represents past, present, or future conditions. The majority of data represents either existing activities or the most recent, up-to-date information available. The future is represented in scenarios of climate change impacts and economic developments and the past in historical datasets and outdated documents, where newer versions are already available.

Table 5: Overview of strengths and limits of the data harmonisation concept focusing on the perspective of users and addressing different data types used in MSP processes.

	Strengths	Limits
Planners, stakeholders,	+ structural basis to document and compare information	- categorisation approach requires training of planners, users, etc.

other users, technology	+ usage of common language, simple wording	- integration of older data is challenging: data with no categorisation needs modulation (time consuming)
	+ fast access	
	+ facilitates the identification of synergies and conflicts	
	+ transparent information for impact assessment of activities	
	+ incorporation of non-spatial data possible	
	+ application in Spatial Decision Support Systems	
	+ facilitation of navigation, organisation, and data exploration	
Administrative boundaries	+ description of horizontal dimension possible	- static data type, categorisation provides little practical benefit
	+ description of time line possible	
Environmental data	+ model derived data allow visualisation of past and future condition	- most data describe particular situation at specific location at certain time in the past
	+ compilation of data on reoccurring environmental phenomena	
Human activities	+ facilitates identification of synergies and conflicts of simultaneously occurring activities	
	+ compilation of data on reoccurring activities	
	+ transparent information for impact assessment of activities	
Socioeconomic and policy-related data	+ information accessible in GISsystem	- static data type, categorization provides little practical benefit
	+ inclusion of information of coastal region in MSP	
	+ description of time line possible	

4.3 Data quality control

A highly challenging aspect in data management, in general and in working with a categorisation system such as presented here, is the quality control. Data portals should have quality control systems as incoming dataset and metadata sometimes have errors despite common guidelines and data quality requirements (Underwood et al., 2018). This includes aspects such as correct and harmonised classification, regular updates, and accessible storage of information.

To maximise the benefits of data categorisation, it must be ensured that attribute domains are used in a harmonious way. Harmonisation is especially important in enabling transnational data exchange processes (Abramic et al., 2018; Jay et al., 2016). Even though the scope and level of detail in data may be more simple in cross-border planning, the coherence and harmonisation of the datasets remains a major task, which begins with a challenge of finding a common language and consistent vocabulary (MSP Data Study, 2016). The predefined attribute domains along with simple descriptions presented here can help to reduce false or ambiguous categorisation.

Datasets and corresponding metadata need updates. A national strategy document may represent 'present' information at the time of uploading. However, the information becomes 'past' or outdated whenever a new updated version of the document is published. In such cases, the old document can be stored in the database as a historical reference, which in some cases may be appropriate, but the time line category of the evidence ought to be changed irrespective of the fact whether a new document is available in the portal or not.

One option to maintain high quality data and to ensure the suitability of datasets for different user needs is to store information within the dataset itself (in the attribute table) as well as in the metadata. As attribute domains in the attribute table are rather short, information in the metadata can give more details and additional explanation. However, providing data and metadata should be easy and intuitive in order to make data sharing attractive. If data and metadata descriptions are too complicated and time-consuming to produce (Kalantari et al., 2014; Olfat et al., 2011) data providers may lose interest in data exchange. Consequently, effort should be paid to avoid or lower any hindrances to the users to provide and share their own data and information in MSP processes.

5 CONCLUSION

The presented conceptual approach categorises the spatial and temporal dimensions in a meaningful way for MSP purposes. It facilitates analysis of potential areas of conflict or for co-location of activities. The simple and intuitive language in predefined attribute domains can contribute to data harmonisation in cross-border planning. To ensure high quality of datasets, the concept may be implemented in both the attribute table and the metadata depending on the needs of the user. The inclusion of non-spatial data within GIS repositories ensures the availability of all relevant data in one database. This may foster a more holistic approach to planning. Overall, the concept provides effective steps towards a more coherent data management in line with the FAIR principles and subsequently better use of information in MSP processes.

6 ACKNOWLEDGEMENTS

The present work has been carried out within the project BONUS BASMATI (Baltic Sea Maritime Spatial Planning for Sustainable Ecosystem Services), which has received funding from BONUS (art. 185), funded jointly by the EU, Innovation Fund Denmark, Swedish Research Council Formas, Academy of Finland, Latvian Ministry of Education and Science and Research centre Jülich, Germany. Part of this work was funded by the research and development project MSP-TRANS (FKZ 3517 84 0100) of the Federal Agency for Nature Conservation, Germany and funded by the Federal Ministry for the Environment, Nature Conservation and Nuclear Safety, Germany. We thank Lotta Maack for GIS expertise and her contributions to the figures. Also we would like to thank Andrea Morf and Christian Koski for their contributions regarding user perspectives.

7 LITERATURE

- Abramic A. et al. (2018) Maritime spatial planning supported by infrastructure for spatial information in Europe (INSPIRE) *Ocean & Coastal Management* 152,23-36. <https://doi.org/10.1016/j.ocecoaman.2017.11.007>
- Bartha, G. & Kocsis, S. (2011). Standardization of geographic data: The european inspire directive. *European Journal of Geography*, 2(2), 79–89.

- Chan, K. M. A. et al. (2012). Where are Cultural and Social in Ecosystem Services? A Framework for Constructive Engagement. *BioScience*, 62(8), 744–756. <https://doi.org/10.1525/bio.2012.62.8.7>
- Cornu, E. L. . et al. (2014). Current Practice and Future Prospects for Social Data in Coastal and Ocean Planning. *Conservation Practice and Policy*, 28(4), 902–911. <https://doi.org/10.1111/cobi.12310>
- Crowder, L., & Norse, E. (2008). Essential ecological insights for marine ecosystem-based management and marine spatial planning. *Marine Policy*, 32(5), 772–778. <https://doi.org/10.1016/J.MARPOL.2008.03.012>
- Dapueto, G. et al. (2015). A spatial multi-criteria evaluation for site selection of offshore marine fish farm in the Ligurian Sea, Italy. *Ocean & Coastal Management*, 116,64–77. <https://doi.org/10.1016/J.OCECOAMAN.2015.06.030>
- Ehler, C. (2008). Conclusions: Benefits, lessons learned, and future challenges of marine spatial planning. *Marine Policy*, 32(5), 840–843. <https://doi.org/10.1016/J.MARPOL.2008.03.014>
- Ehler, C., & Douvère, F. (2009). Marine Spatial Planning: a step-by-step approach toward ecosystem-based management. *Intergovernmental Oceanographic Commission and Man and the Biosphere Programme, IOC Manuals and Guides* 53, 1–98. <http://dx.doi.org/10.25607/OBP-43>
- European Union. (2007). Directive 2007/2/EC of the European Parliament and of the Council of 14 March 2007 establishing an Infrastructure for Spatial Information in the European Community (INSPIRE). Official Journal of the European Union, Vol. 50, L 108, ISSN 1725-2555.
- Fiduccia, A., et al. (2016). Web-Based Spatial Decision Support Systems to Monitor and Manage Coastal Environments. 18th Italian National Conference on Photonic Technologies (Fotoniica 2016) p60(4.) – 60(4.)
- Gee, K. et al. (2017). Identifying culturally significant areas for marine spatial planning. *Ocean & Coastal Management*, 136, 139–147. <https://doi.org/10.1016/j.ocecoaman.2016.11.026>
- Gilliland, P. M., & Laffoley, D. (2008). Key elements and steps in the process of developing ecosystem-based marine spatial planning. *Marine Policy*, 32(5), 787–796. <https://econpapers.repec.org/RePEc:eee:marpol:v:32:y:2008:i:5:p:787-796>
- Greco, M. et al. (2018). Integrated SDSS for Environmental Risk Analysis in Sustainable Coastal Area Planning. *Computational Science and Its Applications - ICCSA 2018* (pp. 671–684). Cham: Springer International Publishing.

- Hattam, C. et al. (2015). Marine ecosystem services : Linking indicators to their classification. *Ecological Indicators*, 49, 61–75. <https://doi.org/10.1016/j.ecolind.2014.09.026>
- HELCOM. (2010). Towards a tool for quantifying anthropogenic pressures and potential impacts on the Baltic Sea marine environment: A background document on the method, data and testing of the Baltic Sea Pressure and Impact Indices. *Balt. Sea Environ. Proc. No. 125*, 1–72. <http://helcom.fi/Lists/Publications/BSEP125.pdf>
- HELCOM. (2016). Endorsment of the method to calculate the Baltic Sea cumulative impact index (BSII). *HELCOM TAPAS Project*, 1–17. [https://portal.helcom.fi/meetings/HOD51-2016-400/MeetingDocuments/67 Endorsement of the method to calculate the Baltic Sea cumulative impact index \(BSII\).pdf](https://portal.helcom.fi/meetings/HOD51-2016-400/MeetingDocuments/67%20Endorsement%20of%20the%20method%20to%20calculate%20the%20Baltic%20Sea%20cumulative%20impact%20index%20(BSII).pdf)
- HELCOM. (2017). Preliminary results of the Baltic Sea Impact Index Input. *Helcom Spice Workshop*, 1–47. [https://portal.helcom.fi/meetings/HELCOM SPICE BSII WS 1-2017-427/MeetingDocuments/Document 3_Preliminary results from the Baltic Sea Impact Index.pdf](https://portal.helcom.fi/meetings/HELCOM%20SPICE%20BSII%20WS%201-2017-427/MeetingDocuments/Document%203_Preliminary%20results%20from%20the%20Baltic%20Sea%20Impact%20Index.pdf)
- HELCOM-VASAB. (2018). Recommendations for transboundary MSP output data. 9th meeting of the BSR MSP Data Expert Subgroup 17-18 April Vilnius, Lithuania. [https://vasab.org/wp-content/uploads/2018/06/MSP-Output-Data recommendations_approved_byData_ESG_April2018.pdf](https://vasab.org/wp-content/uploads/2018/06/MSP-Output-Data%20recommendations%20approved%20byData_ESG_April2018.pdf)
- Janssen, R. et al. (2015). Decision support tools for collaborative marine spatial planning: identifying potential sites for tidal energy devices around the Mull of Kintyre, Scotland. *Journal of Environmental Planning and Management*, 58(4), 719–737. <https://doi.org/10.1080/09640568.2014.887561>
- Jay, S. et al. (2016) Transboundary dimensions for marine spatial planning: Fostering inter-jurisdictional relations and governance. *Marine Policy* 65,85-96. <https://doi.org/10.1016/j.marpol.2015.12.025>
- Kalantari, M. et al. (2014) Geospatial Metadata 2.0 - An approach for Volunteered Geographic Information. *Computers, Environment and Urban Systems* 48,35-48. <https://doi.org/10.1016/j.compenvurbsys.2014.06.005>
- Lam, N. S.-N., & Quattrochi, D. A. (2018). On the Issues of Scale, Resolution, and Fractal Analysis in the Mapping Sciences. *The Professional Geographer*, 44(1), 88–98. <https://doi.org/10.1111/j.0033-0124.1992.00088.x>
- MSP Data Study. (2016). Evaluation of data and knowledge gaps to implement MSP. Technical Study under the Assistance Mechanism for the implementation of Maritie Spatial Planning. *European Commission*, 136. <https://doi.org/10.2826/25289>

- Olfat, H. et al. (2011) A GML-based approach to automate spatial metadata updating *International Journal of Geographical Information Science* 231-250. <https://doi.org/10.1080/13658816.2012.678853>
- Pınarbaşı, K. et al. (2017). Decision support tools in marine spatial planning: Present applications, gaps and future perspectives. *Marine Policy*, 83, 83–91. <https://doi.org/10.1016/j.marpol.2017.05.031>
- Schaefer, N., & Barale, V. (2011). Maritime spatial planning: opportunities & challenges in the framework of the EU integrated maritime policy. *Journal of Coastal Conservation*, 15(2), 237–245. <https://doi.org/10.1007/s11852011-0154-3>
- Shucksmith, R. J., & Kelly, C. (2014). Data collection and mapping – Principles, processes and application in marine spatial planning. *Marine Policy*, 50 Part A, 27–33. <https://doi.org/10.1016/J.MARPOL.2014.05.006>
- Stamoulis, K. A., & Delevaux, J. M. S. (2015). Data requirements and tools to operationalize marine spatial planning in the United States. *Ocean & Coastal Management*, 116, 214–223. <https://doi.org/10.1016/j.ocecoaman.2015.07.011>
- Strain, L. et al. (2014). Marine administration and spatial data infrastructure. *Marine Policy*, 30(4), 431–441. <https://doi.org/10.1016/j.marpol.2005.03.005>
- Strickland-Munro, J. et al. (2016). Valuing the wild, remote and beautiful: Using public participation GIS to inform tourism planning in the Kimberley, Western Australia. *International Journal of Sustainable Development and Planning*, 11(3), 355–364. <https://doi.org/10.2495/SDP-V11-N3-355-364>
- Underwood E, et al. (2018) The use of biodiversity data in spatial planning and impact assessment in Europe. *Research Ideas and Outcomes* <https://doi.org/10.3897/rio.4.e28045>
- Uusitalo, L. et al. (2016). Exploring methods for predicting multiple pressures on ecosystem recovery: A case study on marine eutrophication and fisheries. *Continental Shelf Research*, 121, 48–60. <https://doi.org/10.1016/j.csr.2015.11.002>
- Wilkinson, M. D. et al. (2016). The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data* Article no. 160018. <https://doi.org/10.1038/sdata.2016.18>
- Zaucha, J. (2012). Offshore Spatial Information – Maritime Spatial Planning in Poland. *Regional Studies*, 46(4), 459–473. <https://doi.org/10.1080/00343404.2012.668615>