

California State University, Monterey Bay
Digital Commons @ CSUMB

School of Natural Sciences Faculty Publications
and Presentations

School of Natural Sciences

2013

Genetic Diversity, Population Structure, and Demographic History of Exploited Sea Urchin Populations (*Tripneustes Gratilla*) in the Philippines

Inggat Laya N. Casilagan

Marie Antonette Juinio-Menez

Eric D. Crandall

California State University, Monterey Bay, ecrandall@csumb.edu

Follow this and additional works at: https://digitalcommons.csumb.edu/sns_fac

Recommended Citation

Casilagan, I.L.N.*, Juinio-Meñez, M. A., and E.D. Crandall. 2013. Genetic diversity, population structure, and demographic history of exploited sea urchin populations (*Tripneustes gratilla*) in the Philippines. *Journal of Experimental Marine Biology and Ecology* 449: 284-293.

This Article is brought to you for free and open access by the School of Natural Sciences at Digital Commons @ CSUMB. It has been accepted for inclusion in School of Natural Sciences Faculty Publications and Presentations by an authorized administrator of Digital Commons @ CSUMB. For more information, please contact digitalcommons@csumb.edu.

1 **Genetic diversity, population structure, and demographic history of exploited sea urchin populations**
2 **(*Tripneustes gratilla*) in the Philippines**

3

4 Inggat Laya N. Casilagan ^{1,*}, Marie Antonette Juinio-Meñez ¹, and Eric D. Crandall ^{2,3}

5

6 ¹ The Marine Science Institute, College of Science, University of the Philippines

7 Velasquez St., Diliman, Quezon City 1101 Philippines

8 ² Southwest Fisheries Science Center, Fisheries Ecology Division, 110 Shaffer Rd. Santa Cruz, CA 95060, USA

9 ³ Department of Ocean Sciences, University of California, 110 Shaffer Rd. Santa Cruz, CA 95060, USA

10

11 *Corresponding Author:

12 Email: inggatlaya@upmsi.ph

13 Telephone: (632) 922 3921

14 Fax: (632) 924 7678

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31 **Abstract**

32

33 The sea urchin *Tripneustes gratilla* is ecologically and economically important in the Indo-Pacific
34 region. We use population genetic methods to investigate the population structure and historical demography
35 of exploited populations in the Philippines. Sea urchins sampled in 6 localities in western Luzon and 4 outgroup
36 sites were sequenced for mitochondrial cytochrome oxidase-1 gene (n = 282) and genotyped for seven
37 microsatellite loci (n = 277). No significant genetic structure was found for either class of markers, indicating
38 either extensive gene flow across the archipelago, or that populations have high genetic diversity and have not
39 yet attained equilibrium between genetic drift and migration following large changes in demography.
40 Interestingly, demographic inferences from the two types of markers were discordant. Mitochondrial lineages
41 showed demographic expansion during the Pleistocene while microsatellite data indicated population
42 decline. Estimates for the date of each event suggest that a Pleistocene expansion could have preceded a more
43 recent population decline, but we also discuss other hypotheses for the discordant inferences. The high genetic
44 diversity and broad distribution of haplotypes in populations that recently recovered from fishery collapse
45 indicate that this species is very resilient over evolutionary timescales.

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61 1. Introduction

62 The widely-distributed *Tripneustes gratilla* is a high-value sea urchin in the Indo-Pacific region. This
63 ubiquitous herbivore occupies a key trophic position in nutrient cycling especially in tropical seagrass
64 ecosystems (Koike, et al. 1987; Klumpp et al., 1993; Alcoverro and Mariani, 2002). *T. gratilla*, being an
65 opportunistic grazer, also functions as a keystone species for coral reef and sea grass communities by controlling
66 invasive macroalgae (Conklin and Smith, 2005; Stimson et al., 2007). Population outbreaks in some areas,
67 however, have resulted in overgrazing of seagrasses (Eklöf et al., 2008) and foliose algae (Valentine and Edgar,
68 2010). This is also an economically-valuable sea urchin species and primarily collected for its high quality
69 gonad – a specialty food item primarily in Japan (Lawrence and Agatsuma, 2001; Andrew et al., 2002). Many
70 fisheries for this species have declined over time due to indiscriminate harvesting and lack of management
71 measures which resulted to economic losses (i.e. Shimabukuro, 1991; Talaue-McManus and Kesner, 1995;
72 Andrew et al., 2002). While these cases documented significant fisheries impacts on *T. gratilla* populations,
73 they need to be placed into a broader spatial and temporal context. What does the regional population structure
74 of the species look like? What is the demographic history of the species?

75 A population genetic approach can potentially provide these much needed insights for exploited
76 invertebrate species (Thorpe et al., 2000). Molecular methods were initially used with marine species to better
77 understand the effects of larval dispersal on the population structure (e.g. Waples 1987; Doherty et al., 1995)
78 and determine the spatial scales of population connectivity (Palumbi, 2003; 2004; Hedgecock et al., 2007;
79 Cowen and Sponaugle, 2009). Development of genetic markers and analysis have since extended the utility of
80 genetic data to direct measurement of migration, estimation of effective population size, and examination of
81 population demographic history (Emerson et al., 2001; Pearse and Crandall, 2004; Manel et al., 2005; Hellberg,
82 2009; Hare et al., 2011). To date, genetic assessments on *T. gratilla* populations have been limited to broad scale
83 phylogeographic surveys of its genus (Lessios et al., 2003) and phylogenetic studies (Zigler and Lessios, 2003;
84 Palumbi and Lessios, 2005). Mitochondrial sequence variation showed very weak regional divergence of
85 *Tripneustes* populations across the Indo-Pacific region despite the significant local differentiation among the
86 populations in this region (Lessios et al., 2003). This implies that *T. gratilla* in this region belongs to a large
87 *Tripneustes* metapopulation (Lessios et al., 2003). The only finer-scale genetic survey on *T. gratilla* populations
88 was carried out in western Luzon and eastern Philippines (Malay et al., 2002). This was conducted soon after the
89 collapse of a local artisanal sea urchin fishery (i.e. Talaue-McManus and Kesner, 1995) to aid management of
90 the heavily-exploited *T. gratilla* populations in northwestern Luzon (Malay et al., 2002). Based on six allozyme

91 markers, genetic differentiation of sea urchin populations within the western Luzon region was not significant,
92 indicating extensive gene flow among populations (Malay et al., 2002). This lack of genetic structure of *T.*
93 *gratilla* was initially explained by its potential for long-distance larval dispersal (Malay et al., 2002). This
94 species has a relatively long planktonic larval duration of 20 to 52 days under culture conditions (Shimabukuro,
95 1991; Juinio-Meñez et al., 1998; Lawrence and Agatsuma, 2001). The planktonic stage or echinopluteus, though
96 capable of movement using their ciliated bands (Emlet et al., 2006), is categorized as a weakly swimming larva
97 and considered passive to the forces of oceanographic processes (Chia et al., 1984 cited in Weersing and
98 Toonen, 2009). Thus, there is great potential for *T. gratilla* larvae to be dispersed over large distances before
99 settlement. It is possible, though, that limited but significant genetic differentiation was not detected with the
100 employed genetic marker, sampling design, or analysis (Ward, 2006). The availability of DNA markers for *T.*
101 *gratilla* and recent developments in population genetics analysis presents an opportunity to re-examine the
102 genetic variation in this exploited species.

103 Using multiple molecular markers, this study aims to obtain a more detailed characterization of *T.*
104 *gratilla* populations in the Philippines to make inferences about population structure, effective size, and
105 demographic history and gain insights on the vulnerability of the species to exploitation. Mitochondrial DNA
106 has been noted for its relatively rapid development of population genetic structure due to its small effective
107 population size. Based on this, it has become a preferred genetic marker over allozymes for initial examination
108 of population differentiation (Bowen et al., in press). Given its non-recombining nature, mtDNA has been useful
109 in gaining some insights on species evolutionary history and population demographic history based on maternal
110 lineage (Avisé et al., 1987; Liu and Cordes, 2004 but also see Ballard and Whitlock, 2004). On the other hand,
111 microsatellites are inherited as codominant markers and thus provide insights on genetic differentiation of
112 populations based on gene flow of both sexes (Liu and Cordes, 2004; Selkoe and Toonen, 2006). Highly
113 polymorphic microsatellite loci also have the potential to reveal contemporary gene flow and effective
114 population size (Ovenden et al., 2007; Saenz-Agudelo et al., 2011). The different characteristics of the DNA
115 markers and new analytical approaches employed in this study should aid in elucidating historical and
116 contemporary processes that influenced the current pattern of genetic variations in *T. gratilla*.

117

118 2. Materials and methods

119

120 2.1. Tissue collection and DNA extraction

121 Samples were collected from 6 sites along the western Luzon coast and 4 outgroup sites from other
122 regions in the archipelago (Table 1). Only adult sea urchins with test diameter greater than 60 mm (Bangi, 2001)
123 were collected on snorkel and tube feet and spines were sampled non-destructively. Individual tissue samples
124 were kept in a vial and preserved in 90% ethanol. DNA was extracted from the tissue samples using a 10 %
125 ChelexTM (Biorad) solution (Walsh et al., 1991).

126

127 2.2. DNA sequencing of the CO1 region

128 A region of the mitochondrial cytochrome oxidase subunit 1(CO1) gene was amplified using the
129 universal forward primer: CO1f 5' CCTGCAGGAGGAGGAGAYCC and a *Tripneustes*-specific reverse primer
130 CO1TR1 5'GGCATTCCAGCTAGTCCTARAAA (Lessios et al., 2003). PCR reactions were carried out in a
131 final volume of 25- μ l containing 1 μ l genomic DNA extraction, 2.5 μ l 10x PCR Buffer (PE-II), 2.5 μ l of 8 mM
132 dNTPs, 2.0 μ l of 25 mM MgCl₂, 1.25 μ l of each primer (10 mM), 0.125 μ l of Amplitaq Gold polymerase
133 (Applied Biosystems; 5 units/ μ l) and 14.5 μ l of molecular grade water. The PCR temperature profile was as
134 follows: initial denaturation for 7 min at 95°C, 40 times of the main cycle: 30 s at 95°C (denature), 30 s at 52°C
135 (anneal) and 1 min at 72°C (extension), and 10 min of final elongation at 72°C. After amplification, PCR
136 products were run in 1% agarose gels stained with ethidium bromide to evaluate the quality and quantity of the
137 amplified DNA. Excess dNTPs and primers were removed prior to cycle sequencing using an enzymatic
138 method. Five (5) μ l of the PCR product was incubated with 0.5 μ l (0.5 U) of shrimp alkaline phosphatase and
139 0.5 μ l (5 U) of exonuclease (GE Healthcare) for 30 min at 37°C, 15 min at 80°C, and 1 min at 25°C. Purified
140 DNA products were cycle sequenced in both directions using BigDye 3.0 terminator chemistry (Applied
141 Biosystems, Foster City, CA). The cycle sequencing reaction was carried out in a 12- μ l volume reaction: 2.5 μ l
142 of ABI 5x sequencing buffer, 0.5 μ l of 10 mM primer, 0.5 μ l BigDye, 0.5 μ l DMSO, 7.0 μ l molecular grade
143 water and 1 μ l of cleaned PCR product. The sequencing program started with 25 cycles of 95°C for 10 s, 50°C
144 for 5 s and 60°C for 5 min and a final step of 5 min at 20°C (Barber et al., 2006). Afterwards, the labelled DNA
145 sequences were precipitated with ethanol, resuspended in formamide, and sent to Life Sciences Core
146 Laboratories Center at Cornell University for sequencing on an Automated 3730 DNA Analyzer (Applied
147 Biosystems, Inc.).

148

149 *2.3. Microsatellite PCR amplification and genotyping*

150 Seven out of 11 microsatellite loci developed for *T. gratilla* (Carlon and Lippé, 2007) were used
151 (Supplementary Table S1). The other four loci were excluded due to persistence of null alleles in all
152 subpopulations tested (Carlon and Lippé, 2007). PCR amplifications of three loci: Tgr-B11, Tgr-C117, and Tgr-
153 D134 were carried out in multiplex while the remaining four loci (i.e. Tgr-24, Tgr-A11, Tgr-C11, and Tgr-D5)
154 were individually amplified in separate PCR runs. The multiplex PCR consisted of 1 µl genomic DNA extracts,
155 1 µl 10x PCR buffer, 1 µl of 8 mM dNTPs, 1 µl of 1mg/ml Bovine Serum Albumin (BSA A7030 Sigma), 0.8 µl
156 of 25 mM MgCl₂, 0.05 µl of Amplitaq polymerase (5 units/µl), 2.15 µl of molecular grade water, and the
157 forward and reverse primers (10 mM) of each locus, 0.25 µl Tgr-B11, 1.0 µl Tgr-C117, and 0.25 µl Tgr-D134
158 adding to a final volume of 10 µl.

159 For single-locus PCR, the 10-µl reactions contained 4.15 µl of molecular grade water, 0.5 µl of the
160 forward and reverse primers (10 mM) of a specific locus with same volume of the remaining reagents as
161 described above. The temperature profile of the PCR was the following: an initial denaturation at 94°C for 7 min
162 followed by several cycles (38-40 times) of denaturing at 94°C for 30 s, annealing at T_A for 30 s (Supplementary
163 Table S1), and elongation of 5 min at 72°C. PCR products were then electrophoresed in 1% agarose gels,
164 stained with ethidium bromide, and examined under UV light.

165 For genotyping, microsatellite loci were divided into two pooling sets of PCR products. One pooling
166 set was a mixture of 0.5 µl PCR products of locus Tgr-24 and 1.0 µl of the multiplex PCR products. The other
167 set was comprised of 1.0 µl PCR products of locus Tgr-C11 and 0.5 µl PCR products of each locus, Tgr-D5 and
168 Tgr-A11. In each well of a 96-well plate, a set of pooled PCR products were combined with 9 µl Formamide-
169 GeneScan™ 500 LIZ® (Applied Biosystems) mixture. The master mix solution was initially prepared in a
170 microcentrifuge tube by combining 985 µl of formamide and 15 µl of GeneScan™ 500 LIZ®. The latter served
171 as internal molecular weight standard for allele calling. The PCR products were sent to Life Sciences Core
172 Laboratories Center at Cornell University and analyzed on Automated 3730xl DNA Analyzer (Applied
173 Biosystems, Inc.) with a fluorescent-based detection system.

174

175 *2.4 Sequence data analysis*

176 Electropherograms were visualized, proofread and compiled in Sequencher v4.8 (GeneCode, Ann
177 Arbor, MI) and the resulting sequences were aligned in MUSCLE (Edgar, 2004). Sequence data was collapsed

178 into unique haplotypes using the web tool, FaBOX (Villesen, 2007). A haplotype network was created in TCS
179 v1.21 (Clement et al., 2000) using a statistical parsimony procedure. A minimum spanning tree based on
180 pairwise differences was also generated in Arlequin v3.5 (Excoffier and Lischer, 2010) and visualized in
181 FigTree (Rambaut, 2009). To illustrate the relationships between unique haplotypes, the minimum spanning
182 haplotype tree was edited to reflect key results from TCS v1.21 (Clement et al., 2000). The dominant haplotypes
183 were also compared with sequences deposited in Genbank with BLASTN v2.2.24+ (Zhang et al., 2000) at the
184 nucleotide collection (nr/nt) database. A number of sequences from different localities were downloaded and a
185 neighbor-joining tree was generated using MEGA4 (Tamura et al., 2007) under nucleotide models of Log-Det
186 and Kimura 2-parameter (following Lessios et al., 2003) with 1000 bootstrap replicates.

187 Genetic diversity indexes were calculated for each population with Arlequin v3.5 (Excoffier and
188 Lischer, 2010): number of haplotypes (Nh), number of polymorphic sites (Np), haplotype diversity (h), and
189 nucleotide diversity (π). The same program was used to examine genetic differentiation among populations
190 using an analysis of molecular variance (AMOVA) based on haplotype frequency and sequence divergence (F_{ST}
191 and Φ_{ST} , respectively). Pairwise genetic distances (F_{ST} and Φ_{ST}) between populations were also calculated and
192 significance values were determined by performing 10000 permutations among the individuals between
193 population.

194 The relationship between genetic distance and geographical distance among populations was examined
195 using IBDWS v3.16 (Jensen et al., 2005). Mantel tests were performed with 10,000 permutations on genetic
196 distance obtained from Arlequin v3.5 and the measured geographical distances between localities determined
197 from Google Earth® as the shortest distance over the sea.

198 The sequence data were further analyzed for information on changes in the long-term effective
199 population size of *T. gratilla*. Fu's F_S (1997) generated in Arlequin v3.5 was examined to test for significant
200 departures from the neutral model. These analyses were performed for each population as well as for the pooled
201 Philippines sequences. In addition, a Bayesian skyline plot was generated in BEAST v1.5.3 (Drummond et al.,
202 2005) to estimate past population dynamics over time back to the most recent common ancestor of the gene
203 sequences. The program utilizes a standard Markov chain Monte Carlo (MCMC) sampling procedure in
204 estimating a posterior distribution of effective population size through time from sequence data given any
205 specified nucleotide-substitution model (Drummond et al., 2005). Due to computational constraints for a large
206 dataset, a subsample of 100 sequences randomly chosen from unstructured populations was analyzed on
207 processors provided by the Cornell Computational Biology Service Unit web-computing facility

208 (<http://cbsuapps.tc.cornell.edu/beast.aspx>). The subsampled dataset was run 6 times for 40 million steps to
209 ensure convergence under an HKY+G model of nucleotide substitution, strict clock model (fixed clock rate of
210 1.0), and linear skyline model (skyline groups = 5) with the default priors for model parameters and statistics as
211 specified in BEAUti (κ : Gamma prior [0.05, 40], initial = 1.0; α : Uniform [0, 1 000], initial = 0.5;
212 skyline.popSize: Uniform [0, ∞], initial = 0.0030). The mutation model for this sequence data was determined
213 through jModelTest v.0.1.1 (Posada, 2008) implementing Bayesian Information Criterion selection strategies.
214 Logfiles and treefiles from the replicate runs were examined for convergence in Tracer v1.5 and were combined
215 together using LogCombiner. The skyline plot was created in Tracer v1.5 from the combined treefiles and
216 logfiles (Rambaut and Drummond, 2009). The estimates from BEAST were converted into units of time and
217 effective population size using a lineage mutation rate of 1.4%, 1.75%, and 2.67% per million years (MY) as a
218 heuristic values based from studies on *Tripneustes* (Lessios et al. 2003), *Echinometra* (McCartney et al., 2000)
219 and *Protoreaster nodosus* (Crandall et al., 2012), respectively.

220

221 2.5. Microsatellite analysis

222 Chromatograms were examined in STRand v2.3 (Toonen and Hughes, 2001) to determine the fragment
223 sizes (alleles) of each locus per sample. All samples were checked manually and samples with ambiguous peaks
224 were not scored. Descriptive statistics including number of alleles, allelic richness (standardized to the smallest
225 sample size, n=29), observed (H_o) and expected (H_e) heterozygosity (Nei, 1978) were determined for each
226 population at each locus using FSTAT v. 2.9.3 (Goudet, 1995). For each locus and each sampled population, the
227 inbreeding coefficient (F_{IS}) was calculated and significance was estimated with 840 randomizations as
228 implemented in the same program. Genotypic linkage disequilibrium (LD) of all locus pairs and deviations from
229 Hardy-Weinberg equilibrium (HWE) of each locus in every population were tested using the web version of
230 Genepop v4.0.10 (Rousset, 2008). For the evaluation of HWE, data for each locus were initially analyzed by
231 testing for the general probability of departure from HWE. When a significant departure was found, the more
232 explicit hypotheses of heterozygote excess ($H_e < H_o$) and heterozygote deficit ($H_e > H_o$) were tested. In all tests,
233 parameters were set at 1000 dememorizations, 1000 batches, and 10000 iterations per batch. The classical one-
234 stage corrections for the false discovery rate (FDR) method (Verhoeven et al., 2005; Pike, 2010) were applied to
235 the p-values obtained in tests with multiple comparisons (i.e. HWE, LD, heterozygote deficits, and inbreeding
236 coefficient) to generate q-values. Micro-Checker v2.2.3 (Van Oosterhout and Hutchinson, 2004) was also used
237 to identify possible genotyping errors due to null alleles, large allele dropout or mis-scoring of stutter peaks.

238 Results of analysis in Micro-checker indicated the possible presence of null alleles in all populations at locus
239 Tgr-24, Tgr-A11, Tgr-B11; 4 populations at locus Tgr-D5; and 2 populations at locus Tgr-C11 and Tgr-C117.
240 Among these loci, Tgr-24 was the most likely to have null alleles based on the highest frequency of PCR failure,
241 significant departure from HWE in all populations, and highest level of inbreeding coefficient. Tgr-24 is thus
242 considered a deviant locus in this study and subsequent analyses were carried out without this locus.

243 Genetic differentiation among the sampled sea urchin populations was initially examined by testing the
244 significance of genotypic differentiation for all populations and all pairs of populations in Genepop v4.0.10 on
245 the web (1000 dememorizations, 1000 batches, and 10000 iterations per batch). In Arlequin v3.5, genetic
246 differentiation of populations was evaluated under the assumptions of unstructured AMOVA (or single group of
247 populations) and infinite allele model. Because high levels of heterozygosity can lower the maximum value of
248 F_{ST} , we also measured G'_{ST} and D_{EST} , which correct for this problem using SMOGD (Crawford, 2010). Another
249 test of population differentiation was carried out in BAPS v5 (Corander et al., 2008) which was based on a
250 Bayesian clustering method. This population mixture analysis requires input of the maximum number of
251 genetically diverged groups (K). In this analysis, clustering of groups of individuals was carried out using
252 several K values (1, 2, 3, 4, 5 and 6) and analyses were run three times for each K value. In addition we used the
253 Bayesian coalescent sampler Migrate v3.5.1 (Beerli and Palzewski, 2010) to test a model of genetic structure
254 ($k=2$, Guimaras vs. Northern Luzon sites) against no structure ($k=1$). We used a Brownian motion model of
255 mutation, and windowed, exponential priors on Θ (1×10^{-4} to 1×10^3) and m/μ (1×10^{-3} to 1×10^4). Each model
256 was run for 1 million steps, with 10,000 steps removed as burnin, and 10 heated chains ranging in temperature
257 from 1 to 1×10^5 . Model selection was based on Bayes Factors calculated from the marginal likelihood of each
258 model, as estimated from a Bezier approximation to thermodynamic integration over the heated chains (see
259 Beerli and Palzewski 2010, Crandall et al., 2012). Based on unbiased genetic distance (Nei, 1978), a UPGMA
260 (unweighted pair-group method using arithmetic averages) tree of sampled populations was generated (1000
261 bootstraps) in TFPGA v1.3 (Miller, 1997). A Mantel test was carried out in IBDWS v3.16 (Jensen et al., 2005)
262 to test for a positive relationship between geographical distance (measured as shortest distance by sea) and
263 pairwise genetic differentiation obtained from Arlequin v3.5 based on 10000 random permutations.

264 To estimate contemporary effective population size of the *T. gratilla* population in Northwest Luzon,
265 we employed a method that utilizes the mean squared correlation in allele frequencies (i.e. linkage
266 disequilibrium) implemented in the LDNe software (Waples, 2006; Waples and Do, 2008). We combined data
267 from all localities except for Guimaras and Lucero (which showed a small amount of non-significant structure

268 with other localities), and only included alleles with a frequency greater than 0.02, which Waples and Do (2010)
269 have shown to balance the precision provided by many alleles with the bias created by rare alleles. Confidence
270 intervals were determined with a one-delete jackknife over loci.

271 To infer past demographic events from the microsatellite data, Bottleneck v1.2.02 (Piry et al., 1999)
272 was used to determine whether the sampled populations have experienced a reduction in their effective
273 population size. In principle, a severe reduction of effective population size results in a progressive reduction of
274 number of alleles and heterozygosity at polymorphic loci. However, the number of alleles is reduced faster than
275 the heterozygosity (H_e); hence, a transient excess in H_e is expected to characterize a bottlenecked population. To
276 detect heterozygosity excess ($H_e > H_{eq}$), this program compares the expected heterozygosity (H_e) calculated from
277 allele frequency and expected equilibrium heterozygosity (H_{eq}) (i.e. no bottleneck) derived from the number of
278 alleles found in the samples. The comparison of these heterozygosity parameters (H_e and H_{eq}) is in the context
279 of Nei's gene diversities (1978). Thus, this heterozygosity excess ($H_e > H_{eq}$) should not be confused with excess
280 of heterozygotes ($H_e < H_o$) which compares the proportion of heterozygotes with expectations of Hardy-
281 Weinberg equilibrium. The calculations were performed using the infinite allele model. Since there were fewer
282 than 20 loci, Wilcoxon's test was used to determine the significance of the observed heterozygosity excess of
283 the population (Piry et al., 1999). Bottleneck analyses excluding one or two and all of the putative loci with null
284 alleles (i.e. Tgr-24, Tgr-A11, Tgr-B11) using the original and Brookfield corrected data were also carried out.

285

286 3. Results

287 3.1. Genetic diversity

288 Mitochondrial CO1 sequences were obtained from 282 individuals of *T. gratilla* sampled from 10 sites.
289 The 605-bp sequences were aligned without indels and collapsed into 79 unique haplotypes characterized by 69
290 polymorphic sites (Supplementary Table S2; GenBank accession numbers: **JX661089-JX661167**). Most of
291 these haplotypes occurred only once (57 singleton or 72% of the total unique haplotypes). Based on overall
292 frequency, the haplotypes found to be dominant were sequence 1 (34.8%), 10 (11.7%), and 3 (9.3%) while the
293 rest of the haplotypes occurred less often (<2.5%). The minimum spanning tree revealed three star-like
294 polytomies with the dominant haplotypes separated by only one nucleotide difference (Fig. 1). All haplotypes
295 were closely related as they differed by only 1 to 10 unique mutations. The most frequent and broadly
296 distributed haplotype 1 was also identified by TCS v1.21 (Clement et al., 2000) as the most probable ancestral
297 haplotype. A neighbor-joining tree generated from a number of sequences from Lessios et al. (2003) showed

298 that the dominant haplotypes in this present study were also shared with other populations from different parts
299 of the world (Fig. 2). Interestingly, the most dominant haplotype 1 was shared by *T. gratilla* and *Tripneustes*
300 *depressus* populations from several localities across the Indian and Pacific Oceans (Fig. 2). Haplotype 1 was
301 also the most common haplotype in the global phylogeographic survey comprising about 18% of the pooled
302 sample of *T. gratilla* and *T. depressus* (Lessios et al., 2003).

303 Geographical distribution and relative frequency of unique haplotypes are shown in Fig. 3. The three
304 dominant haplotypes comprised the major proportion of the samples in all sites except Burgos where haplotype
305 3 was not sampled. Minor haplotypes were also shared by 2 to 6 populations. Private or site-specific haplotypes
306 were present in each population but comprised a small proportion of the samples (range 9.5 to 23.3%). In each
307 sampled population, the number of haplotypes ranged from 12 to 20 and number of polymorphic sites ranged
308 from 12 to 24 (Table 2) indicating high degree of genetic diversity. Overall, the haplotype diversity was high
309 (mean $h = 0.8554 \pm 0.0592$) while nucleotide diversity was low (mean $\pi = 0.0031 \pm 0.0005$) (Table 2).

310 A total of 277 sea urchins from 6 localities were genotyped at seven microsatellite loci (Supplementary
311 Table S3) but a number of individuals (no more than 12 per locus) had missing data at some loci due to
312 technical causes (i.e. PCR failure, ambiguous peaks). There was no significant linkage disequilibrium after
313 adjustment for multiple comparisons indicating that none of the loci were physically linked. Overall, there were
314 153 alleles found in all loci ranging from 9 to 41 alleles per locus. When averaged across populations, the
315 number of alleles per locus ranged from 7 to 28 alleles (or 6 to 24 alleles in terms of allelic richness). There
316 were 27 private alleles occurring at low frequency (0.010 to 0.021) across the sampled localities. High genetic
317 diversity is indicated by the mean and total expected heterozygosities of each locus ranging from 0.658 to 0.952
318 and 0.663 to 0.952, respectively.

319 There were significant discrepancies in the observed and expected heterozygotes per locus averaged
320 across the populations, indicating a high level of deviation from Hardy-Weinberg equilibrium (HWE). Out of 42
321 single locus tests per population, 27 (64.3%) did not conform to the expectations of HWE (Probability test,
322 $p < 0.05$, Supplementary Table S3). Further evaluation of HWE showed that majority of the populations that were
323 not in equilibrium had deficits in the number of heterozygotes (34 out of 42 tests were significant) and none had
324 heterozygote excesses (data not shown). In all sampled populations, more than half of the microsatellite loci
325 used in the study had heterozygote deficiency (Supplementary Table S3). Locus Tgr-24 was the most notable for
326 its significant HWE departure and heterozygote deficiency in all populations. Likewise, Tgr-A11 and Tgr-B11
327 also exhibited high degree of HWE deviation. The remaining four loci had fewer departures from HWE and

328 were at equilibrium in at least three sites. Null alleles have been identified as the most probable technical cause
329 of observed heterozygote deficit for these microsatellite loci as also reported in the primer note (Carlson and
330 Lippé, 2007). As we found no evidence for nulls at 4 loci and estimates of F_{IS} were positive at all loci, the
331 presence of null alleles can only partly explain the HWE departures observed in *T. gratilla* populations and
332 other processes (e.g. selection, demographic effects) may still have significant role in reducing heterozygosity.

333

334 3.2. Population genetic structure

335 Analysis of molecular variance (AMOVA) based on mitochondrial sequence data found no significant
336 population differentiation among population within western Luzon ($F_{ST}=-0.004$, $p=0.696$ and $\Phi_{ST}=-0.004$,
337 $p=0.727$). Genetic differentiation among sea urchin populations sampled across the archipelago were also not
338 significant ($F_{ST}=-0.003$, $p=0.736$ or $\Phi_{ST}=-0.003$, $p=0.767$). Pairwise F_{ST} values were less than 0.02 and none
339 were significant (data not shown). The genetic and geographic distance were not positively correlated ($Z=-$
340 69.6316 , $r=-0.1081$, Mantel test, $p=0.7202$).

341 Similar results were found in the analysis of microsatellite data. The fixation index values were also
342 very low ($F_{ST}<0.01$) and genetic differentiation among populations were not significant (AMOVA, global
343 $F_{ST}=0.001$, $p=0.719$). Per-locus estimates for G'_{ST} and D_{EST} were all below 0.02 (Supplementary Tables S4 and
344 S5). The UPGMA dendrogram also showed that the clustering of *T. gratilla* populations did not conform to the
345 expected grouping based on geographical distance between locations (Supplementary Fig. S1). This is
346 consistent with a lack of correlation between genetic and geographic distance ($Z=1.4041$, $r=-0.3958$, Mantel
347 test, $p=0.8591$). For example, Lucero and Victory populations were not clustered together despite their
348 proximity of less than 10 km apart. Further analyses also showed that genotypic differentiation for all
349 populations and pair-wise population comparisons were not significant (Fisher's method: $\chi^2=6.072$, $p=0.965$).
350 The Bayesian approach employed in BAPS v5 was also not able to partition the sampling localities. A model
351 with a single population ($k=1$) had a higher marginal likelihood value ($\log ML=-8067.51$) than the rest of
352 population models tested with probability values ranging from -8224.19 to -8916.66. Similarly, the $k=1$ model
353 in Migrate had a much higher marginal likelihood (-132,118) than a $k=2$ model (-775,726), although both
354 models showed good convergence as evaluated by effective sample sizes greater than 400.

355

356

357

358 3.3. Demographic history inferences

359 Fu's F_S values were significantly negative (Table 2, $p=0.0001$) in all sites, indicating an excess of rare
360 haplotypes and the rejection of the null model of neutral evolution. Similarly, the Bayesian skyline plot revealed
361 population growth in local sea urchin populations (Fig. 4). The abrupt increase of effective population size was
362 within the range of 30000-55000 years ago (median time) assuming a lineage mutation rate maximum at 2.67%
363 and minimum at 1.4%, respectively. Long-term coalescent N_e was estimated to range between 0.6 and 13.5
364 million effective females depending on which mutation rate was used. LDNe estimated contemporary effective
365 size for Northwest Luzon to be 6535.5 from six microsatellite loci, with a lower bound of 448.4 and an infinite
366 upper bound. This infinite upper bound is consistent with the absence of significant linkage disequilibrium
367 found by Genepop (Waples and Do, 2010). Bottleneck analysis of the microsatellite data revealed an excess in
368 expected heterozygosity, which indicates a decline in the effective population size of *T. gratilla*. Expected
369 heterozygosity (H_e) was generally found to be higher than the expected equilibrium heterozygosity (H_{eq}) in 33
370 out of 42 comparisons (Table 3) and the multi-locus statistical test showed that the observed heterozygosity
371 excess was significant in all sampled populations (Wilcoxon test, $p=0.008-0.016$, Table 3). Analyses that
372 excluded Tgr-A11 or Tgr-B11, which were also suspected with null alleles, still showed significant
373 heterozygosity excess in all populations (data not shown). In addition, this result stayed highly significant when
374 genotype data were corrected for null-alleles (Brookfield, 1996) and with Tgr-24 re-added to the dataset
375 (Supplementary Table S6).

376

377 4. Discussion

378 4.1. Non-equilibrium populations and absence of genetic structure in the Philippines

379 Western Luzon populations of *T. gratilla* were found to be genetically homogeneous and not
380 significantly distinct from populations in other regions of the Philippines. These results from seven
381 microsatellite loci and a mitochondrial locus are consistent with the findings of the previous genetic survey that
382 used allozyme markers (Malay et al., 2002). The observed genetic homogeneity of *T. gratilla* populations along
383 the western Luzon coasts is consistent with its life history features that predict long-distance larval dispersal.
384 The low and non-significant F_{ST} values ($F_{ST} < 0.02$) obtained for this species indicate the persistence of enough
385 gene flow among the sampled populations of about 10 or more migrants per generation (Lowe and Allendorf,
386 2010). Based on a passive particle dispersal model in the Western Luzon region (Bernardo, 2011), the estimated
387 range of dispersal distance for *T. gratilla* larvae was about 116 to 1060 km after 52 days depending on source

388 location and season (L. Bernardo, unpubl data). Considering that *T. gratilla* is a year-round spawner (Tuason,
389 1980), larvae can be effectively supplied over a wide range of oceanographic conditions (e.g. Addison et al.,
390 2008), exposing them to current vectors that may move in opposite directions depending on the monsoon
391 seasons (e.g. Juinio-Meñez and Villanoy, 1994), thereby further enhancing the extent of larval exchange within
392 the region.

393 The lack of genetic structure across the archipelago, however, could not be entirely explained by
394 contemporary oceanographic processes. This is in contrast with the findings of other local genetic studies in the
395 Philippines on marine species with biphasic life-histories that revealed broad to fine-scale genetic structure
396 within the archipelago that conformed with current patterns or biogeographic divisions (e.g. *Linckia laevigata* -
397 Juinio-Meñez et al., 2003; *Siganus fuscescens* - Magsino and Juinio-Meñez, 2008; Ravago-Gotanco and Juinio-
398 Meñez, 2010). Perhaps a better explanation for the absence of detectable genetic structure in our data is that the
399 underlying structure has been obscured by non-equilibrium processes such as the demographic changes
400 described below (Excoffier et al., 2009; Marko and Hart, 2011). Based on the large effective population size
401 estimated from microsatellites for the Northwest Luzon population ($N_e = 6535.5$), and assuming an ecologically
402 high rate of gene flow ($m = 0.1$), we estimate that it would take ~ 4700 generations for F_{ST} values to move
403 halfway to their equilibrium value following any of these demographic changes (Crow and Aoki 1984). The
404 inferred absence of equilibrium between gene flow and genetic drift is initially evidenced by the lack of
405 significant correlation between geographic and genetic distance for both markers (Slatkin, 1993). This is further
406 supported by the observed sharing of the dominant mitochondrial haplotypes with samples from Reunion Island
407 in Western Indian Ocean and even with its congeneric species, *T. depressus* in the Galapagos Islands in the
408 eastern Pacific (Lessios et al., 2003). This lack of divergence across the Eastern Pacific Barrier and the
409 incomplete sorting of the mitochondrial lineage for the genus *Tripneustes* indicate the persistence of genetic
410 patterns shaped by evolutionary events in the past (Benzie, 1999; Lessios et al., 2003).

411 For the microsatellite data, significant departures from Hardy-Weinberg equilibrium due to
412 heterozygote deficiency also substantiate the non-equilibrium state of *T. gratilla* populations. Heterozygote
413 deficiency has been associated primarily with ecological processes such as inbreeding, recent admixture,
414 selection, or accumulation of genetically distinct cohorts (e.g. Watts et al., 1990; Addison and Hart, 2004; van
415 Oppen et al., 2008). Considering that populations in this study were genetically homogeneous and have high
416 inbreeding coefficients without linkage disequilibrium, the most probable explanation for the heterozygote
417 deficiency would be the significant genetic differentiation among cohorts or the temporal Wahlund effect (i.e.

418 Watts et al., 1990). This has been suggested to be a consequence of large variance in reproductive success or
419 differential survival during the planktonic larval stage or immediately after settlement at early benthic stage
420 (Watts et al., 1990; Hedgecock, 1994; Hedgecock et al., 2007; Hedgecock et al., 2011). This hypothesis, also
421 known as sweepstakes reproductive success, suggests that this large reproductive variance is due to asynchrony
422 between oceanographic and biological conditions that may influence larval development, dispersal, and
423 recruitment (Hedgecock, 1994; Hedgecock et al., 2007, Hedgecock et al., 2011). *T. gratilla* is predisposed to
424 large variance in reproductive or recruitment success based on its life history features (i.e. high fecundity,
425 broadcast spawning, long planktonic larval duration), patchy distribution, and dynamic habitat (Shimabukuro,
426 1991; Lawrence and Agatsuma, 2001). In support of this idea, preliminary genetic analysis of *T. gratilla* recruits
427 and adults sampled from local populations in Santiago Island, Bolinao indicate genetic variability among
428 cohorts (Casilagan, 2011). Overall, these non-equilibrium results at the archipelagic scale are in agreement with
429 the chaotic population structure and absence of equilibrium that has been found across the entire species range
430 (Lessios et al., 2003).

431

432 4.2. Discordance in demographic trends inferred by mtDNA and microsatellites

433 Estimates of long-term coalescent effective population size from the mtDNA data were orders of
434 magnitude larger than estimates of contemporary effective size based on linkage disequilibrium in the
435 microsatellite data. This is unsurprising, because the two markers and methods are estimating over very different
436 temporal and spatial scales. The mtDNA data integrates over the coalescent history of the genetic sample, and
437 therefore reflects the population size of the global *T. gratilla* population, which may include the entire Indo-
438 Pacific, as indicated by the vast ranges of shared haplotypes in Fig. 2 (Lessios et al., 2003). On the other hand,
439 linkage disequilibrium is a transient phenomenon that occurs among loci, and N_e estimates from the
440 microsatellite data reflect only the effective number of parents from which the current sample was drawn. Thus,
441 this estimate represents the effective size of the breeding population in Northwest Luzon in 2007-2008. The
442 linkage disequilibrium method has been shown to be robust to migration from outside the sampled region unless
443 the percentage of migrants from outside is greater than 5%-10% (Waples and England, 2011).

444 Similarly, demographic analyses of mitochondria and microsatellite data suggested contrasting
445 demographic events in the history of *T. gratilla* populations in the Philippines. The mitochondrial lineages
446 indicated population expansion while microsatellite data exhibited genetic signatures of a population decline. In
447 particular, the highly negative F_u 's F_s , star-like haplotype network, high haplotype diversity and low nucleotide

448 diversity provided initial evidence of demographic expansion in the mtDNA. As shown by the Bayesian skyline
449 plot, the initiation of the abrupt population growth occurred during Pleistocene. Demographic expansion during
450 the Pleistocene also been reported for other marine taxa in the Indo-Pacific region such as gastropods (Crandall
451 et al., 2008a), echinoderms (Crandall et al., 2008b; Kochzius et al., 2009), crustaceans (Benzie et al., 2002), and
452 fish (Rohfritsch and Borsa, 2005; Liu et al., 2007; Ravago-Gotanco and Juinio-Meñez, 2010). The Pleistocene
453 era was characterized by changing sea levels and temperatures due to glaciations and deglaciations (Roy et al.,
454 1996; Rohling et al., 1998; Siddall et al., 2003). Particularly in Southeast Asia, sea levels were reduced to more
455 than 120 m below the present levels (Voris, 2000) which would have exposed reef flats causing local extinction
456 especially of species in shallow habitats. Following the Last Glacial Maximum (~20000 years BP), flooding of
457 the shelves provided newly-available habitat for re-colonization and subsequent population expansion (e.g.
458 Crandall et al., 2008a, 2008b; Ravago-Gotanco and Juinio-Meñez, 2010). The populations of *T. gratilla*, being a
459 shallow-water echinoid, were likely been influenced by these historical events.

460 On the other hand, microsatellite data indicate demographic decline in *T. gratilla* populations.
461 Significant heterozygosity excess was observed in most loci in all populations, and indicates a deficiency in
462 alleles that characterizes bottlenecked populations. It might at first be thought that, because microsatellites
463 evolve relatively faster than the mitochondrial markers, the bottleneck might be of anthropogenic origin,
464 associated with the recent collapse in the *T. gratilla* fishery (Talaue-McManus and Kesner, 1995). However, the
465 method used here detects bottlenecks that happened around 0.2-4.0 N_e generations ago (Cornuet and Luikart,
466 1996). Based on our estimates of contemporary effective size, this means that the bottleneck could have
467 occurred between 90 to more than 26,000 generations ago or similar values in years ago, assuming 1 year
468 generation time (Bangi, 2001; Lawrence and Agatsuma, 2001). This estimate suggests that the bottleneck
469 detected by the microsatellites cannot be attributed to the most recent collapse of the *T. gratilla* fishery, which
470 occurred in the 1990's, but might be attributed to earlier undocumented anthropogenic pressure, or else to
471 environmental changes during the late Pleistocene.

472 The seemingly discordant inferences of a population expansion in mtDNA and a bottleneck in
473 microsatellites can be reconciled in a number of ways. First, microsatellites are better suited for detecting
474 population declines than they are for detecting expansions (Cornuet and Luikart, 1996), so both inferences may
475 be correct. Given the rough range of dates that we inferred for each type of marker (expansion at 30-55 kya in
476 mtDNA, contraction at 0.009 – 26 kya in microsatellites), it is probable that a population decline detected by the
477 microsatellites occurred after the expansion detected in the mtDNA. Second, due to lack of recombination, it is

478 also possible that the mitochondrial genome had undergone a selective sweep or introgression (Ballard and
479 Whitlock, 2004). The sharing of dominant haplotypes among Indo-Pacific *T. gratilla* and eastern Pacific *T.*
480 *depressus* suggests recent introgression among these species, especially since a recent global survey of
481 *Tripneustes* using microsatellite markers revealed that *T. gratilla* can be genetically distinguished from the *T.*
482 *depressus* at nuclear loci (Carlson, pers comm). Hence, the observed genetic signature of population growth could
483 also be a consequence of selective sweep following introgression. Finally, it is possible that departures from the
484 neutral model detected in both mtDNA sequence and microsatellites could be the result of purifying selection.
485 While the mtDNA and microsatellite variation that we measured is putatively neutral, selection could have
486 occurred on nearby linked genes (i.e. background selection). Background selection has been shown to produce
487 patterns similar to population growth in sequence data (Fu, 1997) as well as to cause a loss of rare alleles that
488 might indicate a bottleneck in the microsatellite data (Charlesworth et al., 1993). Discordant inferences between
489 mtDNA and microsatellites have been noticed in a number of other marine taxa as well, and likely have to do
490 with the large variance in genealogies that is possible in species with large effective population sizes (DiBattista
491 et al., 2012). To disentangle the signals of historical processes related to demographic fluctuation from those of
492 selective processes, it might be necessary to obtain sequence data from multiple nuclear regions (e.g. Calderón
493 et al., 2008)

494

495 4.3. Conservation implications

496 Conservation of *T. gratilla* as a species is a relevant concern because it is the top target echinoid
497 species for commercial harvesting in the tropical and subtropical region. In this context, identification of
498 evolutionary significant units (ESUs, Moritz, 1994) and evaluating species vulnerability would be important.
499 The genetic analysis did not reveal distinct ESUs or indications of cryptic speciation based on the absence of
500 deep phylogenetic divergence. Based on the species' wide distribution range and the estimates of population
501 genetic diversity and variability (i.e. high genetic diversity, broad distribution of major haplotypes, and
502 extensive genetic exchange), the vulnerability of the species to extinction is low. Our data, however, also
503 suggest that populations in the Philippines have undergone large demographic fluctuations in the past, perhaps
504 similar to the anthropogenic changes that we are seeing now. Although recent recovery of overharvested
505 populations was observed (Juinio-Meñez et al., 2008), the persistent harvesting and habitat degradation still pose
506 a serious threat of local depletion. With the absence of genetic structuring particularly in western Luzon region,
507 a precautionary approach to management (e.g. McCook et al., 2009) should be practiced with a degree of

508 “spatial bet-hedging” (sensu Larson and Julian, 1999). This would entail region-wide management interventions
509 to protect adequate spawning stocks and ensure reliable recruitment in the localities. This can be carried out, for
510 example, by establishing network of marine protected areas and sea urchin grow-out cage culture sites along the
511 western Luzon coasts which serves as reproductive reserves and recruitment sites (Malay et al., 2002; Junio-
512 Meñez et al., 2008; 2009). Consideration of genetic impacts is also still vital especially in implementing the
513 culture-based management interventions such as the release or grow-out culture of hatchery-produced juveniles
514 (Ward, 2006). As with all fisheries species, hatchery-based supportive breeding efforts should aim to maximize
515 genetic diversity in the captive population released into the wild (Ryman and Laikre 1991). Integration of these
516 insights with other information (i.e. recruitment patterns, demographic data) would facilitate the development of
517 an effective management scheme that would ensure sustainability of the *T. gratilla* fishery.

518

519

520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549

Acknowledgments

This research study was part of the Coral Triangle - Partnerships for International Research and Education (CT-PIRE; OISE-0730256) Project of National Science Foundation (NSF) and we would like to thank K. Carpenter and P. Barber for their support. We also extend our gratitude to H. Bangi, W. Licuanan, M. Samson, and M. F. Nievaes for the logistical support and J. Evangelio, R. Palla, U. R. Casilagan, M. A. Paña, C. M. Edullantes, G. Peralta, J. Garcia, R. Dumalan, C. Ragos, T. Catbagan and R. Albino for the assistance in the tissue sampling and sea urchin surveys. We are also grateful to R. J. Ravago-Gotanco, M. C. Lagman, D. Carlon, A. Ackiss, C. Starger, E. Sabrocco, S. Ibarra, A. Lluisma and G. Concepcion for the general assistance in the laboratory work and analysis. We also thank the two anonymous reviewers for their valuable insights that help improve the manuscript. This study is part of I.L. Casilagan Master's thesis and was also supported by scholarship grant from Department of Science and Technology through the Accelerated Science and Technology Human Resources Development Program (DOST-ASTHRDP) and Marine Environment and Resources Foundation (MERF), Inc. This is UPMSI Contribution no. XXXX

550

551

552 **References**

553 Addison, J.A., Hart, M.W., 2004. Analysis of population genetic structure of the green sea urchin
554 (*Strongylocentrotus droebachiensis*) using microsatellites. *Mar. Biol.* 144, 243–251.

555 Addison, J.A., Ort, B.S., Mesa, K.A., Pogson, G.H., 2008. Range-wide genetic homogeneity in the California
556 sea mussel (*Mytilus californianus*): a comparison of allozymes, nuclear DNA markers, and mitochondrial
557 DNA sequences. *Mol. Ecol.* 17, 4222–4232.

558 Alcoverro, T., Mariani, S., 2002. Effects of sea urchin grazing on seagrass (*Thalassodendron ciliatum*) beds of a
559 Kenyan lagoon. *Mar. Ecol. Prog. Ser.* 226, 255–263.

560 Andrew, N., Agatsuma, Y., Ballesteros, E., Bazhin, A., Creaser, E., Barnes, D., Botsford, L.W., Bradbury, A.,
561 Campbell, A., Dixon, J., Einarsson, S., Gerring, P., Hebert, K., Hunter, M., Hur, S., Johnson, C., Juinio-
562 Meñez, M.A., Kalvass, P., Miller, R., Moreno, C., Palleiro, J., Rivas, D., Robinson, S., Schroeter, S.C.,
563 Steneck, R.S., Vadas, R., Woodby, D., Xiaoqi, Z., 2002. Status and management of world sea urchin
564 fisheries. *Oceanogr. Mar. Biol. Annu.* 40, 343–425.

565 Arnaud-Haond, S., Vonau, V., Rouxel, C., Bonhomme, F., Jean, P., Goyard, E., Boudry, P., 2008. Genetic
566 structure at different spatial scales in the pearl oyster (*Pinctada margaritifera cumingii*) in French
567 Polynesian lagoons: beware of sampling strategy and genetic patchiness. *Mar. Biol.* 155, 147–157.

568 Avise, J.C., Arnold, J., Ball, R.M., Bermingham, E., Lamb, T., Neigel, J.E., Reeb, C.A., Saunders, N.C., 1987.
569 Intraspecific Phylogeography: The population genetics and systematics. *Annu. Rev. Ecol. Syst.* 18, 489–
570 522.

571 Ballard, J.W.O., Whitlock, M.C., 2004. The incomplete natural history of mitochondria. *Mol. Ecol.* 13, 729 –
572 744.

573 Bangi, H.G.P., 2001. The effect of adult nutrition on somatic and gonadal growth, egg quality, and larval
574 development of the sea urchin *Tripneustes gratilla* (Echinodermata: Echinoidea). MS Thesis, University
575 of the Philippines – Diliman, 106pp.

576 Barber, P., Erdmann, M., 2000. Molecular systematics of the *Gonodactylidae* (Stomatopoda) using
577 mitochondrial cytochrome oxidase C (subunit I) DNA sequence data. *J. Crustacean Biol.* 20, 20–36.

578 Barber, P., Erdmann, M., Palumbi, S.R., 2006. Comparative phylogeography of three codistributed
579 stomatopods: origins and timing of regional lineage diversification in the coral triangle. *Evolution* 60,
580 1825–1839.

581 Beerli P, Palczewski M. 2010. Unified Framework to Evaluate Panmixia and Migration Direction Among
582 Multiple Sampling Locations. *Genetics* 185,313–326.

583 Benzie, J., Ballment, E., Forbes, A., Demetriades, N., Sugama, K., Moria, S., 2002. Mitochondrial DNA
584 variation in Indo-Pacific populations of the giant tiger prawn, *Penaeus monodon*. *Mol. Ecol.* 11, 2553–
585 2569.

586 Benzie, J.A.H., 1999. Genetic structure of coral reef organisms: ghosts of dispersal past. *Coral Reefs* 145, 131–
587 145.

588 Bernardo, L. P., 2011. Development of a particle dispersal model for Bohol Sea (Philippines). MS Thesis,
589 University of the Philippines – Diliman, 139pp.

- 590 Bowen, B.W., Shanker, K., Yasuda, N., Malay, M.C.D, von der Heyden, S., Paulay, G., Rocha, L.A., Selkoe,
591 K.A., Barber, P.H., Williams, S.T., Lessios, H.A., Crandall, E.D., Bernardi, G., Meyer, C.P., Carpenter,
592 K.E., and R.J. Toonen. Phylogeography Unplugged: Comparative geographic surveys in the genomic era.
593 In press at *Bulletin of Marine Science*.
- 594 Brookfield, J.F.Y., 1996. A simple new method for estimating null allele frequency from heterozygote
595 deficiency. *Mol. Ecol.* 5, 453-455.
- 596 Calderón, I., Giribet, G., Turon, X., 2008. Two markers and one history: phylogeography of the edible common
597 sea urchin *Paracentrotus lividus* in the Lusitanian region. *Mar. Biol.* 154, 137–151.
- 598 Carlon, D.B., Lippé, C., 2007. Eleven new microsatellite markers for the tropical sea urchin *Tripneustes gratilla*
599 and cross-amplification in *Tripneustes ventricosa*. *Mol. Ecol. Notes* 7, 1002–1004.
- 600 Casilagan, I.L.N., 2011. Spatial and temporal aspects of genetic variation in sea urchin, *Tripneustes gratilla*
601 populations in Western Luzon, Philippines. MS Thesis, University of the Philippines – Diliman, 95pp.
- 602 Charlesworth, B., Morgan, M., Charlesworth, D., 1993. The effect of deleterious mutations on neutral molecular
603 variation. *Genetics* 134, 1289–1303.
- 604 Chia, F. S., Buckland-Nicks, J., Young, C. M. 1984. Locomotion of marine invertebrate larvae: a review. *Can J*
605 *Zool*, 62(7), 1205-1222.
- 606 Clement, M., Posada, D., Crandall, K.A., 2000. TCS: a computer program to estimate gene genealogies. *Mol.*
607 *Ecol.* 9, 1657–1659.
- 608 Conklin, E.J., Smith, J.E., 2005. Abundance and spread of the invasive red algae, *Kappaphycus spp.*, in Kane
609 ohe Bay, Hawai'i and an experimental assessment of management options. *Biol. Invasions* 7, 1029–
610 1039.
- 611 Corander, J., Marttinen, P., Sirén, J., Tang, J., 2008. Enhanced Bayesian modelling in BAPS software for
612 learning genetic structures of populations. *BMC Bioinformatics* 9, 539.
- 613 Cowen, R.K., Sponaugle, S., 2009. Larval dispersal and marine population connectivity. *Annu. Rev. Mar. Sci.* 1,
614 443–466.
- 615 Crandall, E.D., Frey, M.A., Grosberg, R.K., Barber, P.H., 2008. Contrasting demographic history and
616 phylogeographical patterns in two Indo-Pacific gastropods. *Mol. Ecol.* 17, 611–26.
- 617 Crandall, E.D., Jones, M.E., Muñoz, M., Akinronbi, B., Erdmann, M. V., Barber, P.H., 2008. Comparative
618 phylogeography of two seastars and their ectosymbionts within the Coral Triangle. *Mol. Ecol.* 17, 5276–
619 5290.
- 620 Crandall, E.D., Sbrocco, E.J., Deboer, T.S., Barber, P.H., Carpenter, K.E., 2012. Expansion dating: calibrating
621 molecular clocks in marine species from expansions onto the Sunda Shelf following the Last Glacial
622 Maximum. *Mol. Biol. Evol.* 29, 707–19.
- 623 Crandall ED, Treml EA, Barber PH. 2012. Coalescent and biophysical models of stepping-stone gene flow in
624 neritid snails. *Mol Ecol* 21, 5579–5598.
- 625 Crawford NG. 2010. SMOGD: software for the measurement of genetic diversity. *Mol. Ecol. Resour.* 10, 556-
626 557.
- 627 Crow J.F., Aoki, K., 1984. Group selection for a polygenic behavioral trait: Estimating the degree of population
628 subdivision. *Proc. Natl Acad. Sci. USA* 81, 6073–6077.

- 629 De Loma, T.L., Conand, C., Harmelin-Vivien, M., Ballesteros, E., 2002. Food selectivity of *Tripneustes gratilla*
630 (L.)(Echinodermata: Echinoidea) in oligotrophic and nutrient-enriched coral reefs at La Reunion (Indian
631 Ocean). *B. Mar. Sci.* 70, 927–938.
- 632 Dibattista, J.D., Rocha, L.A., Craig, M.T., Feldheim, K.A., Bowen, B.W., 2012. Phylogeography of two closely
633 related Indo-Pacific butterflyfishes reveals divergent evolutionary histories and discordant results from
634 mtDNA and microsatellites. *J. Hered.* 103, 617–29.
- 635 Doherty, P., Planes, S., Mather, P., 1995. Gene flow and larval duration in seven species of fish from the Great
636 Barrier Reef. *Ecology* 76, 2373–2391.
- 637 Drummond, A.J., Rambaut, A., Shapiro, B., Obexer-Ruff, G., 2005. Bayesian coalescent inference of past
638 population dynamics from molecular sequences. *Mol. Biol. Evol.* 22, 1185–1192.
- 639 Edgar, R.C., 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic
640 Acids Res.* 32, 1792–7.
- 641 Eklöf, J.S., De la Torre-Castro, M., Gullström, M., Uku, J., Muthiga, N., Lyimo, T., Bandeira, S.O., 2008. Sea
642 urchin overgrazing of seagrasses: A review of current knowledge on causes, consequences, and
643 management. *Estuar. Coast. Shelf. S.* 79, 569–580.
- 644 Emerson, B.C., Paradis, E., Thébaud, C., 2001. Revealing the demographic histories of species using DNA
645 sequences distributions. *Trends Ecol. Evol.* 16, 707–716.
- 646 Emler, R.B., Young, C.M., George, S.B., 2006. Phylum Echinodermata: Echinoidea, in: Young, C.M., Sewell,
647 M.A., Rice, M.E. (Eds.), *Atlas of Marine Invertebrate Larvae*. Barcelona, pp. 531–551.
- 648 Excoffier, L., Foll, M., Petit, R.J., 2009. Genetic consequences of range expansions. *Annu. Rev. Ecol. Evol. S.*
649 40, 481–501.
- 650 Excoffier, L., Lischer, H.E.L., 2010. Arlequin suite ver 3.5: a new series of programs to perform population
651 genetics analyses under Linux and Windows. *Mol. Ecol. Resour.* 10, 564–567.
- 652 Fu, Y., 1997. Statistical tests of neutrality of mutations against population growth, hitchhiking and background
653 selection. *Genetics* 147, 915–925.
- 654 Hare, M. P., Nunney, L., Schwartz, M. K., Ruzzante, D. E., Burford, M., Waples, R. S., Ruegg, K., Palstra, F.
655 2011. Understanding and estimating effective population size for practical application in marine species
656 management. *Conserv Biol* 25, 438-449.
- 657 Hedgecock, D., 1994. Temporal and spatial genetic structure of marine animal populations in the California
658 Current. *California Cooperative Oceanic Fisheries Investigations Reports* 35, 73–81.
- 659 Hedgecock, D., Barber, P.H., Edmands, S., 2007. Genetic approaches to measuring connectivity. *Oceanography*
660 20, 70–79.
- 661 Hedgecock, D., Pudovkin, A.I. 2011. Sweepstakes reproductive success in highly fecund marine fish and
662 shellfish: a review and commentary. *Bull. Mar. Sci.* 87(4), 971-1002.
- 663 Hellberg, M.E., 2009. Gene flow and isolation among populations of marine animals. *Annu. Rev. Ecol. Syst.* 40,
664 291-310.
- 665 Jensen, J.L., Bohonak, A.J., Kelley, S.T., 2005. Isolation by distance, web service. *BMC Genet.* 6, 13.

- 666 Jones, G.P., Almany, G.R., Russ, G.R., Sale, P.F., Steneck, R.S., Van Oppen, M.J.H., Willis, B.L., 2009. Larval
667 retention and connectivity among populations of corals and reef fishes : history, advances and challenges.
668 Coral Reefs 28, 307–325.
- 669 Juinio-Meñez, M., Pastor, D., Bangi, H.G.P., 2009. Indications of Recruitment Enhancement in the Sea Urchin
670 *Tripneustes gratilla* Due to Stock Restoration Efforts, in: Proceedings of the 11th International Coral Reef
671 Symposium. Ft. Lauderdale, Florida, pp. 1017-1021.
- 672 Juinio-Meñez, M.A., Bangi, H.G.P., Malay, M.C.D., Pastor, D., 2008. Enhancing the recovery of depleted
673 *Tripneustes gratilla* stocks through grow-out culture and restocking. Rev. Fish. Sci. 16, 35–43.
- 674 Juinio-Meñez, M.A., Macawaris, N., Bangi, H.G.P., 1998. Community-based sea urchin (*Tripneustes gratilla*)
675 grow-out culture as a resource management tool, in: Jamieson, G.S., Campbell, A. (Eds.), Proceedings of
676 the North Pacific Symposium on Invertebrate Stock Assessment and Management. Can. Spec. Publ. Fish.
677 Aquat. Sci., pp. 393–399.
- 678 Juinio-Meñez, M.A., Magsino, R.M., Ravago-Gotanco, R.G., Yu, E.T., 2003. Genetic structure of *Linckia*
679 *laevigata* and *Tridacna crocea* populations among Palawan shelf & shoal reefs. Mar. Biol. 142, 717–726.
- 680 Juinio-Meñez, M.A., Villanoy, C.L., 1994. Sea urchin recruitment studies for management considerations, in:
681 Soemodihardjo, S., Tsukamoto, K., Romimoharto, K. (Eds.), Proceedings of the Joint Seminar on Marine
682 Science at the 4th Meeting of the LIPI-JSPS. Jakarta, Indonesia, pp. 15–20.
- 683 Kaplan, D.M., Planes, S., Fauvelot, C., Brochier, T., Lett, C., Bodin, N., Loc'h, F. Le, Tremblay, Y., Georges,
684 J.-Y., 2010. New tools for the spatial management of living marine resources. Current Opinion in
685 Environmental Sustainability 1, 1–6.
- 686 Klumpp, D.W., Salita-Espinosa, J.T., Fortes, M.D., 1993. Feeding ecology and trophic role of sea urchins in a
687 tropical seagrass community. Aquat. Bot. 45, 205–229.
- 688 Kochzius, M., Seidel, C., Hauschild, J., Kirchhoff, S., Mester, P., Meyer-Wachsmuth, I., Nuryanto, A., Timm,
689 J., 2009. Genetic population structures of the blue starfish *Linckia laevigata* and its gastropod ectoparasite
690 *Thyca crystallina*. Mar. Ecol. Prog. Ser. 396, 211–219.
- 691 Koike, I., Mukai, H., Nojima, S., 1987. The role of the sea urchin, *Tripneustes gratilla* (Linnaeus), in
692 decomposition and nutrient cycling in a tropical seagrass bed. Ecol. Res. 2, 19–29.
- 693 Larson, R.J., Julian, R., 1999. Spatial and temporal genetic patchiness in marine populations and their
694 implications for fisheries management. California Cooperative Oceanic Fisheries Investigations Report
695 40, 94–99.
- 696 Lawrence, J., Agatsuma, Y., 2001. The ecology of *Tripneustes*, in: Lawrence, J. (Ed.), Edible Sea Urchins:
697 Biology and Ecology. Elsevier Science, Amsterdam, pp. 395–413.
- 698 Lessios, H.A., Kane, J., Robertson, D.R., 2003. Phylogeography of the pantropical sea urchin *Tripneustes*:
699 Contrasting patterns of population structure between Oceans. Evolution 57, 2026–2036.
- 700 Levin, L.A., 2006. Recent progress in understanding larval dispersal: new directions and digressions. Integr. and
701 Comp. Biol. 46, 282–297.
- 702 Liu, J.-X., Gao, T.-X., Wu, S.-F., Zhang, Y.-P., 2007. Pleistocene isolation in the Northwestern Pacific marginal
703 seas and limited dispersal in a marine fish, *Chelon haematocheilus* (Temminck & Schlegel, 1845). Mol.
704 Ecol. 16, 275–88.
- 705 Liu, Z.J., Cordes, J.F., 2004. DNA marker technologies and their applications in aquaculture genetics.
706 Aquaculture 238, 1 – 37.

- 707 Lowe, W.H., Allendorf, F.W., 2010. What can genetics tell us about population connectivity? *Mol. Ecol.* 19,
708 3038–3051.
- 709 Magsino, R.M., Juinio-Meñez, M.A., 2008. The influence of contrasting life history traits and oceanic processes
710 on genetic structuring of rabbitWsh populations *Siganus argenteus* and *Siganus fuscescens* along the
711 eastern Philippine coasts. *Mar. Biol.* 154, 519–532.
- 712 Malay, M., Juinio-Meñez, M., Villanoy, C., 2002. Population genetic structure of the sea urchin *Tripneustes*
713 *gratilla* from selected sites in western Luzon and Eastern Philippines, in: Proceedings 9th International
714 Coral Reef Symposium. Bali, Indonesia, pp. 1–5.
- 715 Manel, S., Gaggiotti, O. E., Waples, R. S. 2005. Assignment methods: matching biological questions with
716 appropriate techniques. *Trends Ecol. Evol.* 20, 136–142.
- 717 Marko PB, Hart MW. 2011. The complex analytical landscape of gene flow inference. *Trends Ecol. Evol.* 26,
718 448–456.
- 719 McCartney, M., Keller, G., Lessios, H.A., 2000. Dispersal barriers in tropical oceans and speciation in Atlantic
720 and eastern Pacific sea urchins of the genus *Echinometra*. *Mol. Ecol.* 9, 1391–1400.
- 721 McCook, L.J., Almany, G.R., Berumen, M.L., Day, J.C., Green, a. L., Jones, G.P., Leis, J.M., Planes, S., Russ,
722 G.R., Sale, P.F., Thorrold, S.R., 2009. Management under uncertainty: guide-lines for incorporating
723 connectivity into the protection of coral reefs. *Coral Reefs* 28, 353–366.
- 724 Miller, M.P., 1997. Tools for Population Genetic Analyses (TFPGA). a Windows program for the analysis of
725 allozyme and molecular population genetic data. Ver. 1.3 Available via <http://bioweb.usu.edu/mpmbio>
- 726 Moritz, C., 1994. Defining 'evolutionarily significant units' for conservation. *Trends Ecol. Evol.* 9, 373–374.
- 727 Nei, M., 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals.
728 *Genetics* 89, 583–590.
- 729 Ovenden, J.R., Peel, D., Street, R., Courtney, A.J., Hoyle, S.D., Peel, S.L., Podlich, H., 2007. The genetic
730 effective and adult census size of an Australian population of tiger prawns (*Penaeus esculentus*). *Mol.*
731 *Ecol.* 16, 127–38.
- 732 Palsbøll, P.J., Bérubé, M., Allendorf, F.W., 2007. Identification of management units using population genetic
733 data. *Trends Ecol. Evol.* 22, 11–6.
- 734 Palumbi, S.R., 2003. Population genetics, demographic connectivity, and the design of marine reserves. *Ecol.*
735 *Appl.* 13, 146–158.
- 736 Palumbi, S.R., 2004. Marine reserves and ocean neighborhoods: The spatial scale of marine populations and
737 their management. *Annu. Reviews* 29, 31–68.
- 738 Palumbi, S.R., Lessios, H.A., 2005. Evolutionary animation : How do molecular phylogenies compare to Mayr's
739 reconstruction of speciation patterns in the sea? *P. Nat. Acad. Sci. USA* 102, 6566–6572.
- 740 Pearse, D.E., Crandall, K.A., 2004. Beyond FST: Analysis of population genetic data for conservation.
741 *Fisheries Sci.* 5, 585–602.
- 742 Pike, N., 2011. Using false discovery rates for multiple comparisons in ecology and evolution. *Methods Ecol.*
743 *Evol.* 2, 278–282.
- 744 Piry, S., Luikart, G., Cornuet, J.M., 1999. BOTTLENECK: A Computer Program for Detecting Recent
745 Reductions in the Effective Population Size Using Allele Frequency Data. *J. Hered.* 502–503.

- 746 Posada, D., 2008. jModelTest: phylogenetic model averaging. *Mol. Biol. Evol.* 25, 1253–1256.
- 747 Purcell, J.F.H., Cowen, R.K., Hughes, C.R., Williams, D.A., 2006. Weak genetic structure indicates strong
748 dispersal limits : a tale of two coral reef fish. *Proc. R. Soc. B.* 273, 1483–1490.
- 749 Rambaut, A. (2009) FigTree v1.3.1 <http://tree.bio.ed.ac.uk/software/figtree>
- 750 Rambaut, A., Drummond, A.J. (2009) Tracer v1.5. Available via <http://tree.bio.ed.ac.uk/software/tracer>
- 751 Ravago-Gotanco, R.G., Juinio-Meñez, M.A., 2010. Phylogeography of the mottled spinefoot *Siganus*
752 *fuscescens*: Pleistocene divergence and limited genetic connectivity across the Philippine archipelago.
753 *Mol. Ecol.* 1–15.
- 754 Reece, J.S., Bowen, B.W., Joshi, K., Goz, V., Larson, A., 2010. Phylogeography of two moray eels indicates
755 high dispersal throughout the indo-pacific. *J. Hered.* 101, 391–402.
- 756 Rohfritsch, A., Borsa, P., 2005. Genetic structure of Indian scad mackerel *Decapterus russelli*: Pleistocene
757 vicariance and secondary contact in the Central Indo-West Pacific Seas. *Heredity* 95, 315–26.
- 758 Rohling, E.J., Fenton, M., Jorissen, F.J., Bertrand, P., Ganssen, G., Caulet, J.P., 1998. Magnitudes of sea-level
759 lowstands of the past 500,000 years. *Nature* 394, 162–165.
- 760 Rousset, F., 2008. GENEPOP'007: a complete re-implementation of the GENEPOP software for Windows and
761 Linux. *Mol. Ecol. Resour.* 8, 103–106.
- 762 Roy, K., Valentine, J.W., Jablonski, D., Kidwell, S.M., 1996. Scales of climatic variability and time averaging
763 in Pleistocene biotas: implications for ecology and evolution. *Trends Ecol. Evol.* 11, 458–463.
- 764 Ryman, N., Laikre, L., 1991. Effects of supportive breeding on the genetically effective population size.
765 *Conserv. Biol.* 5, 325–329.
- 766 Saenz-Agudelo, P., Jones, G.P., Thorrold, S.R., Planes, S., 2011. Connectivity dominates larval replenishment
767 in a coastal reef fish metapopulation. *P. Roy. Soc. B.-Biol.* 278, 2954–61.
- 768 Schwartz, M.K., Luikart, G., Waples, R.S., 2006. Genetic monitoring as a promising tool for conservation and
769 management. *Trends Ecol. Evol.* 22, 25–33.
- 770 Selkoe, K.A., Toonen, R.J., 2006. Microsatellites for ecologists : a practical guide to using and evaluating
771 microsatellite markers. *Ecol. Lett.* 9, 615–629.
- 772 Shimabukuro, S., 1991. *Tripneustes gratilla* (Sea urchin), in: Shokita, S.K., Kakasu, K., Tomori, A., Toma, T.
773 (Eds.), *Aquaculture in Tropical Areas*. Midori Shobo Co. Ltd., Japan, pp. 313–318.
- 774 Siddall, M., Rohling, E., Almogi-Labin, A., Hemleben, C., Meischner, D., Schmelzer, I., Smeed, D.A., 2003.
775 Sea-level fluctuations during the last glacial cycle. *Nature* 423, 853–858.
- 776 Slatkin, M., 1993. Isolation by distance in equilibrium and non-equilibrium populations. *Evolution* 47, 264–279.
- 777 Stimson, J., Cunha, T., Philippoff, J., 2007. Food preferences and related behavior of the browsing sea urchin
778 *Tripneustes gratilla* (Linnaeus) and its potential for use as a biological control agent. *Mar. Biol.* 151,
779 1761–1772.
- 780 Talaue-McManus, L., Kesner, K., 1995. Valuation of a Philippine municipal sea urchin fishery and implications
781 of its collapse, in: *Philippine Coastal Resources Under Stress*. pp. 229–239.

- 782 Tamura, K., Dudley, J., Nei, M., Kumar, S., 2007. MEGA4: Molecular Evolutionary Genetics Analysis
783 (MEGA) software version 4.0. *Mol. Biol. Evol.* 24, 1596–9.
- 784 Thorpe, J.P., Sole-Cava, A.M, Watts, P. C., 2000. Exploited marine invertebrates: genetics and fisheries.
785 *Hydrobiologia* 420, 165-184.
- 786 Toonen, R.J., Hughes, S., 2001. Increased throughput for fragment analysis on ABI Prism 377 Automated
787 Sequencer using a membran comb and STRand software. *Biotechniques* 31, 1320–1324.
- 788 Tuason, A.Y., 1980. The reproductive biology of *Tripneustes gratilla* (Linnaeus) (Echinodermata: Echinoidea) –
789 with some notes on *Diadema setosum* (Leske). MS Thesis, University of the Philippines – Diliman, 77pp.
- 790 Vaitilingon, D., Rasolofonirina, R., Jangoux, M., 2003. Feeding preferences, seasonal gut repletion indices, and
791 diel feeding patterns of the sea urchin *Tripneustes gratilla* (Echinodermata: Echinoidea) on a coastal
792 habitat. *Mar. Biol.* 143, 451–458.
- 793 Valentine, J., Edgar, G., 2010. Impacts of a population outbreak of the urchin *Tripneustes gratilla* amongst Lord
794 Howe Island coral communities. *Coral Reefs* 29, 399–410.
- 795 Van Oosterhout, C., Hutchinson, W., 2004. MICRO-CHECKER: software for identifying and correcting
796 genotyping errors in microsatellite data. *Mol. Ecol.* 4, 535– 538.
- 797 Van Oppen, M.J.H., Lutz, A., De'ath, G., Peplow, L., Kininmonth, S., 2008. Genetic traces of recent long-
798 distance dispersal in a predominantly self-recruiting coral. *PlosOne* 3, e3401.
- 799 Verhoeven, K.J., Simonsen, K.L., & McIntyre, L.M. (2005). Implementing false discovery rate control:
800 increasing your power. *Oikos* 108, 643-647.
- 801 Villesen, P., 2007. FaBox: an online toolbox for fasta sequences. *Mol. Ecol. Notes* 7, 965–968.
- 802 Voris, H.K., 2000. Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time
803 durations. *J. Biogeogr.* 27, 1153–1167.
- 804 Walsh, P.S., Metzger, D.A., Higuchi, R., 1991. Chelex 100 as a medium for simple extraction of DNA for PCR-
805 based typing from forensic material. *Biotechniques* 10, 506.
- 806 Waples, R.S., 1987. A multispecies approach to the analysis of gene flow in marine shore fishes. *Evolution* 41,
807 385–400.
- 808 Waples, R.S., 2006. A bias correction for estimates of effective population size based on linkage disequilibrium
809 at unlinked gene loci*. *Conserv. Genet.* 7, 167–184.
- 810 Waples, R.S., Do, C., 2008. Ldne: a Program for Estimating Effective Population Size From Data on Linkage
811 Disequilibrium. *Mol. Ecol. Resour.* 8, 753–756.
- 812 Waples, R.S., England, P.R., 2011. Estimating contemporary effective population size on the basis of linkage
813 disequilibrium in the face of migration. *Genetics* 189, 633–44.
- 814 Waples, R.S., Punt, A.E., Cope, J.M., 2008. Integrating genetic data into management of marine resources: how
815 can we do it better? *Fish Fish.* 9, 423–449.
- 816 Ward, R.D., 2006. The importance of identifying spatial population structure in restocking and stock
817 enhancement programmes. *Fish.Res.* 80, 9–18.
- 818 Watts, R.J., Johnson, M.S., Black, R., 1990. Effects of recruitment on genetic patchiness in the urchin
819 *Echinometra mathaei* in Western Australia. *Mar. Biol.* 105, 145–151.

820 Weersing, K., Toonen, R.J., 2009. Population genetics, larval dispersal, and connectivity in marine systems.
821 Mar. Ecol. Prog. Ser. 393, 1–12.

822 Zhang, Z., Schwartz, S., Wagner, L., Miller, W., 2000. A greedy algorithm for aligning DNA sequences. J.
823 Comput. Biol. 7, 203–214.

824 Zigler, K.S., Lessios, H., 2003. 250 Million Years of Bindin Evolution. Biol. Bull. 205, 8–15.

825

826

827

828

829 **Fig. 1** Minimum spanning tree for the CO1 haplotypes of *T. gratilla*. Filled circle represents a unique haplotype
830 which was sized proportionally to its absolute frequency, ranging from 1– 98. The dominant haplotypes were
831 identified by their sequence number. The asterisk denotes the most probable root haplotype while the square
832 represents unsampled haplotypes as revealed by TCS v1.21 (Clement et al. 2000).

833

834 **Fig. 2** Neighbor-joining tree illustrating the relationship of the major CO1 haplotypes in this study and the CO1
835 sequences from Lessios et al. (2003) based on LogDet substitution model. Numbers next to nodes indicate
836 bootstrap support from 1000 iterations and nodes with less than 50% support have been collapsed.

837

838 **Fig. 3** Relative frequency of mitochondrial haplotypes per population. Represented in the pie graph are the
839 major haplotypes (sequence 1, 10, and 3), minor haplotypes (pooled haplotypes with frequency of 2 to 7), and
840 private haplotypes (haplotypes found in only one site).

841

842 **Fig. 4** Bayesian skyline plots of effective population size (N_e) scaled by generation time for CO1 mitochondrial
843 DNA. The plots run from the present to their median time to most recent common ancestor (T_{MRCA}). Grey dotted
844 lines represent the 95% CI for Net.

845

846

847

848

849

850

851

852

853

854

855

856

857

858

Table 1

Sampling location, population code, collection dates, and the number of individuals (n) screened at mitochondrial (CO1) and microsatellite markers.

Sampling sites	Lat (N)	Long (E)	Population Code	Collection date	Mitochondrial	Microsatellite	
					n	Collection date	n
Study Region: Western Luzon							
Bubon, Burgos, Ilocos Norte	18°30'31"	120°34'45"	Burgos (BUR)	Nov-08	29	Nov-08	34
Dadalaquiten, Sinit, Ilocos Sur	17°53'34"	120°26'28"	Sinit (SIN)	Dec-08	28	<i>n.d.</i>	<i>n.d.</i>
Lucero, Bolinao, Pangasinan	16°24'09"	119°54'30"	Lucero (LUC)	Nov-08	21	Nov-09	50
Victory, Bolinao, Pangasinan	16°23'35"	119°57'59"	Victory (VIC)	Nov-08	24	Nov-09	49
Panglit Is., Masinloc, Zambales	15°29'42"	119°55'09"	Masinloc (MAS)	Apr-09	30	Apr-09	48
Matuod, Lian, Batangas	13°59'11"	120°37'40"	Lian (LIA)	Feb-09	30	Feb-09	47
Outgroup Sites							
Bacon, Sorsogon	13°02'24"	124°02'56"	Sorsogon (SOR)	Jul-08	21	<i>n.d.</i>	<i>n.d.</i>
Lawi, Jordan, Guimaras	10°32'45"	122°31'16"	Guimaras (GUI)	Jan-09	47	Dec-09	49
Cantaan, Guinsiliban, Camiguin	09°06'25"	124°48'14"	Camiguin (CAM)	Mar-09	32	<i>n.d.</i>	<i>n.d.</i>
Simunul, Tawi-Tawi	04°54'03"	119°50'56"	Tawi-tawi (TAW)	May-09	20	<i>n.d.</i>	<i>n.d.</i>

Table 2

Summary statistics of *Tripneustes gratilla* populations based on CO1 sequences. Population, sampling size (n), measures of genetic diversity (Nh = no. of haplotypes; Np = no. of polymorphic sites; h = haplotype diversity; π = nucleotide diversity; s.d. = standard deviation), and Fu's neutrality test (Fu 1997). All F_S values were found to be significant ($p=0.0001$) indicating deviation from neutral equilibrium model.

Population	n	Nh	Np	h (s.d.)		π (s.d.)		Neutrality test (F_S)
Burgos	29	13	15	0.7660	(0.0816)	0.0030	(0.0020)	-7.851
Sinit	28	15	16	0.9101	(0.0370)	0.0033	(0.0022)	-10.583
Bolinao	45	20	21	0.8737	(0.0415)	0.0030	(0.0020)	-17.172
Lucero	21	11	11	0.9143	(0.0380)	0.0029	(0.0020)	
Victory	24	14	16	0.8333	(0.0767)	0.0031	(0.0020)	
Masinloc	30	13	12	0.7678	(0.0749)	0.0022	(0.0015)	-10.235
Lian	30	15	21	0.8736	(0.0498)	0.0035	(0.0022)	-9.610
Sorsogon	21	12	13	0.9095	(0.0479)	0.0036	(0.0023)	-6.915
Guimaras	47	18	23	0.8372	(0.0403)	0.0026	(0.0018)	-14.628
Camiguin	32	17	24	0.8347	(0.0646)	0.0036	(0.0023)	-12.468
Tawi-tawi	20	13	13	0.9263	(0.0431)	0.0033	(0.0021)	-9.810

Table 3

Bottleneck analysis on *Tripneustes gratilla* microsatellite data from 6 loci (without Tgr-24) under the assumption of infinite allele model. Asterisk (*) indicate heterozygosity excess ($H_e > H_{eq}$) based on the comparison of expected heterozygosity (H_e) and expected equilibrium heterozygosity (H_{eq}) of each locus per population. Wilcoxon's test determined the significance of heterozygosity excess across loci in a population. All p-values were significant ($p < 0.05$).

Population	Tgr-A11	Tgr-B11	Tgr-C11	Tgr-C117	Tgr-D134	Tgr-D5	Wilcoxon's test (p-value)
Burgos	*	*	*	*	*	*	0.008
Lucero	*	*	*	-	*	*	0.008
Victory	*	*	*	-	*	*	0.016
Masinloc	*	*	*	*	*	*	0.008
Lian	*	*	*	*	*	*	0.008
Guimaras	*	-	*	*	*	*	0.016




Fig. 1 Minimum spanning tree for the CO1 haplotypes of *T. gratilla*. Filled circle represents a unique haplotype which was sized proportionally to its absolute frequency, ranging from 1– 98. The dominant haplotypes were identified by their sequence number. The asterisk denotes the most probable root haplotype while the square represents unsampled haplotypes as revealed by TCS v1.21 (Clement et al. 2000).




Fig. 2 Neighbor-joining tree illustrating the relationship of the major CO1 haplotypes in this study and the CO1 sequences from Lessios et al. (2003) based on LogDet substitution model. Numbers next to nodes indicate bootstrap support from 1000 iterations and nodes with less than 50% support have been collapsed.




Fig. 3 Relative frequency of mitochondrial haplotypes per population. Represented in the pie graph are the major haplotypes (sequence 1, 10, and 3), minor haplotypes (pooled haplotypes with frequency of 2 to 7), and private haplotypes (haplotypes found in only one site).




Fig. 4 Bayesian skyline plots of effective population size (N_e) scaled by generation time for CO1 mitochondrial DNA. The plots run from the present to their median time to most recent common ancestor ($T_{MRC A}$). Grey dotted lines represent the 95% CI.