

[Chem. Pharm. Bull., 35, 2382 (1987)]

**Sulfur-Containing Acylamino Acids. I. Syntheses and Angiotensin I
Converting Enzyme-Inhibitory Activities of Sulfur-Containing *N*-
Mercaptoalkanoyl Amino Acids.**

TAKETOSHI KOMORI*, KATSUMI ASANO, YASUTO SASAKI, HIROMI HANAI,
RUMI SEO, MASANORI TAKAOKA, SHIRO MORIMOTO, MIKIO HORI

N-Mercaptoalkanoyl derivatives of sulfur-containing amino acids were synthesized and examined for inhibitory effects on angiotensin I converting enzyme (ACE) extracted from rabbit lung. Inhibition of ACE was determined by means of a spectrometric assay with hippuryl-L-histidyl-L-leucine as a substrate. Among the synthesized sulfur-containing compounds, *N*-(2-benzyl-3-mercaptopropanoyl)-*S*-methyl-L-cysteine and *N*-(2-benzyl-3-mercaptopropanoyl)-*S*-ethyl-L-cysteine showed the most potent inhibitory effects on ACE activity.

[Chem. Pharm. Bull., 35, 2388 (1987)]

**Sulfur-Containing Acylamino Acids. II. Syntheses and Angiotensin I
Converting Enzyme-Inhibitory Activities of *N*-Mercaptoalkanoyl-*S*-
ethyl-L-cysteine.**

TAKETOSHI KOMORI,* KATSUMI ASANO, YASUTO SASAKI, HIROMI HANAI,
SHIRO MORIMOTO, MIKIO HORI

N-Mercaptoalkanoyl derivatives of sulfur-containing amino acids were synthesized as candidate angiotensin I converting enzyme (ACE) inhibitors. Among them, *N*-(3-mercapto-2-(4-methoxybenzyl)propanoyl)-*S*-ethyl-L-cysteine was found to be the most potent inhibitor of ACE, with an IC_{50} value of $0.045\mu M$. The maximum hypotensive effect of this compound was almost equal to that of captopril in anesthetized rats.

[Synthesis, 1987, 278]

**Synthesis of 6-Substituted Purines from 3,7-Dimethyl-6-methylthio-
2-oxopurine.**

MIKIO HORI*, TADASHI KATAOKA, HIROSHI SHIMIZU, EIJI IMAI,
MASAHARU YOKOMOTO, YASUHIKO ANDO

Nucleophilic substitution of 3,7-dimethyl-6-methylthio-2-oxopurine with alcoholates or carbanions of active methylene compounds affords the corresponding 6-alkoxy- or 6-alkylidene-3,7-dimethyl-2-oxopurines, respectively, in good yields. The new method is valuable for the synthesis of the novel compounds with a carbon chain at the 6-position of the purine ring. This nucleophilic substitution at 6-position was performed by the direct substitution of methylthio group.