

[J. Chem. Soc., Perkin Trans. 1, 1990, 3017]

**A New Method for Radical Generation: Reductive C-Se or C-S Bond
Cleavage of Cyclic Onium Salts.**

TADASHI KATAOKA, KAZUHIRO TSUTSUMI, KENJI KANO, KAZUYA MORI,
MIHO MIYAKE, MANABU YOKOTA, HIROSHI SHIMIZU, MIKIO HORI*

2-Methyl-3,4-dihydro-1*H*-2-benzoselenopyranium salt was reduced by some metallic reagents or magnesium metal via the single electron transfer (SET) process to give 2-(2-(methylseleno)ethyl)-toluene. Magnesium metal was a good SET reducing agent. Some other selenonium or sulphonium salts were similarly reduced by magnesium. Particularly, sonication accelerated the reactions. ϵ -Ene-selenonium salt was treated with activated magnesium to give a cyclized pyrrolidine derivative.

[Chem. Pharm. Bull., 38, 8 (1990)]

Synthesis of 2,6-Epithio-3-benzazocine Derivatives.

MIKIO HORI*, TADASHI KATAOKA, HIROSHI SHIMIZU, EIJI IMAI,
TOKUO KOIDE, NORIYUKI IWATA, MASAYASU KURONO

2,6-Epithio-3-benzazocines (9-thiabenzomorphans) in which the carbon atom at the 11-position is replaced by a sulfur atom, were synthesized by treatment of 1-(2-ethoxycarbonylaminoethyl)isothiochroman sulfoxides with acetic anhydride or by heating 3-acetoxyisothiochromans in Dowtherm A. The hetero-acetal moiety of this novel heterocycle was stable to lithium aluminum hydride and boron tribromide. Reduction of the novel heterocycle with lithium aluminum hydride gave the *N*-methyl derivative and demethylation of the 8-methoxy derivative with boron tribromide gave the 8-hydroxy derivative.

[Chem. Pharm. Bull., 38, 779 (1990)]

**2-Methylisoselenochromanium Salts: Spectroscopic Properties and
Reactions.**

MIKIO HORI*, TADASHI KATAOKA, HIROSHI SHIMIZU,
KAZUHIRO TSUTSUMI

Some 2-methylisoselenochromanium salts were prepared. Proton nuclear magnetic resonance spectra in CDCl₃ and electron impact mass spectra showed that the selenonium tosylate and mesylate are selenuranes and that the tetrafluoroborate and the triflate are selenonium salts. Their structures at the selenium atom are influenced by their counter anions. Selenonium salts reacted with some nucleophiles to give a styrene derivative and methyl phenethyl selenides.