

[Biochem. J., 266, 583 (1990)]

**Demonstration of 3 $\alpha$  (17 $\beta$ )-hydroxysteroid dehydrogenase distinct from 3 $\alpha$ -hydroxysteroid dehydrogenase in hamster liver.**

MARI OHMURA, AKIRA HARA, MAKOTO NAKAGAWA, HIDEO SAWADA\*

NAD-linked and NADP-linked 3 $\alpha$ -hydroxysteroid dehydrogenases were purified from hamster liver cytosol. The two monomeric enzymes with Mr-38,000 differed in pI values, activation energy, heat stability and peptide fragment patterns. The NADP-linked enzyme catalyzed the oxidation of various 3 $\alpha$ -hydroxysteroids, whereas the NAD-linked enzyme oxidized both 3 $\alpha$ - and 17 $\beta$ -hydroxysteroids. The thermal stabilities of the 3 $\alpha$ - and 17 $\beta$ -hydroxysteroid dehydrogenase activities of the NAD-linked enzyme were identical, and the two enzyme activities were inhibited by mixing 17 $\alpha$ - and 3 $\beta$ -hydroxysteroid substrates, respectively. Synthetic steroids and 3 $\beta$ -hydroxysteroids competitively inhibited 3 $\alpha$ - and 17 $\beta$ -hydroxysteroid dehydrogenase activities of the enzyme.

[J. Biochem., 108, 250 (1990)]

**Purification and properties of multiple forms of dihydrodiol dehydrogenase from human liver.**

AKIRA HARA, HIROYUKI TANIGUCHI, TOSHIHIRO NAKAYAMA, HIDEO SAWADA\*

Two acidic and three basic forms of monomeric dihydrodiol dehydrogenase with molecular weights in the range of 36,000-39,000 were purified from human liver. One acidic enzyme was immunologically identified as aldehyde reductase. Two of the basic enzymes exhibited a 20 $\alpha$ -hydroxysteroid dehydrogenase activity and was sensitive to 1,10-phenanthroline, whereas the third basic enzyme oxidized some 3 $\alpha$ -hydroxysteroids at low rates and was inhibited by cyclopentane-1,1-diacetic acid. Another acidic enzyme showed a high 3 $\alpha$ -hydroxysteroid dehydrogenase activity and was the most sensitive to inhibition by medroxyprogesterone acetate. The Km values of the enzymes, except aldehyde reductase, for hydroxysteroids were lower than for xenobiotic alcohols.

[Biochem. Biophys. Res. Commun., 171, 661 (1990)]

**Primary Structure of Vitamin K-dependent Human Protein Z.**

HIROSHI SEJIMA, TATSUYA HAYASHII, YOSHIHIRO DEYASHIKI\*,  
JUNJI NISHIOKA, KOJI SUZUKI

The primary structure of a vitamin K-dependent human protein Z was determined by a combination of analyses of 41 amino acid residues of the NH<sub>2</sub>-terminal region and 1265 nucleotide base pairs of a cDNA encoding the residual COOH-terminal part of the protein and the 3' noncoding region. Human protein Z has 360 amino acid residues which is less than that of bovine protein Z containing 396 residues. Human protein Z was composed of an NH<sub>2</sub>-terminal domain rich in  $\gamma$ -carboxyglutamic acids, two epidermal growth factor-like domains and a COOH-terminal serine protease-like domain as was bovine protein Z.