

[Yakugaku Zasshi, 110, 453 (1990)]

Studies on the Methods for Chemical Identification of Forsythiae Fructus and Plantaginis Herba.

SANSEI NISHIBE, MICHIKO SASAHARA, MIYOSHI MIYAGETA, YUKIO NORO
TOMOKO KAWAMURA and TOSHIHIRO TANAKA*

On Forsythiae Fructus, a color reaction with 4-aminoantipyrine due to phenylethanoid glycoside, namely, forsythiaside and acteoside, is distinctive by showing red. Moreover, a color reaction with hydroxyamine and ferric chloride due to lignan, arctigenin, is distinctive by showing reddish purple, and a color reaction with sulfuric acid due to lignan, phillurin, is distinctive by showing red. The three origins of Forsythiae Fructus are distinguishable by difference of color reaction due to lignans. On Plantaginis Herba, a color reaction with 4-aminoantipyrine due to plantamajoside is remarkable and identification is also applicable by thin layer chromatography.

[Yakushigaku Zasshi, 25, 125 (1990)]

Yokusai Iinuma's Herbarium of "Sabina Called in Market"

MIZUO MIZUNO, TOSHIHIRO TANAKA*, EIJI SAKAI, YAN KUANG,
MASAHARU ENDO and JIN MURATA

The authors obtained the herbarium which Yokusai Iinuma had recorded as a "Sabina called in market" in his "Somoku-Zusetsu". The leaf of this plant was characterized with linear, entire margin, stomatal line, spiral arranged and twisted base. In the transverse section of leaf, fibrovascular bundle were vertical arranged to the longer axis and the fiber were continuously covered beneath the epidermis. These features corresponded with those of *Podocarpus imbricatus*. Another similar plants were collected and compared histologically. Considering with the ancient literatures, this herbarium recorded by Iinuma was identified to be *Podocarpus imbricatus*.

[Phytochemistry, 29, 857 (1990)]

Butanolides from *Litsea japonica*

HITOSHI TANAKA, TAKESHI NAKAMURA, KAZUHIKO ICHINO,
KAZUO ITO and TOSHIHIRO TANAKA*

Six lactones, litsenolide D₁, litsenolide D₂, litsenolide E₁, litsenolide E₂, hamabiwalactone A and hamabiwalactone B, were isolated from fresh leaves of *Litsea japonica* (japanese name "Hamabiwa") along with the known six litsenolides (litsenolide A₁, A₂, B₁, B₂, C₁ and C₂). Their structures were elucidated on the basis of spectroscopic evidence. The chemical structures of the new litsenolides (litsenolide D₁, D₂, E₁ and E₂) were determined to be α -alkylidene- β -hydroxy- γ -methyl- α, β' -unsaturated- γ -lactone. And the structures of hamabiwalactones were determined to be endocyclic α, β -unsaturated- γ -lactones. The isolation of hamabiwalactones from lauraceous plants is the first example.