

[Phytochemistry, 34, 1653-1654 (1993)]

[Lab. of Pharmacognosy]

Isoflavonoids in Roots of *Sophora fraseri*.

MUNEKAZU IINUMA*, MASAYOSHI OHYAMA, TOSHIYUKI TANAKA,

MERVYN P. HEGARTY, ELWYN E. HEGARTY

Munekazu Inuma*, Masayashi Ohyama, Toshiyuki Tanaka, Mervyn P.

Hegarty, Elwyn E. Hegarty

Six known phenolic compounds [sophoronol, sophoraisoflavanone A, maackiain, sophoracarpan B, calycosin and 2-(2',4'-dihydroxyphenyl)-5,6-methylenedioxybenzofuran], and a new isoflavanone were isolated from the roots of *Sophora fraserii*. The structure of the new compound, named fraserinone A, was shown to be 5,7,4'-trihydroxy-5'-(1,1-dimethylallyl)-2'-methoxyisoflavanone by means of spectroscopic analysis.

[Phytochemistry, 33, 841-846 (1993)]

[Lab. of Pharmacognosy]

Flavonol Glycosides Production in Cell Suspension Cultures of *Vancouveria hexandra*.

HIROBUMI YAMAMOTO*, KUANG YAN, KYOKO IEDA, TOSHIYUKI TANAKA,

MUNEKAZU IINUMA, MIZUO MIZUNO

Cell suspension cultures of *Vancouveria hexandra* produce a large amount of des-O-methylanhydroicaritin glycosides in addition to a small amount of anhydroicaritin glycosides. These cells also produced large amounts of kaempferol glycosides which do not occur in the original plant. Higher phosphate increased contents of 8-isoprenylated flavonol glycosides, but had practically no effect on that of kaempferol glycosides. This cell culture required 2,4-dichlorophenoxyacetic acid for the production of flavonol glycosides and high 2,4-D resulted the increase of anhydroicaritin glycoside contents without no effects on the other flavonol glycosides contents.

[J. Nat. Prod., 56, 1638-1641 (1993)]

[Lab. of Pharmacognosy]

Chemical Constituents in the Genus *Achlys*.

MUNEKAZU IINUMA*, HIDEKI TOSA, TOSHIYUKI TANAKA,

MIZUO MIZUNO, FRANK A. LANG

The structure of a new isocoumarin derivative, achlisocoumarin IV, isolated from the underground parts of *Achlys triphylla* was characterized by means of its spectroscopic properties. The chemical constituents of *A. triphylla* and *A. triphylla* subsp. *japonica* were compared by HPLC to find their chemotaxonomic similarities and differences. Results showed a chemotaxonomically close relationship.