

[Life Sciences, 55, PL233-237 (1994)]

[Lab. of Pharmacology]

**The model of arthritis induced by superantigen in mice.**

HIROICHI NAGAI\*, YUKO TAKAOKA, HIROYUKI KAMADA, HIROSHI MORI

Subcutaneous injection of Staphylococcal enterotoxine B (SEB) produced by *Staphylococcus aureus*, caused severe arthritis in DBA/1J mice which had been previously immunized with bovine type II collagen. The severity of this arthritis was dose-dependent and prolonged joint inflammation with erosion of bone was observed. Anti-type II collagen antibodies were detected in the serum of arthritic mice. Effector T cells against type II collagen were also detected by means of delayed type hypersensitivity in the skin. Moreover, a significant decrease in the ratio between T cells and B cells and an increase in the ratio between CD4<sup>+</sup> cells and CD8<sup>+</sup> cells were observed in spleen cells from arthritic mice. Prednisolone suppressed the induction and development of clinical signs of the arthritis in mice.

[Japan. J. Pharmacol., 66, 87-94 (1994)]

[Lab. of Pharmacology]

**Antiallergic effect of ZCR-2060: antihistaminic action.**

TOORU ABE, TAKESHI OMATA, KENJI YOSHIDA, TOSHIHIRO MATSUMURA, YUGO IKEDA,  
YOSHIHIDE SEGAWA, KAZUO MATSUDA, HIROICHI NAGAI\*

The antihistaminic effect of 2-[2-[4-(diphenylmethyl)-1-piperadiny]ethoxy]benzoic acid maleate (ZCR-2060), a newly synthesized antiallergic agent, was investigated. ZCR-2060 selectively antagonized histamine-induced contraction of isolated guinea pig ileum and trachea. <sup>3</sup>H-Mepyramine specific binding to membranes from guinea pig lung and brain was markedly inhibited by ZCR-2060 concentration-dependently. ZCR-2060 significantly inhibited the histamine-induced cutaneous reaction in rats, and its action was long-lasting. Thiopental-induced sleep and spontaneous ambulatory activity in mice, however, were unaffected by ZCR-2060 even at higher doses. ZCR-2060 is a potent, selective and long-acting histamine H<sub>1</sub>-receptor antagonist with no side effect.

[Japan. J. Pharmacol., 66, 95-103 (1994)]

[Lab. of Pharmacology]

**Antiallergic effects of ZCR-2060: effect on allergic cutaneous reactions  
and rhinitis models in mice and rats.**

TOORU ABE, TAKESHI OMATA, KENJI YOSHIDA, YOSHIHIDE SEGAWA, KAZUO MATSUDA,  
HIROICHI NAGAI\*

The antiallergic action of 2-[2-[4-(diphenylmethyl)-1-piperadiny]ethoxy]benzoic acid maleate (ZCR-2060) was investigated in mice and rats. ZCR-2060 markedly inhibited passive cutaneous anaphylaxis, histamine-, compound 48/80- and calcium ionophore A23187-induced cutaneous reactions, and IgE-mediated biphasic skin reaction. The antigen- and histamine-induced nasal vascular permeability increase in rats were clearly inhibited in a dose-dependent fashion by ZCR-2060. Moreover, ZCR-2060 significantly inhibited antigen-induced anaphylactic histamine release from rat peritoneal mast cells and carrageenin-induced paw edema in rats.