

[Natural Medicines, 48, 208-212 (1994)]

[Lab. of Herbal Garden]

Pharmacognostical Studies of *Houttuyniae Herba* (1) Flavonoid Glycosides
Contents of *Houttuynia cordata* THUNB..

TOMOKO KAWAMURA, YOUICHI HISATA, KAZUYO OKUDA, YUKIO NORO,
TOSHIHIRO TANAKA,* MASASHI YOSHIDA, EIJI SAKAI

The amounts of five flavonoid glycosides contained in *Houttuynia ordata*, quercetin, hyperin, rutin, isoquercitrin and afzerin were determined by HPLC. Leaves had a high quercitrin content and the spikes had high quercitrin and hyperin contents, but the stems contained only a little and the rhizomes scarcely any of these flavonoid glycosides. The flavonoid glycoside contents and the glycoside profiles in the leaves remained constant as the grew. The commercial crude drugs of Chinese origin tended to have higher hyperin contents than Japanese samples.

[Natural Medicines, 48, 237-243 (1994)]

[Lab. of Herbal Garden]

Studies on Genus *Urtica* in Turkey (1) Identification of *Urtica* in
Turkey by Internal Structure of Axis and Petiole.

MORIYUKI IIDA, YASUYUKI ASAMI, TOSHIHIRO TANAKA,* GISHO HONDA, MAMORU
TABATA, EKREM SEZIK, ERDEM YESHILADA

Of the 5 species growing in Turkey, 4 species, *U. pilulifera*, *U. urens* and *U. haussknechii* are used as medicinal plants. Microscopic studies revealed that 1) the transverse section of axis of all the species except *U. pilulifera* was rhomb, 2) the petiole of all of the species except *U. urens* had 5-7 vascular bundles, 3) clustered crystal of calcium oxalate was observed in the axis and petiole of *U. dioica* and *U. haussknechii*, only in the node of *U. urens* and not at all in *U. pilulifera*.

[Natural Medicines, 48, 317-321 (1994)]

[Lab. of Herbal Garden]

Report on Strain of *Zanthoxylum* sp. Called 'Budo-Zanshou' Cultivated in
Wakayama Prefecture.

KAZUHITO KAWAHARA, CHIHO KITANI, TOSHIHIRO TANAKA*

A strain of *Zanthoxylum piperitum* has been cultivated in the limited are of the Shimizu-Chou (Wakayama Pref.). The cultivation area is approximately 50 ha, and the crop of fresh fruit is about 150 tons a year at the present. The pericarp of the fruit of this strain is thicker, the oil cavity is bigger, and the (processing) yield of pericarp from the fruit is higher than those of *Z. piperitum* forma *inermis*, called 'Asakura-zanshou' in Japanese Market. Thus, the pericarp was experimentally proved to contain a larger amount of essential oil, which is the essential effective component. This strain should be accepted as an effective crude drug.