

[Tetrahedron Lett., **36**, 245 - 248 (1995)]

[Lab. of Pharm. Chemistry]

**Selective C-S Bond Cleavage of 3-Aryl- $\beta$ -sultams with EtAlCl<sub>2</sub>**

TADASHI KATAOKA,\* TETSUO IWAMA

Selective C-S bond cleavage of a  $\beta$ -sultam ring was achieved by the reactions of 3-aryl- $\beta$ -sultams with EtAlCl<sub>2</sub>. Aryl ketones or aldehydes were provided *via* processes of the C-S bond cleavage, 1,2-aryl shift and imine formation. These reactions were influenced by the cation stabilizing capability of C-4 substituents, by the electron density of C-3 aryl group and by the steric relation between C-3 and C-4 substituents, and anchimeric assistance played an important role for the selective C-S bond cleavage.

[Pharmacology Letters, **56**, 395 - 400 (1995)]

[Lab. of Pharm. Chemistry]

**Opioid Receptor Interaction and Adenylyl Cyclase Inhibition of Dihydroetorphine: Direct Comparison with Etorphine**

MASAYUKI NIWA, LUAY Y.AL-ESSA, SHUICHIRO OHTA, KEN-ICH KOHNO,

MASAKATSU NOZAKI, KAITO TSURUMI, TATSUNORI IWAMURA,\* TADASHI KATAOKA.

To find out the reason of weak addiction property of dihydroetorphine, we compared the affinities of dihydroetorphine to the type selective opioid receptor and inhibition effect on the adenylyl cyclase activity with those of etorphine. Dihydroetorphine and etorphine have almost the same binding affinities to all types ( $\mu$ ,  $\delta$  and  $\kappa$ ) of opioid receptors and antagonist binding sites, and have similar inhibition activities to forskolin stimulated adenylyl cyclase. However, dihydroetorphine showed significantly smaller value of DTNB-index compared with that of etorphine. This differentiation may explain partly the high analgesic with low dependence properties of dihydroetorphine.

[Chem. Lett., 459 -460 (1995)]

[Lab. of Pharm. Chemistry]

**Acid-promoted C<sub>1</sub>-C<sub>2</sub> Bond Fission and Subsequent 1,5-Sulfeny Shift of 1-Acceptor-1-sulfenyl-substituted 2-Vinylcyclopropanes. Formation of 6-Sulfenyl- $\alpha$ ,  $\beta$ - $\gamma$ ,  $\delta$ -unsaturated Carboxylic Esters and Nitriles**

TADASHI KATAOKA,\* HARUTOSHI MATSUMOTO, TETSUO IWAMA, HIROSHI SHIMIZU

1-Acceptor-1-sulfenyl-substituted 2-vinylcyclopropanes were isomerized to 6-sulfenyl- $\alpha$ ,  $\beta$ - $\gamma$ ,  $\delta$ -unsaturated carboxylic esters and nitriles *via* a C<sub>1</sub>-C<sub>2</sub> bond fission and *via* a 1,5-sulfenyl shift using an acid catalyst. The reaction efficiently took place using a sulfonic acid such as *p*-toluenesulfonic acid monohydrate and trifluoromethanesulfonic acid in benzene, a nonpolar solvent. The substituent effect of an arylthio group was slightly observed. The cross-over experiment revealed that the 1,5-sulfenyl shift proceeded intermolecularly.