

[Int. Arch. Allergy Immunol., **107**, 587-591 (1995)]

[Lab. of Pharmacology]

Relationship between histamine release and leukotrienes production from human basophils derived from atopic dermatitis donors.

MICHITAKA SHICHIJO, MOTOHIRO EBISAWA, KATSUHIRO MIURA, YUMIKO TOIDA,
TAKEBUMI ONDA, HIROHISA SAITO, HIROICHI NAGAI*, YOJI IIKURA

We examined the relationship between histamine and leukotrienes (LTs) production from basophils in the presence or absence of 1 ng/ml of interleukin-3 (IL-3). Normal basophils released a smaller amount of histamine and LTs than atopic dermatitis (AD) basophils stimulated with an optimal concentration of anti-IgE antibody. When we examined the relationship between these mediator release from AD donors, we found a significant exponential correlation between these two mediators. Although IL-3 enhanced both histamine release and LTs production from AD donors, the relationship between these two mediators was not affected.

[Pharmacol. Commun., **6**, 361-367 (1995)]

[Lab. of Pharmacology]

Inhibitory mechanisms of prednisolone for delayed type hypersensitivity reactions in mice.

NAOKI INAGAKI, TAKANORI SENGOKU, SATORU NIWA, HIROSHI MORI, HIROICHI NAGAI*, AKIHIDE KODA

The inhibitory mechanisms of prednisolone (Pred) on the effector phase of delayed type hypersensitivity (DTH) reactions caused by picryl chloride and tuberculin were investigated in mice. Pred administered intraperitoneally 6-22h after challenge significantly inhibited both DTH reactions. The inhibition caused by Pred given 14h after challenge was not affected by 17 α -methyltestosterone or cycloheximide. On the contrary, the agents completely abrogated the inhibition caused by Pred given 22h after challenge.

[Int. Arch Allergy Immunol. **108**, 189-195 (1995)]

[Lab. of Pharmacology]

Effects of some immunosuppressors on allergic bronchial inflammation and airway hyperresponsiveness in mice.

HIROICHI NAGAI*, SHUJI YAMAGUCHI, HIROYUKI TANAKA, NAOKI INAGAKI

The effects of two new immunosuppressors, FK-506 (FK) and mizoribine (MZ), on antigen-induced bronchial inflammation and reactivity to acetylcholine in mice were studied in comparison with those of cyclosporin A (CsA) and cyclophosphamide (CP). The administration of each of the four immunosuppressors clearly inhibited the airway eosinophilia. Moreover, FK, MZ and CP clearly inhibited the interleukin-5 (IL-5) production and CsA showed the tendency to inhibit IL-5 production. Whereas FK, MZ and CsA clearly inhibited the airway hyperreactivity, CP did not show a significant effect on this airway hyperreactivity. These two drugs inhibited the airway hyperreactivity, probably by another mechanism to inhibit airway eosinophilia and IL-5 production.