

[*Phytochemistry*, **53**, 1015-1019 (2000)]

[Lab. of Pharmacognosy]

Stilbenoids in the Stem Bark of *Hopea parviflora*.

Toshiyuki TANAKA,* Testuro ITO, Yoshimi IDO, SON T.-K., Ken-ichi NAKAYA,
Munekazu IINUMA, Masayoshi OHYAMA and Vhelladurai VEILI

From the bark of *Hopea parviflora*, two stilbenoids, named (+)-parviflorol and (-)-ampelopsin A, were isolated in addition to three known compounds: (+)-balanocarphol, (-)- ϵ -viniferin and (-)-hopeaphenol. Their structures were determined by analysis of spectral data, including 2D NMR and NOE experiments, respectively.

[*Phytochemistry*, **54**, 63-69 (2000)]

[Lab. of Pharmacognosy]

Oligostilbenoids in Stem Bark of *Vatica rassak*.

Toshiyuki TANAKA,* Testuro ITO, Ken-ichi NAKAYA, Munekazu IINUMA and Seodarsono RISWAN

Three resveratrol oligomers, vaticanols A, B and C, as well as three known stilbenoids, resveratrol, piceid and ϵ -viniferin were isolated from the stem bark of *Vatica rassak* (Dipterocarpaceae). Their structures were determined by the analysis of NMR spectral data including the application of 2D methods.

[*Chem. Pharm. Bull.*, **48**, 1001-1005 (2000)]

[Lab. of Pharmacognosy]

Stilbenoids Isolated from Stem Bark of *Shorea hemsleyana*.

Testuro ITO, Toshiyuki TANAKA,* Yoshimi IDO, Ken-ichi NAKAYA,
Munekazu IINUMA and Seodarsono RISWAN

Two new stilbene glucosides [(+)- α -viniferin 13b- β -glucopyranoside and resveratrol 12-C- β -glucoside] and two new resveratrol oligomers, hemsleyanol A and B were isolated from the bark of *Shorea hemsleyana* along with four known resveratrol oligomers. The structures of the isolated, including the relative configuration, were established by spectroscopic data involving long-range coupling and nuclea Overhauser effect experiments.

[*Chem. Pharm. Bull.*, **48**, 1959-1963 (2000)]

[Lab. of Pharmacognosy]

Four New Stilbenoids C-Glucosides Isolated from the Stem Bark of *Shorea hemsleyana*.

Testuro ITO, Toshiyuki TANAKA,* Yoshimi IDO, Ken-ichi NAKAYA,
Munekazu IINUMA and Seodarsono RISWAN

Four new C-glucopyranoside of resveratrol oligomers, hemslyanosides A-D, were isolated from the bark of *Shorea hemsleyana*. The structures were established on the basis of spectroscopic evidence, including ^1H - ^1H and ^{13}C - ^1H long-range couplings and nuclear Overhauser effect experiments in the NMR spectrum.