

SUT Journal of Mathematics  
Vol. 51, No. 2 (2015), 181–193

## Odd harmonious labeling of some new families of graphs

P. Jeyanthi, S. Philo and Kiki A. Sugeng

(Received April 12, 2015; Revised October 31, 2015)

**Abstract.** A graph  $G(p, q)$  is said to be odd harmonious if there exists an injection  $f : V(G) \rightarrow \{0, 1, 2, \dots, 2q - 1\}$  such that the induced function  $f^* : E(G) \rightarrow \{1, 3, \dots, 2q - 1\}$  defined by  $f^*(uv) = f(u) + f(v)$  is a bijection. A graph that admits odd harmonious labeling is called odd harmonious graph. In this paper, we prove that shadow and splitting of graph  $K_{2,n}$ ,  $C_n$  for  $n \equiv 0 \pmod{4}$ , the graph  $H_{n,n}$ , double quadrilateral snakes  $DQ(n)$ ,  $n \geq 2$ , the graph  $P_{r,m}$  if  $m$  is odd, banana tree and the path union of cycles  $C_n$  for  $n \equiv 0 \pmod{4}$  are odd harmonious.

*AMS 2010 Mathematics Subject Classification.* 05C78.

*Key words and phrases.* Harmonious labeling, odd harmonious labeling, shadow graph, splitting graph, double quadrilateral snake.

### §1. Introduction

Throughout this paper by a graph we mean a finite, simple and undirected one. For standard terminology and notation we follow Harary [6]. A graph  $G = (V, E)$  with  $p$  vertices and  $q$  edges is called a  $(p, q)$  – graph. The graph labeling is an assignment of integers to the set of vertices or edges or both, subject to certain conditions. An extensive survey of various graph labeling problems is available in [3]. Labeled graphs serves as useful mathematical models for many applications such as coding theory, including the design of good radar type codes, synch-set codes, missile guidance codes and convolution codes with optimal autocorrelation properties. They facilitate the optimal nonstandard encoding of integers. Graham and Sloane [4] introduced harmonious labeling during their study of modular versions of additive bases problems stemming from error correcting codes. A graph  $G$  is said to be harmonious if there exists an injection  $f : V(G) \rightarrow Z_q$  such that the induced function  $f^* : E(G) \rightarrow Z_q$  defined by  $f^*(uv) = (f(u) + f(v)) \pmod{q}$  is a

bijection and  $f$  is called harmonious labeling of  $G$ . The concept of odd harmonious labeling was followed Liang and Bai [7]. A labeling is said to be odd harmonious if there exists an injection  $f : V(G) \rightarrow \{0, 1, 2, \dots, 2q - 1\}$  such that the induced function  $f^* : E(G) \rightarrow \{1, 3, \dots, 2q - 1\}$  defined by  $f^*(uv) = f(u) + f(v)$  is a bijection. A graph that admits odd harmonious labeling is called odd harmonious graph. The odd harmoniousness of graph is useful for the solution of undetermined equations. The same authors have obtained necessary conditions for the existence of odd harmonious labeling of graphs:

1. If  $G$  is an odd harmonious graph, then  $G$  is a bipartite graph.
2. If a  $(p, q)$ -graph  $G$  is odd harmonious, then  $2\sqrt{q} \leq p \leq 2q - 1$ .

Several results have been published on odd harmonious labeling see [1, 2, 5, 8, 9, 10]. Motivated by these results, in this paper we prove that the shadow and splitting of the graphs  $K_{2,n}$ ,  $C_n$  for  $n \equiv 0 \pmod{4}$ , the graph  $H_{n,n}$ , double quadrilateral snakes  $DQ(n)$ ,  $n \geq 2$ , the graph  $P_{r,m}$  if  $m$  is odd, banana tree and the path union of cycles  $C_n$  for  $n \equiv 0 \pmod{4}$  are odd harmonious.

**Definition 1.** A function  $f$  is said to be a strongly odd harmonious labeling of a graph  $G$  with  $q$  edges if  $f$  is an injection from the vertices of  $G$  to the integers from 0 to  $q$  such that the induced mapping  $f^*(uv) = f(u) + f(v)$  from the edges of  $G$  to the odd integers between 1 to  $2q - 1$  is a bijection.

**Definition 2.** [9] The shadow graph  $D_2(G)$  of a connected graph  $G$  is constructed by taking two copies of  $G$  say  $G'$  and  $G''$  and join each vertex  $u'$  in  $G'$  to the neighbours of the corresponding vertex  $v'$  in  $G''$ .

**Definition 3.** [9] For a graph  $G$  the splitting graph  $spl(G)$  of a graph  $G$  is obtained by adding a new vertex  $v'$  corresponding to each vertex  $v$  of  $G$  such that  $N(v) = N(v')$ .

**Definition 4.** The graph  $H_{n,n}$  has the vertex set  $V(H_{n,n}) = \{v_1, v_2, \dots, v_n, u_1, u_2, \dots, u_n\}$  and the edge set  $E(H_{n,n}) = \{v_i u_j : 1 \leq i \leq n, n - i + 1 \leq j \leq n\}$ .

**Definition 5.** Let  $Q(n)$  be the quadrilateral snake obtained from the path  $v_1, v_2, v_3, \dots, v_{n+1}$  by joining  $v_i$  and  $v_{i+1}$  to the new vertices  $u_i$  and  $w_i$ . That is, every edge of a path is replaced by a cycle  $C_4$ .

**Definition 6.** Let  $Q(n)$  be the quadrilateral snake obtained from the path  $v_1, v_2, v_3, \dots, v_{n+1}$ . The double quadrilateral snake  $DQ(n)$  is obtained from  $Q(n)$  by adding the vertices  $s_1, s_2, s_3, \dots, s_n; t_1, t_2, t_3, \dots, t_n$  and the edges  $v_i s_i, t_i v_{i+1}, s_i t_i$  for  $1 \leq i \leq n$ .

**Definition 7.** Let  $u$  and  $v$  be the fixed vertices and connect  $u$  and  $v$  by means of  $b \geq 2$  internally disjoint paths of length  $a \geq 2$  each. The resulting graph embedded in a plane is denoted by  $P_{a,b}$ . Let  $v_0^i, v_1^i, v_2^i, \dots, v_a^i$  be the vertices of the  $i^{th}$  copy of the path of length  $a$  where  $i = 1, 2, \dots, b, v_0^i = u$  and  $v_a^i = v$  for all  $i$ . We observe that the graph  $P_{a,b}$  has  $(a - 1)b + 2$  vertices and  $ab$  edges.

**Definition 8.** For graphs  $G_1, G_2, \dots, G_n (n \geq 2)$ , we call a graph obtained by adding an edge from  $G_i$  to  $G_{i+1}$  for  $i = 1, 2, \dots, n - 1$  a path-union of  $G_1, G_2, \dots, G_n$  (the resulting graph may depends on how the edges are chosen).

**Definition 9.** [3] A banana tree is a graph obtained by connecting a vertex  $v$  to one leaf of each of any number of stars. Let  $K_{1,n_1}, K_{1,n_2}, \dots, K_{1,n_k}$  be a family of disjoint stars. Let  $v$  be a new vertex and the tree obtained by joining  $v$  to one pendant vertex of each star is called a banana tree. The class of all such trees is denoted by  $BT(n_1, n_2, \dots, n_k)$ . If  $n_1 = n_2 = \dots = n_k$  we denote  $BT(n_1, n_2, \dots, n_k)$  as  $BT_k(n)$

### §2. Main Results

**Theorem 2.1.** The shadow graph  $D_2(K_{2,n})$  is an odd harmonious graph.

*Proof.* Consider the two copies of  $K_{2,n}$ . Let  $v_1, v_2, v_3, \dots, v_n$  be the vertices of the first copy of  $K_{2,n}$  adjacent with  $u$  and  $v$ . Let  $v'_1, v'_2, v'_3, \dots, v'_n$  be the vertices of the second copy of  $K_{2,n}$  adjacent with  $u$  and  $v$ . Let  $G = D_2(K_{2,n})$ . Then  $V(G) = \{u, v, u', v', v_i, v'_i | 1 \leq i \leq n\}$  and  $E(G) = \{uv_i, vv_i, u'v'_i, v'v'_i, uv'_i, vv'_i, u'v_i, v'v_i | 1 \leq i \leq n\}$ . Hence  $|V(G)| = 2n + 4$  and  $|E(G)| = 8n$ . We define a labeling  $f : V(G) \rightarrow \{0, 1, 2, \dots, 16n - 1\}$  as follows:

$f(u) = 0; f(v) = 4n; f(u') = 8n; f(v') = 12n; f(v_i) = 2i - 1, 1 \leq i \leq n; f(v'_i) = 2n - 1 + 2i, 1 \leq i \leq n$ . The induced edge labelings are as follows: For  $1 \leq i \leq n$

$f^*(uv_i) = 2i - 1; f^*(uv'_i) = 2n + 2i - 1; f^*(vv_i) = 4n + 2i - 1; f^*(vv'_i) = 6n + 2i - 1; f^*(u'v_i) = 8n + 2i - 1; f^*(u'v'_i) = 10n + 2i - 1; f^*(v'v_i) = 12n + 2i - 1; f^*(v'v'_i) = 14n + 2i - 1$ . Hence  $f^*(E(G)) = \{1, 3, \dots, 2n - 1, 4n + 1, 4n + 3, \dots, 6n - 1, 10n + 1, 10n + 3, \dots, 12n - 1, 14n + 1, 14n + 3, \dots, 16n - 1, 2n + 1, 2n + 3, \dots, 4n - 1, 6n + 1, 6n + 3, \dots, 8n - 1, 8n + 3, \dots, 10n - 1, 12n + 1, 12n + 3, \dots, 14n - 1\} = \{1, 3, \dots, 16n - 1\}$

Thus  $f$  admits odd harmonious labeling on  $D_2(K_{2,n})$ . □

**Illustration 1.** The odd harmonious labeling of the graph  $D_2(K_{2,5})$  is given in Figure 1




Figure 1: Odd harmonious labeling of  $D_2(K_{2,5})$

**Corollary 2.2.** *The graph  $spl(K_{2,n})$  is an odd harmonious graph.*

*Proof.* Let  $G = spl(K_{2,n})$ . Then  $G$  is isomorphic to a graph obtained from the graph  $D_2(K_{2,n})$  by deleting the edges  $u'v'_i, v'v'_i (1 \leq i \leq n)$ . Hence  $|V(G)| = 2n + 4$  and  $|E(G)| = 6n$ . By using the labeling given in Theorem 2.1, we get the induced edge labels as  $\{1, 3, \dots, 12n - 1\}$ . Hence  $spl(K_{2,n})$  is an odd harmonious graph.  $\square$

**Illustration 2.** *The odd harmonious labeling of the graph  $spl(K_{2,4})$  is given in Figure 2*




Figure 2: Odd harmonious labeling of  $spl(K_{2,4})$

**Theorem 2.3.** *The shadow graph  $D_2(C_n)$  is an odd harmonious graph if  $n \equiv 0 \pmod{4}$ .*

*Proof.* Let  $v_1, v_2, v_3, \dots, v_n$  be the vertices of the first copy of  $C_n$  and  $u_1, u_2, u_3, \dots, u_n$  are the vertices of the second copy of  $C_n$ . Let  $G = D_2(C_n)$  and  $n \equiv 0 \pmod{4}$ . Then  $V(G) = \{v_i, u_i | 1 \leq i \leq n\}$  and  $E(G) = \{v_i v_{i+1}, u_i u_{i+1}, v_n v_1,$

$u_n u_1 | 1 \leq i \leq n - 1 \} \cup \{v_i u_{i+1}, v_n u_1 | 1 \leq i \leq n - 1 \} \cup \{v_1 u_n, v_i u_{i-1} | 2 \leq i \leq n \}$ . Here  $|V(G)| = 2n$  and  $|E(G)| = 4n$ . We define  $f : V(G) \rightarrow \{0, 1, 2, \dots, 8n - 1\}$  as follows:

$$f(v_i) = \begin{cases} i - 1 & \text{if } 1 \leq i \leq \frac{n}{2}, \\ i + 1 & \text{if } i \text{ is odd and } \frac{n}{2} + 1 \leq i \leq n \\ i - 1 & \text{if } i \text{ is even and } \frac{n}{2} + 1 \leq i \leq n \end{cases}$$

$$\text{For } 1 \leq i \leq n \quad f(u_i) = \begin{cases} f(v_i) + 2n & \text{if } i \text{ is odd,} \\ f(v_i) + 4n & \text{if } i \text{ is even.} \end{cases}$$

The induced edge labelings are as follows:  $f^*(\{v_i v_{i+1}, u_i u_{i+1}, v_n v_1, u_n u_1 | 1 \leq i \leq n - 1 \}) = \{1, 3, \dots, 2n - 1\} \cup \{6n + 1, 6n + 3, \dots, 8n - 1\}$

$f^*(\{v_i u_{i+1}, v_n v_1 | 1 \leq i \leq n - 1 \}) = \{2n + 3, 4n + 5, 2n + 7, \dots, 5n - 3, 3n + 1, 5n + 3, 3n + 5, \dots, 6n - 1, 4n + 1\}$

$f^*(\{v_1 u_n, v_i u_{i-1} | 2 \leq i \leq n \}) = \{5n - 1, 2n + 1, 4n + 3, 2n + 5, 4n - 7, \dots, 3n - 3, 5n + 1, 3n + 3, \dots, 6n - 3, 4n - 1\}$ .

Hence  $f^*(E(G)) = \{1, 3, \dots, 8n - 1\}$ . Thus  $D_2(C_n)$  is an odd harmonious graph. □

**Illustration 3.** The odd harmonious labeling of  $D_2(C_8)$  is given in Figure 3




Figure 3: Odd harmonious labeling of  $D_2(C_8)$

**Theorem 2.4.** The graph  $spl(C_n)$  if  $n \equiv 0 \pmod{4}$  is an odd harmonious graph .

*Proof.* Let  $G = spl(C_n)$ . Then  $G$  is isomorphic to a graph obtained from the graph  $D_2(C_n)$  by deleting the edges  $u_i u_{i+1}, u_n u_1 (1 \leq i \leq n - 1)$ . Hence  $|V(G)| = 2n$  and  $|E(G)| = 3n$ . By using the labeling given in Theorem 2.3, we get the induced edge labels as  $\{1, 3, \dots, 6n - 1\}$ . Hence  $spl(C_n)$  is an odd harmonious graph. □

**Illustration 4.** The odd harmonious labeling of  $spl(C_8)$  is shown in Figure 4




Figure 4: Odd harmonious labeling of  $spl(C_8)$

**Theorem 2.5.** *The graph  $D_2(H_{n,n})$  is an odd harmonious graph .*

*Proof.* Let  $v_1, v_2, v_3, \dots, v_n, u_1, u_2, u_3 \dots, u_n$  be the vertices of first copy of  $H_{n,n}$  and  $v'_1, v'_2, v'_3, \dots, v'_n, u'_1, u'_2, u'_3 \dots, u'_n$  be the vertices of second copy of  $H_{n,n}$ . Let  $G = D_2(H_{n,n})$ . Then  $|V(G)| = 4n$  and  $|E(G)| = 2n(n + 1)$ . We define  $f : V(G) \rightarrow \{0, 1, 2, \dots, 4n(n + 1) - 1\}$  as follows:

$$\begin{aligned}
 f(v_i) &= i(i - 1), 1 \leq i \leq n \\
 f(u_j) &= (2n + 1) - 2j, 1 \leq j \leq n \\
 f(v'_i) &= i(i - 1) + n(n + 1), 1 \leq i \leq n \\
 f(u'_j) &= (2n + 1) - 2j + 2n(n + 1), 1 \leq j \leq n
 \end{aligned}$$

The induced edge labelings are as follows:

$$\begin{aligned}
 f^*(v_i u_j) &= i(i - 1) + (2n + 1) - 2j \\
 f^*(v'_i u'_j) &= i(i - 1) + 3n(n + 1) + (2n + 1) - 2j \\
 f^*(v'_i u_j) &= i(i - 1) + n(n + 1) + (2n + 1) - 2j \\
 f^*(v_i u'_j) &= i(i - 1) + (2n + 1) - 2j + 2n(n + 1).
 \end{aligned}$$

The induced edge labeling  $f^*(E(G)) = \{1, 3, 5, \dots, 4n(n + 1) - 1\}$ . Thus  $f$  is an odd harmonious labeling of  $G$ . Hence the graph  $D_2(H_{n,n})$  is an odd harmonious graph. □

**Illustration 5.** *The odd harmonious labeling of  $D_2(H_{3,3})$  is given in Figure*

5




Figure 5: Odd harmonious labeling of  $D_2(H_{3,3})$

**Theorem 2.6.** *The graph  $spl(H_{n,n})$  is an odd harmonious graph.*

*Proof.* Let  $v_1, v_2, v_3, \dots, v_n, u_1, u_2, u_3, \dots, u_n$  be the vertices of first copy of  $H_{n,n}$ . Let  $v'_1, v'_2, v'_3, \dots, v'_n, u'_1, u'_2, u'_3, \dots, u'_n$  be the new vertices added to the corresponding vertices  $v_1, v_2, v_3, \dots, v_n, u_1, u_2, u_3, \dots, u_n$  respectively. Let  $G = spl(H_{n,n})$ . Then  $|V(G)| = 4n$  and  $|E(G)| = |E(H_{n-1,n-1})| + 3n$ .

We define  $f : V(G) \rightarrow \{0, 1, 2, \dots, 2(|E(H_{n-1,n-1})| + 3n) - 1\}$  as follows:

$$\begin{aligned} f(v_i) &= i(i-1), 1 \leq i \leq n \\ f(u_j) &= (2n+1) - 2j, 1 \leq j \leq n \\ f(v'_i) &= i(i-1) + n(n+1), 1 \leq i \leq n \\ f(u'_j) &= (2n+1) - 2j + 2n(n+1), 1 \leq j \leq n \end{aligned}$$

The induced edge labelings are as follows:

$$\begin{aligned} f^*(v_i u_j) &= i(i-1) + (2n+1) - 2j \\ f^*(v'_i u'_j) &= i(i-1) + n(n+1) + (2n+1) - 2j \\ f^*(v_i u'_j) &= i(i-1) + (2n+1) - 2j + 2n(n+1). \end{aligned}$$

The induced edge labeling is  $f^*(E(G)) = \{1, 3, 5, \dots, 2(|E(H_{n-1,n-1})| + 3n) - 1\}$ . Thus  $f$  is an odd harmonious labeling of  $G$ . Hence the graph  $spl(H_{n,n})$  is an odd harmonious graph.  $\square$

**Illustration 6.** *The odd harmonious labeling of  $spl(H_{3,3})$  is shown in Figure 6*




Figure 6: Odd harmonious labeling of  $spl(H_{3,3})$

**Theorem 2.7.** *The double quadrilateral snake  $DQ(n)$  is an odd harmonious graph for  $n \geq 2$ .*

*Proof.* Let  $v_1, v_2, v_3, \dots, v_n, v_{n+1}; u_1, u_2, u_3, \dots, u_n; w_1, w_2, w_3, \dots, w_n; s_1, s_2, s_3, \dots, s_n; t_1, t_2, t_3, \dots, t_n$  be the vertices of  $DQ(n)$ . Let  $G = DQ(n)$ . Hence  $|V(G)| = 5n + 1$  and  $|E(G)| = 7n$ . We define  $f : V(G) \rightarrow \{0, 1, 2, \dots, 14n - 1\}$  as follows:

$$\begin{aligned} &\text{If } n \text{ is even, then } f(v_1) = 0 \text{ and } f(v_{n+1}) = 7n \\ &\text{If } n \text{ is odd, then } f(v_1) = 0 \text{ and } f(v_{n+1}) = 7n - 4 \\ &\text{Also } f(v_i) = \begin{cases} f(v_{i-1}) + 3 & \text{if } i \text{ is even,} \\ f(v_{i-1}) + 11 & \text{if } i \text{ is odd} \end{cases}, 2 \leq i \leq n + 1 \\ &\text{Now } f(u_i) = \begin{cases} f(v_i) + 1 & \text{if } i \text{ is odd,} \\ f(v_i) + 9 & \text{if } i \text{ is even} \end{cases}, 1 \leq i \leq n \\ &f(w_i) = f(v_i) + 6, 1 \leq i \leq n \\ &f(s_i) = \begin{cases} f(v_i) + 5 & \text{if } i \text{ is odd,} \\ f(v_i) + 13 & \text{if } i \text{ is even} \end{cases}, 1 \leq i \leq n \\ &f(t_i) = f(v_i) + 8, 1 \leq i \leq n \end{aligned}$$

The induced edge labelings are as follows:

$$\begin{aligned} f^*(v_i v_{i+1}) &= 14i - 11, 1 \leq i \leq n \\ f^*(v_i u_i) &= 14i - 13, 1 \leq i \leq n \\ f^*(u_i w_i) &= 14i - 7, 1 \leq i \leq n \\ f^*(v_i s_i) &= 14i - 9, 1 \leq i \leq n \\ f^*(s_i t_i) &= 14i - 1, 1 \leq i \leq n \\ f^*(w_i v_{i+1}) &= 14i - 5, 1 \leq i \leq n \\ f^*(t_i v_{i+1}) &= 14i - 3, 1 \leq i \leq n \end{aligned}$$

In view of the above defined labeling pattern,  $f$  admits odd harmonious labeling for  $DQ(n)$ . Hence  $DQ(n), n \geq 2$  is an odd harmonious graph.  $\square$

**Illustration 7.** *The odd harmonious labeling of  $DQ(4)$  is shown in Figure 7*




Figure 7: Odd harmonious labeling of  $DQ(4)$

**Theorem 2.8.** *If  $m$  is odd, then  $P_{r,m}$  is an odd harmonious graph for all the values of  $r > 1$ .*



*Proof.* let  $v_0^i, v_1^i, v_2^i, \dots, v_{2r}^i$  be the vertices of the  $i^{th}$  copy of the path of length  $r$  where  $i = 1, 2, \dots, m$ ,  $v_0^i = u$  and  $v_{2r}^i = v$  for all  $i$ . We observe that the number of vertices of the graph  $P_{r,m}$  has  $(r - 1)m + 2$  vertices and the number of edges of the graph is  $rm$ . We define  $f : V(G) \rightarrow \{0, 1, 2, \dots, 2rm - 1\}$  as follows:

$$f(u) = 0; f(v) = rm; f(v_1^1) = 1$$

$$\text{If } j = 1, \text{ then } f(v_j^i) = f(v_{j-1}^{i-1}) + 2, i = 2, 3, \dots, m.$$

$$\text{If } j = 2, \text{ then } f(v_j^i) = \begin{cases} f(v_{j-1}^i) + 2m + 1, & i = 1, 2, \dots, (\frac{m-1}{2}), \\ f(v_{j-1}^i) + 1 & (\frac{m-1}{2}) + 1 \leq i \leq m \end{cases}$$

If  $j = 3, 4, \dots, r - 1$ , then  $f(v_j^i) = f(v_{j-2}^i) + 2m, i = 1, 2, \dots, m$ . The induced edge labelings are as follows:

$$f^*(uv_1^i) = 2i - 1, 1 \leq i \leq m.$$

**Case i.**  $1 \leq i \leq (\frac{m-1}{2})$ .

$$f^*(v_j^i v_{j+1}^i) = 2jm + 4i - 1, j = 1, 2, 3, 4, 5, \dots (r - 2)$$

**Case ii.**  $(\frac{m-1}{2}) + 1 \leq i \leq m$ .

$$f^*(v_j^i v_{j+1}^i) = 2(j - 1)m + 4i - 1, j = 1, 2, 3, 4, 5, \dots (r - 2)$$

$$\text{If } j = r - 1 \text{ is even, then } f^*(v_{r-1}^i v) = \begin{cases} 4m + rm + 2i, & 1 \leq i \leq (\frac{m-1}{2}), \\ 2m + rm + 2i & (\frac{m-1}{2}) + 1 \leq i \leq m \end{cases}$$

If  $j = r - 1$  is odd, then  $f^*(v_{r-1}^i v) = 4m + rm + 2i - 1, 1 \leq i \leq m$ . In view of the above defined labeling pattern  $f$  is an odd harmonious labeling for  $P_{r,m}$ . Hence  $P_{r,m}$  is an odd harmonious graph for all the values of  $r > 1$ .  $\square$

**Illustration 8.** The odd harmonious labeling of  $P_{5,5}$  is shown below:




Figure 8: Odd harmonious labeling of  $P_{5,5}$

**Theorem 2.9.** Let  $G_1(p_1, q_1), G_2(p_2, q_2), \dots, G_m(p_m, q_m)$  be a strongly odd harmonious graphs and  $u_i$  and  $v_i$  be the vertices of  $G_i$  labeled with 0 and  $q_i$  ( $1 \leq i \leq m$ ) respectively. Then the path union graph  $G$  obtained by joining  $u_i$  with  $v_{i+1}$  by an edge for each  $i, 1 \leq i \leq m - 1$  is a strongly odd harmonious graph.

*Proof.* The path union graph  $G$  has  $q = q_1 + q_2 + q_m + m - 1$  edges. Let  $f_i$  be the strongly odd harmonious labeling of  $G_i$  and  $V(G_i) = \{x_{ij} : j = 1, 2, \dots, p_i\}$ ,  $1 \leq i \leq m$ . Then  $V(G) = \bigcup_{i=1}^m V(G_i)$ . Define a vertex labeling  $f : V(G) \rightarrow \{0, 1, 2, \dots, q\}$  as  $f(x_{ij}) = f_i(x_{ij}) + \sum_{k=1}^{i-1} q_k + i - 1$ . For each  $i$ , the vertex labels of  $G_i$  satisfy the inequality  $\sum_{k=1}^{i-1} q_k + i - 1 \leq f(x_{ij}) \leq \sum_{k=1}^i q_k + i - 1$ . Since  $f_i$  is an injection,  $f$  is also injection. The maximum vertex label of  $G$  is  $\sum_{k=1}^m q_k + m - 1$ . The set of edge labels of  $G_i$  is  $\left\{ 2 \sum_{k=1}^{i-1} q_k + 2i - 1, 2 \sum_{k=1}^{i-1} q_k + 2i + 1, \dots, 2 \sum_{k=1}^i q_k + 2i - 3 \right\}$  for  $1 \leq i \leq m$ . The label of the bridge between  $G_i$  and  $G_{i+1}$  is  $2 \sum_{k=1}^i q_k + 2i - 1$  for  $1 \leq i \leq m - 1$ . Thus the set of edge labels of  $G$  is  $\{1, 3, \dots, 2q_1 - 1, 2q_1 + 1, 2q_1 + 3, \dots, 2q_1 + 2q_2 + 1, 2q_1 + 2q_2 + 3, \dots, 2 \sum_{k=1}^m q_k + 2m - 3\}$ . Hence  $f$  is a strongly harmonious odd labeling.  $\square$

**Theorem 2.10.** *Let  $G_1(p_1, q_1)$  be a strongly odd harmonious graph and  $G_2(p_2, q_2)$  be an odd harmonious graph. Then the graph  $G$  obtained by joining the vertex labeled by  $q_1$  in  $G_1$  with the vertex labeled by 0 in  $G_2$  is also odd harmonious graph.*

*Proof.* The graph  $G$  has  $q = q_1 + q_2 + 1$  edges. Let  $f_1$  be the strongly odd harmonious labeling of  $G_1$  and  $f_2$  be the odd harmonious labeling of  $G_2$ . Define a labeling  $f : V(G) \rightarrow \{0, 1, 2, \dots, q\}$  by  $f(x) = \begin{cases} f_1(x) & \text{if } x \in V(G_1) \\ f_2(x) + q_1 + 1 & \text{if } x \in V(G_2). \end{cases}$  Since  $f_1$  and  $f_2$  are injective functions and  $0 \leq f_1(x) \leq q_1$  and  $0 \leq f_2(x) \leq 2q_2 - 1$ ,  $f$  is injective and  $0 \leq f(x) \leq 2q_1 + 2q_2 - 1$ . The edge labels of  $G_1$  are remain constant and the edge labels  $G_2$  are increased by  $2q_1 + 2$ . Hence the edge labels are  $\{1, 3, \dots, 2q_1 - 1, 2q_1 + 1, 2q_1 + 3, 2q_1 + 5 \dots, 2q_1 + 2q_2 + 1\}$ . Thus,  $f$  is an odd harmonious labeling of  $G$ .  $\square$

**Illustration 9.** *The odd harmonious labeling of path union of 5 cycles  $C_8$  is shown in Figure 9*




Figure 9: Odd harmonious labeling of path union of 5 cycles  $C_8$

**Theorem 2.11.** *The banana tree  $BT_k(n)$  is odd harmonious for all the values of  $n \geq 1$  and  $k \geq 1$*

*Proof.* Consider  $k$  disjoint stars  $K_{1,n}$ . Let  $v$  be a new vertex and  $BT_k(n)$  the banana tree obtained by joining  $v$  to one pendant vertex of each star. Let  $V(BT_k(n)) = \{a_i, v_{ij}, v | 1 \leq i \leq k \text{ and } 1 \leq j \leq n\}$  and  $E(BT_k(n)) = \{av_{ij} | 1 \leq j \leq k \text{ and } 1 \leq j \leq n\} \cup \{vv_{in} | i = 1, 2, \dots, k\}$ .

We define  $f : V(G) \rightarrow \{0, 1, 2, \dots, 2(nk + k + 1) - 1\}$  as follows:  $f(v) = 0$  and  $f(v_{in}) = 2i - 1, i = 1, 2, 3, \dots, k$

$$f(a_i) = 2(2k - 2i + 1), i = 1, 2, 3, \dots, k$$

$$f(v_{ij}) = 2i - 1 + 2jk, i = 1, 2, 3, \dots, k \text{ and } j = 1, 2, 3, \dots, (n - 1).$$

The induced edge labelings are as follows:  $f^*(vv_{in}) = 2i - 1, i = 1, 2, 3, \dots, k$


$$f^*(a_i v_{in}) = 4k - 2i + 1, i = 1, 2, 3, \dots, k$$

$$f^*(a_i v_{ij}) = 4k + 2jk - 2i + 1, i = 1, 2, 3, \dots, k \text{ and } j = 1, 2, 3, \dots, (n - 1).$$

In view of the above defined labeling pattern  $f$  is an odd harmonious labeling for  $BT_k(n)$ . Hence  $BT_k(n)$  is an odd harmonious graph for all values of  $n \geq 1$  and  $k \geq 1$ .

□

**Illustration 10.** *The odd harmonious labeling of  $BT_3(5)$  is shown in Figure 10*

Figure 10: Odd harmonious labeling of  $BT_3(5)$ 

### References

- [1] M. E. Abdel-Aal, Odd Harmonious Labelings of Cyclic Snakes, *International Journal on applications of Graph Theory in Wireless Adhoc networks and Sensor Networks*, 5 (3) (2013) 1–13.
- [2] M. E. Abdel-Aal, New Families of Odd Harmonious Graphs, *International Journal of Soft Computing, Mathematics and Control*, 3(1) (2014) 1–13.
- [3] J. A. Gallian, A dynamic Survey of Graph Labeling, *The Electronics Journal of Combinatorics*, 17 (2014) #DS6.
- [4] R. L. Graham and N. J. A Sloane, On Additive bases and Harmonious Graphs, *SIAM J. Algebr. Disc. Meth.*, 4(1980), 382–404.
- [5] A. Gusti Saputri, K. A. Sugeng and D. Froncek, The Odd Harmonious Labeling of Dumbbell and Generalized Prism Graphs, *AKCE Int. J. Graphs Comb.*, 10 (2) (2013), 221–228.
- [6] F. Harary, *Graph Theory*, Addison-Wesley, Massachusetts, 1972.
- [7] Z. Liang, Z. Bai, On the Odd Harmonious Graphs with Applications, *J. Appl. Math. Comput.*, 29 (2009), 105–116.
- [8] P. Selvaraju, P. Balaganesan and J.Renuka, Odd Harmonious Labeling of Some Path Related Graphs, *International J. of Math.Sci. & Engg.Appls.*, 7(III)(2013), 163–170.
- [9] S. K. Vaidya and N. H. Shah, Some New Odd Harmonious Graphs, *International Journal of Mathematics and Soft Computing*, 1(2011), 9–16.

- [10] S. K. Vaidya, N. H. Shah, Odd Harmonious Labeling of Some Graphs , *International J.Math. Combin.*, 3(2012), 105–112.

P. Jeyanthi  
Research Centre  
Department of Mathematics  
Govindammal Aditanar College for Women  
Tiruchendur - 628 215  
Tamil Nadu, India  
*E-mail:* jeyajeyanthi@rediffmail.com

S. Philo  
Department of Mathematics  
PSN College of Engineering and Technology, Melathediyoor  
Tirunelveli, Tamil Nadu, India  
*E-mail:* lavernejudia@gmail.com

Kiki A. Sugeng  
Department of Mathematics  
Faculty of Mathematical and Natural Sciences  
Universitas Indonesia, Depok 16424  
Indonesia  
*E-mail:* kiki@ui.ac.id