

SUT Journal of Mathematics  
Vol. 42, No. 2 (2006), 295–303

## Quasi-conformally flat manifolds satisfying certain condition on the Ricci tensor

U. C. De and Yoshio Matsuyama

(Received September 22, 2006)

**Abstract.** The object of the present paper is to study a non-flat quasi-conformally flat Riemannian manifold whose Ricci tensor  $S$  satisfies the condition  $S(X, Y) = \gamma T(X)T(Y)$ , where  $\gamma$  is the scalar curvature and  $T$  is a 1-form defined by  $T(X) = g(X, \xi)$ ,  $\xi$  is a unit vector field.

*AMS 2000 Mathematics Subject Classification.* 53C25.

*Key words and phrases.* Quasi-conformally flat manifold, concircular vector field.

### §1. Introduction

The notion of a quasi-conformal curvature tensor was given by Yano and Sawaki [10]. According to them a quasi-conformal curvature tensor  $C^*$  is defined by

$$C^*(X, Y)Z = aR(X, Y)Z + b[S(Y, Z)X - S(X, Z)Y + g(Y, Z)QX - g(X, Z)QY] - \frac{\gamma}{n} \left[ \frac{a}{n-1} + 2b \right] [g(Y, Z)X - g(X, Z)Y], \quad (1.1)$$

where  $a$  and  $b$  are constants and  $R$ ,  $Q$  and  $\gamma$  are the Riemannian curvature tensor of type (1, 3), the Ricci operator defined by  $g(QX, Y) = S(X, Y)$  and the scalar curvature, respectively. If  $a = 1$  and  $b = -\frac{1}{n-2}$ , then (1.1) takes the

---

The work was done when the first author visited Chuo University during 1-30 June, 2006 as a visiting Professor.

form

$$\begin{aligned} C^*(X, Y)Z &= R(X, Y)Z - \frac{1}{n-2}[S(Y, Z)X - S(X, Z)Y + g(Y, Z)QX \\ &\quad - g(X, Z)QY] + \frac{\gamma}{(n-1)(n-2)}[g(Y, Z)X - g(X, Z)Y] \\ &= C(X, Y)Z, \end{aligned}$$

where  $C$  is the conformal curvature tensor [4]. Thus the conformal curvature tensor  $C$  is a particular case of the tensor  $C^*$ . For this reason  $C^*$  is called the quasi-conformal curvature tensor. A manifold  $(M^n, g)$  ( $n > 3$ ) shall be called quasi-conformally flat if  $C^* = 0$ . It is known [1] that a quasi-conformally flat manifold is either conformally flat if  $a \neq 0$  or Einstein if  $a = 0$  and  $b \neq 0$ . Since they give no restrictions for manifolds if  $a = 0$  and  $b = 0$ , it is essential for us to consider the case of  $a \neq 0$  or  $b \neq 0$ .

A Riemannian manifold of quasi-constant curvature was given by B. Y. Chen and K. Yano [3] as a conformally flat manifold with the curvature tensor  $\tilde{R}$  of type (0, 4) satisfies the condition

$$\begin{aligned} \tilde{R}(X, Y, Z, W) &= p[g(Y, Z)g(X, W) - g(X, Z)g(Y, W)] \\ &\quad + q[g(X, W)T(Y)T(Z) + g(Y, Z)T(X)T(W) \\ &\quad - g(X, Z)T(Y)T(W) - g(Y, W)T(X)T(Z)], \end{aligned} \quad (1.2)$$

where  $\tilde{R}(X, Y, Z, W) = g(R(X, Y)Z, W)$ ,  $R$  is the curvature tensor of type (1, 3),  $p, q$  are scalar functions and  $T$  is a non-zero 1-form defined by

$$g(X, \tilde{\xi}) = T(X), \quad (1.3)$$

where  $\tilde{\xi}$  is a unit vector field. It can be easily seen that if the curvature tensor  $\tilde{R}$  is of the form (1.2), then the manifold is conformally flat. On the other hand, G. Vrăncăanu [8] defined the notion of almost constant curvature by the same expression (1.2). Later A. L. Mocanu [6] pointed out that the manifold introduced by Chen and Yano and the manifold introduced by Vrăncăanu are the same. Hence a Riemannian manifold is said to be of quasi-constant curvature if the curvature tensor  $\tilde{R}$  satisfies the relation (1.2). If  $q = 0$ , then the manifold reduces to a manifold of constant curvature.

The present paper deals with the quasi-conformally flat manifold  $(M^n, g)$  ( $n > 3$ ) whose Ricci tensor  $S$  satisfies

$$S(X, Y) = \gamma T(X)T(Y), \quad (1.4)$$

where  $T$  is a non-zero 1-form defined by  $g(X, \xi) = T(X)$ ,  $\xi$  is a unit vector field. For the scalar curvature  $\gamma$  we suppose that  $\gamma \neq 0$  for each point of

$M$ . Under the assumption above we know that  $M$  is not Einstein. Hence we consider the case of  $a \neq 0$  (See §3). We shall prove the following:

**Theorem 1.** *A quasi-conformally flat manifold satisfying the condition (1.4) under the assumption of  $\gamma \neq 0$  is a manifold of quasi-constant curvature.*

**Theorem 2.** *In a quasi-conformally flat Riemannian manifold satisfying the condition (1.4) under the same assumption as Theorem 1, the integral curves of the vector field  $\xi$  are geodesic.*

**Theorem 3.** *In a quasi-conformally flat manifold satisfying (1.4) under the same assumption as Theorem 1, the vector field  $\xi$  is a proper concircular vector field (See §4).*

**Theorem 4.** *If a quasi-conformally flat manifold satisfies (1.4) under the same assumption as Theorem 1, then the manifold is a locally product manifold.*

**Theorem 5.** *A quasi-conformally flat manifold satisfying (1.4) under the same assumption as Theorem 1 can be expressed as a locally warped product  $I \times_{e^a} M^*$  where  $M^*$  is an Einstein manifold (See §4).*

### §2. Preliminaries

From (1.1) we obtain

$$\begin{aligned} (\nabla_W C^*)(X, Y)Z &= a(\nabla_W R)(X, Y)Z + b[(\nabla_W S)(Y, Z)X - (\nabla_W S)(X, Z)Y \\ &\quad + g(Y, Z)(\nabla_W Q)(X) - g(X, Z)(\nabla_W Q)(Y)] \\ &\quad - \frac{d\gamma(W)}{n} \left[ \frac{a}{n-1} + 2b \right] [g(Y, Z)X - g(X, Z)Y], \end{aligned} \tag{2.1}$$

where  $\nabla$  is the covariant differentiation with respect to the Riemannian metric  $g$ . We know that  $(\operatorname{div} R)(X, Y)Z = (\nabla_X S)(Y, Z) - (\nabla_Y S)(X, Z)$ . Hence contracting (2.1) we obtain

$$\begin{aligned} (\operatorname{div} C^*)(X, Y)Z &= (a + b)((\nabla_X S)(Y, Z) - (\nabla_Y S)(X, Z)) \\ &\quad + \frac{1}{n} \left[ \frac{(n-4)b}{2} - \frac{a}{n-1} \right] (g(Y, Z)d\gamma(X) - g(X, Z)d\gamma(Y)). \end{aligned} \tag{2.2}$$

Here we consider quasi-conformally flat manifold i.e.,  $C^* = 0$ . Hence  $\operatorname{div} C^* = 0$ , where 'div' denotes the divergence. If  $a + b \neq 0$ , then from (2.2) it follows that

$$\begin{aligned} &(\nabla_X S)(Y, Z) - (\nabla_Y S)(X, Z) \\ &= \frac{1}{n(a+b)} \left[ \frac{a}{n-1} - \frac{(n-4)b}{2} \right] [g(Y, Z)d\gamma(X) - g(X, Z)d\gamma(Y)]. \end{aligned}$$

This can be written as

$$(\nabla_X S)(Y, Z) - (\nabla_Y S)(X, Z) = \alpha[g(Y, Z)d\gamma(X) - g(X, Z)d\gamma(Y)], \quad (2.3)$$

where  $\alpha = \frac{1}{n(a+b)} \left[ \frac{a}{n-1} - \frac{(n-4)b}{2} \right] = \text{constant}$ .

### §3. Quasi-conformally flat manifold satisfying the condition (1.4)

From (1.1) we get

$$\begin{aligned} \tilde{C}^*(X, Y, Z, W) &= a\tilde{R}(X, Y, Z, W) + b[S(Y, Z)g(X, W) - S(X, Z)g(Y, W) \\ &\quad + S(X, W)g(Y, Z) - S(Y, W)g(X, Z)] \\ &\quad - \frac{\gamma}{n} \left[ \frac{a}{n-1} + 2b \right] [g(Y, Z)g(X, W) - g(X, Z)g(Y, W)]. \end{aligned} \quad (3.1)$$

If the manifold is quasi-conformally flat under the assumption of  $\gamma \neq 0$ , then we get

$$\gamma(a + (n-2)b) = 0.$$

Then we note that  $\left[ \frac{(n-4)b}{2} - \frac{a}{n-1} \right] = \frac{3na}{2(n-1)(n-2)}$ . Since  $a \neq 0$  under the assumption of  $\gamma \neq 0$ , we know that  $a + b \neq 0$  and  $\alpha \neq 0$ . Moreover, from (1.4) we have

$$\begin{aligned} \tilde{R}(X, Y, Z, W) &= \frac{b}{a} [S(X, Z)g(Y, W) - S(Y, Z)g(X, W) + S(Y, W)g(X, Z) - S(X, W)g(Y, Z)] \\ &\quad + \frac{\gamma}{na} \left[ \frac{a}{n-1} + 2b \right] [g(Y, Z)g(X, W) - g(X, Z)g(Y, W)] \end{aligned} \quad (3.2)$$

Using (1.4) in (3.2), we obtain

$$\begin{aligned} \tilde{R}(X, Y, Z, W) &= \frac{\gamma b}{a} [g(Y, W)T(X)T(Z) - g(X, W)T(Y)T(Z) + g(X, Z)T(Y)T(W) \\ &\quad - g(Y, Z)T(X)T(W)] + \frac{\gamma}{na} \left[ \frac{a}{n-1} + 2b \right] [g(Y, Z)g(X, W) - g(X, Z)g(Y, W)], \end{aligned}$$

which implies that the manifold is a manifold of quasi-constant curvature. Hence we can state that

**Theorem 1.** *A quasi-conformally flat manifold satisfying the condition (1.4) under the assumption of  $\gamma \neq 0$  is a manifold of quasi-constant curvature.*

**§4. The results concerning the product manifold**

From (1.4) we have

$$\begin{aligned} (\nabla_Z S)(X, Y) &= d\gamma(Z)T(X)T(Y) + \gamma[(\nabla_Z T)(X)T(Y) + T(X)(\nabla_Z T)(Y)]. \end{aligned} \quad (4.1)$$

Substituting (4.1) in (2.3), we get

$$\begin{aligned} &d\gamma(Z)T(X)T(Y) + \gamma[(\nabla_Z T)(X)T(Y) + T(X)(\nabla_Z T)(Y)] \\ &\quad - d\gamma(X)T(Z)T(Y) - \gamma[(\nabla_X T)(Z)T(Y) + T(Z)(\nabla_X T)(Y)] \\ &= \alpha[g(X, Y)d\gamma(Z) - g(Z, Y)d\gamma(X)]. \end{aligned} \quad (4.2)$$

Putting  $Y = Z = e_i$  in the above expression where  $\{e_i\}$  is an orthonormal basis of the tangent space at each point of the manifold and taking summation over  $i$ ,  $1 \leq i \leq n$ , we get

$$\alpha(1 - n)d\gamma(X) = d\gamma(\xi)T(X) + \gamma(\nabla_\xi T)(X) + \gamma T(X)(\delta T) - d\gamma(X), \quad (4.3)$$

where we put  $\delta T = \sum_{i=1}^n (\nabla_{e_i} T)(e_i)$ . Again putting  $Y = Z = \xi$  in (4.2), it yields

$$\gamma(\nabla_\xi T)(X) = (\alpha - 1)[d\gamma(\xi)T(X) - d\gamma(X)]. \quad (4.4)$$

Substituting (4.4) in (4.3), we get

$$\alpha(n - 2)d\gamma(X) - \alpha d\gamma(\xi)T(X) + \gamma\delta T = 0. \quad (4.5)$$

Now putting  $X = \xi$  in (4.5), it yields

$$\alpha(n - 3)d\gamma(\xi) + \gamma\delta T = 0. \quad (4.6)$$

From (4.5) and (4.6) it follows that

$$\alpha d\gamma(X) = \alpha d\gamma(\xi)T(X).$$

Since  $\alpha \neq 0$ , we have

$$d\gamma(X) = d\gamma(\xi)T(X). \quad (4.7)$$

Putting  $Y = \xi$  in (4.2) and using (4.7), we obtain

$$(\nabla_X T)(Z) - (\nabla_Z T)(X) = 0, \quad (4.8)$$

since  $\gamma \neq 0$ . This means that the 1-form  $T$  defined by  $g(X, \xi) = T(X)$  is closed, i.e.,  $dT(X, Y) = 0$ . Hence it follows that

$$g(\nabla_X \xi, Y) = g(\nabla_Y \xi, X) \quad (4.9)$$

for all  $X, Y$ . Now putting  $Y = \xi$  in (4.9), we get

$$g(\nabla_X \xi, \xi) = g(\nabla_\xi \xi, X). \quad (4.10)$$

Since  $g(\nabla_X \xi, \xi) = 0$ , from (4.10) it follows that  $g(\nabla_\xi \xi, X) = 0$  for all  $X$ . Hence  $\nabla_\xi \xi = 0$ . This means that the integral curves of the vector field  $\xi$  are geodesic. Therefore we can state the following:

**Theorem 2.** *In a quasi-conformally flat Riemannian manifold satisfying the condition (1.4) under the assumption of  $\gamma \neq 0$ , the integral curves of the vector field  $\xi$  are geodesic.*

From (4.4), by virtue of (4.7) we get

$$(\nabla_\xi T)(Z) = 0, \quad (4.11)$$

since  $\gamma \neq 0$ . Now we consider the scalar function

$$f = \alpha \frac{d\gamma(\xi)}{\gamma}.$$

We have

$$\nabla_X f = \frac{\alpha}{\gamma^2} [d\gamma(\xi)T(\nabla_X \xi)\gamma - d\gamma(X)d\gamma(\xi)] + \frac{\alpha}{\gamma} d^2\gamma(\xi, X), \quad (4.12)$$

where the Hessian  $d^2\gamma$  is defined by  $d^2\gamma(X, Y) = X(Y\gamma) - (\nabla_X Y)\gamma$ . On the other hand, (4.7) implies that

$$d^2\gamma(Y, X) = d^2\gamma(\xi, Y)T(X) + d\gamma(\xi)T(\nabla_Y \xi)T(X) + d\gamma(\xi)(\nabla_Y T)(X),$$

from which we get

$$d^2\gamma(\xi, Y)T(X) = d^2\gamma(\xi, X)T(Y), \quad (4.13)$$

since  $(\nabla_X T)(Y) = (\nabla_Y T)(X)$  and  $d^2\gamma(Y, X) = d^2\gamma(X, Y)$ . Putting  $X = \xi$  in (4.13), it follows that

$$d^2\gamma(\xi, Y) = d^2\gamma(\xi, \xi)T(Y),$$

since  $T(\xi) = 1$ . Thus

$$\nabla_X f = \mu T(X), \quad (4.14)$$

where  $\mu = \frac{\alpha}{\gamma} [d^2\gamma(\xi, \xi) - \frac{d\gamma(\xi)}{\gamma} d\gamma(\xi)]$  and we used (4.7). Using (4.14), it is easy to show that

$$\omega(X) = \frac{\alpha}{\gamma} d\gamma(\xi)T(X) = fT(X)$$

is closed. In fact,

$$d\omega(X, Y) = 0.$$

Using (4.7) and (4.8) in (4.2), we get

$$\begin{aligned} & \gamma[T(Z)(\nabla_X T)(Y) - T(X)(\nabla_Z T)(Y)] \\ &= \alpha d\gamma(\xi)[g(Y, Z)T(X) - g(X, Y)T(Z)]. \end{aligned}$$

Now putting  $Z = \xi$  in the above expression it yields

$$-(\nabla_X T)(Y) = \alpha \frac{d\gamma(\xi)}{\gamma} [T(X)T(Y) - g(X, Y)], \tag{4.15}$$

by (4.11). Thus (4.15) can be rewritten as follows:

$$(\nabla_X T)(Y) = -fg(X, Y) + \omega(X)T(Y), \tag{4.16}$$

where  $\omega$  is closed. But this means that the vector field  $\xi$  defined by  $g(X, \xi) = T(X)$  is a proper concircular vector field ([7], [9]). Hence we can state the following:

**Theorem 3.** *In a quasi-conformally flat manifold satisfying (1.4) under the assumption of  $\gamma \neq 0$ , the vector field  $\xi$  is a proper concircular vector field.*

From (4.16) it follows that

$$\nabla_X \xi = -fX + \omega(X)\xi. \tag{4.17}$$

Let  $\xi^\perp$  denote the  $(n - 1)$ -dimensional distribution in a quasi-conformally flat manifold orthogonal to  $\xi$ . If  $X$  and  $Y$  belong to  $\xi^\perp$ , then

$$g(X, \xi) = 0 \tag{4.18}$$

and

$$g(Y, \xi) = 0. \tag{4.19}$$

Since  $(\nabla_X g)(Y, \xi) = 0$ , it follows from (4.17) and (4.19) that

$$g(\nabla_X Y, \xi) = g(\nabla_X \xi, Y) = -fg(X, Y).$$

Similarly, we get

$$g(\nabla_Y X, \xi) = g(\nabla_Y \xi, X) = -fg(X, Y).$$

Hence

$$g(\nabla_X Y, \xi) = (\nabla_Y X, \xi). \tag{4.20}$$

Now  $[X, Y] = \nabla_X Y - \nabla_Y X$  and therefore by (4.20) we obtain

$$g([X, Y], \xi) = g(\nabla_X Y - \nabla_Y X, \xi) = 0.$$

Hence  $[X, Y]$  is orthogonal to  $\xi$ . That is,  $[X, Y]$  belongs to  $\xi^\perp$ . Thus the distribution  $\xi^\perp$  is involutive [2]. Hence from Frobenius' theorem [2] it follows that  $\xi^\perp$  is integrable. This implies that if a quasi-conformally flat manifold satisfies (1.4), then it is a product manifold. We can therefore state the following theorem:

**Theorem 4.** *If a quasi-conformally flat manifold satisfies (1.4) under the assumption of  $\gamma \neq 0$ , then the manifold is a locally product manifold.*

If a quasi-conformally flat manifold satisfies (1.4) under the assumption of  $\gamma \neq 0$ , then in view of Theorem 3,  $\xi$  is a concircular vector field. Also,  $M$  is a quasi-constant curvature manifold and satisfies (1.2) and from Theorem 4 we know that  $\xi^\perp$  is integrable and it holds

$$g(\nabla_X Y, \xi) = -(\nabla_X T)(Y)$$

for the local vector fields  $X, Y$  belonging to  $\xi^\perp$ . Thus from (4.15) the second fundamental form  $k$  for each leaf satisfies

$$k(X, Y) = -\alpha \frac{d\gamma(\xi)}{\gamma} g(X, Y)\xi.$$

Hence we know that each leaf is totally umbilic. Therefore each leaf is a manifold of constant curvature. Hence it must be a warped product  $I \times_{e^q} M^*$  where  $M^*$  is an Einstein manifold. Thus we can state the following result (See [9], [5]):

**Theorem 5.** *A quasi-conformally flat manifold satisfying (1.4) under the assumption of  $\gamma \neq 0$  can be expressed as a locally warped product  $I \times_{e^q} M^*$  where  $M^*$  is an Einstein manifold.*

## References

- [1] AMUR, K. and MARALABHAVI, Y. B., *On quasi-conformal flat spaces*, Tensor, N. S. **31** pp. 194-198 (1977).
- [2] BRICKELL, F and CLARK, R. S., *Differentiable manifold*, Van Nostrand Reinhold Comp. London (1978).
- [3] CHEN, B. Y. and YANO, K., *Hypersurfaces of a conformally flat space*, Tensor, N. S. **26** pp. 318-322 (1972).



- [4] EISENHART, L. P., *Riemannian Geometry*, Princeton University Press (1949).
- [5] GEBAROWSKI, A., *Nearly conformally symmetric warped product manifolds*, Bulletin of the Inst. of Math. Academia Sinica **4** pp.359-371 (1992).
- [6] MOCANU, A. L., *Les variétés à courbure quasi-constant de type Vranceanu*, Proc. Nat. Conf. Geom. and Top., Tirgovisté (1987).
- [7] SCHOUTEN, J. A., *Ricci-Calculus*, Springer, Berlin (1954).
- [8] VRÂNCEANU, G. *Leçons des Geometrie Differential*, Vol. 4, Ed. de l'Academie, Bucharest (1968).
- [9] YANO, K., *Concircular Geometry I*, Proc. Imp. Acad. Tokyo **16** pp. 195-200 (1940).
- [10] YANO, K. and SAWAKI, S., *Riemannian manifolds admitting a conformal transformation group*, J. Diff. Geom. **2** pp.161-184 (1968).

U. C. De  
Department of Mathematics, University of Kalyani,  
Kalyani 741235, W. B., India  
*E-mail*: uc\_de@yahoo.com

Yoshio Matsuyama  
Department of Mathematics, Chuo University,  
Kasuga, Bunkyo-ku, Tokyo 112-8551, Japan  
*E-mail*: matuyama@math.chuo-u.ac.jp