

6-1952

Thallium dithizonate

Bruno Jaselskis

Union College - Schenectady, NY

Follow this and additional works at: <https://digitalworks.union.edu/theses>

 Part of the [Chemistry Commons](#)

Recommended Citation

Jaselskis, Bruno, "Thallium dithizonate" (1952). *Honors Theses*. 1914.
<https://digitalworks.union.edu/theses/1914>

This Open Access is brought to you for free and open access by the Student Work at Union | Digital Works. It has been accepted for inclusion in Honors Theses by an authorized administrator of Union | Digital Works. For more information, please contact digitalworks@union.edu.

THALLIUM DITHIZONATE

by

Bruno Jaselskis

A thesis presented to the Department of Chemistry of
Union College in partial fulfillment of the requirements
for the degree of Bachelor of Science in Chemistry.

By Bruno Jaselskis

Approved by Galen W. Ewing

May 24, 1952

UNION COLLEGE
LIBRARY

UN92

J39t

1952

c.2

Attempts to establish a useful dithizone method in the determination of thallos ion have been carried on at Union College by Snyder (19). Because the previous work had not produced satisfactory results and because in general thallos dithizonates have not been studied, the following work, under Dr. W.G. Ewing's guidance, was undertaken.

Since dithizonates could be of keto or enol forms, or the mixture of both of them as will later be explained, Job's continuous variation method appeared to be applicable. In order to check the relative amount of thallos ion in thallos dithizonate, polarographic work seemed to be adequate. In addition, a few other methods were used as recommended by Mellen (10). Because the dithizone is soluble in strongly polar organic solvents, e. g. chloroform, carbon tetrachloride etc. but thallos carbonate is soluble in water, two phases had to be used and this caused serious inconvenience.

For advice and encouragement, special thanks are to be expressed to Dr. W. G. Ewing and to all the Chemistry Department staff at Union College, who in one way or the other made this work easier.

158356



11 Nov 52

Dept of Chemistry

Emil Fischer prepared diphenylthiocarbozone, common name dithizone $S=C \begin{matrix} H & H \\ | & | \\ N & - & N & - & C_6H_5 \\ | & & | \\ N & = & N & - & C_6H_5 \end{matrix}$ at the end of XIX century (3). He also noticed that the dithizone gave highly colored products with heavy metals. Hellmut Fischer (4) continued investigation and found that dithizone gave colored complexes even at very low concentration of the metal. But selectivity due to the similar colors was rather poor. Other chemists found that by proper choice of pH the selectivity could be increased because heavy metal dithizonates were formed at different pH's, and their solubilities in the organic phase varied markedly.

From a theoretical point of view, dithizone may be considered as a weak dibasic acid with much stronger monobasic acid character. The monobasic acid is prevalent up to pH 12 as it is observed by Buch (1).

Considering that dithizone dissociates in the following way :



(For convenience dithizone is written as H_2Dz and dithizonate ion as HDz^- or Dz^{2-})

The dissociation constant for the first reaction is:

$$K_d = \frac{[H^+]_w \cdot [HDz^-]_w}{[H_2Dz]_w} \dots 1$$

where subscript "o" stands for the organic phase and "w" for the aqueous phase.

The partition coefficient (P) between the organic phase and water is:


$$P_d = \frac{[H_2Dz]_o}{[H_2Dz]_w} \dots 2$$

Considering that H^+ and $[H_2Dz]_w$ are constant and small following is true:

$$\frac{[H_2Dz]_o}{[HDz^-]_w} = \frac{P_d}{K_d} [H^+]_w$$

The value of $\frac{P_d}{K_d} = F = 3.6 \times 10^0$ as computed by Irving and coworkers (7) and mentioned by Sandell (12).

If the metal dithizonate dissociates



then the equilibrium constant is:

$$K_m = \frac{[Me^+]_w [HDz^-]_w}{[MeHDz]_w}$$


and the partition coefficient is:

$$P_m = \frac{[MeHDz]_o}{[MeHDz]_w}$$


Dividing equation 5 by equation 4 and substituting equation 3 for $\frac{[H_2Dz]_o}{[HDz^-]_w}$ the following expression is obtained:

$$\frac{[MeHDz]_o}{[Me^+]_w} = \frac{P_m}{K_m} \frac{[H_2Dz]_o}{F [H^+]_w}$$

Metal dithizonates formed by monobasic dithizone ion are known as keto complexes and are formed by all heavy metals which react with dithizone. Enol type complexes are easily formed by copper, mercury, and silver, but even so at higher pH and in the deficiency of dithizone (13). The structural formulas assigned by H. Fisher (5) are:




Keto metal
dithizonate



Enol metal
dithizonate

Dithizone is easily oxidized to diphenylthiocarbodiazone



Presumably the fading of complexes may be due to this oxidation.


The French Chemist P. Job (8), (9) successfully applied the colorimetric method in the study of complexes. Most of his publications appeared in the nineteen twenties in France. His work provoked some interest in the other countries. In the United States, Vosburgh & Cooper (21) studied the theoretical aspects and practical applications of the Job's continuous variation method in the investigation of the stability of complexes and their coordination numbers.

According to Job's continuous variation method, two reagents A and B, both M molar, are mixed in varying proportions, so that the concentration of A is dependant upon B as:

$$A = M(1-x) \dots\dots\dots 1 \quad \text{where } x \text{ is the volume of B}$$

$$B = Mx \quad \dots\dots\dots 2 \quad \text{and } x \leq 1$$

At equilibrium this reaction may be expressed as:



If the concentration of A, B, and AB_n are respectively c_1, c_2, c_3 , than at equilibrium:

$$c_1 = M(1-x) - c_3 \quad \dots\dots\dots 4$$

$$c_2 = Mx - nc_3 \quad \dots\dots\dots 5$$

$$Kc_3 = c_1 \cdot c_2^n \quad \dots\dots\dots 6$$

$$c_3 \text{ is at a maximum when } \frac{dc_3}{dx} = 0 \quad \dots\dots\dots 7$$

Differentiating equations 4, 5, 6 with respect to x and combining the resulting equations with 4 and 7 the coordination number "n" is obtained:

$$n = \frac{x}{1-x} \quad \text{-----8}$$

In order to show that a minimum or a maximum in the absorption of monochromatic light occurs when C₃ is at the maximum, consider that a_{m1}; a_{m2}; a_{m3} are the molar absorptancy indices of reagents and product at a given wave length. Then the absorbance (A) is:


$$A = l (a_{m1} C_1 + a_{m2} C_2 + a_{m3} C_3) \quad \text{Where "l" is the thickness of the cell.}$$

Denoting γ as the difference between the absorbance observed and the absorbance of an identical solution if the reaction has not taken place. The following value is obtained:

$$\gamma = l [a_{m1} C_1 + a_{m2} C_2 + Ma_{m1} (1-x) - Ma_{m2} x] \quad \text{-----9}$$

By differentiating this equation with respect to "x" it may be shown that "y" occurs at a maximum when C₃ is at the maximum provided that a_{m3} > a_{m1}; and "y" is at a minimum when C₃ is at a maximum if a_{m3} < a_{m1}. Consequently, as a maximum or minimum in y will occur at the same value of "x" for various wave lengths, if there is only one complex formed.

In case a second complex is formed the equilibrium reaction is:



When the equilibrium constant is:

$$K = \frac{C_3 C_2^q}{C_4} \quad \text{-----10}$$

Where C₄ is the concentration of AB (n ≠ q)

It may be shown that the coordination number "n" for this complex is:

$$n'' = \frac{x}{1-x} + \frac{v(v+n)}{M(1-x)} \cdot C_4^{0.4} \quad \text{-----11}$$

In case there is only one complex formed or if the amount of the second complex is so small that it may be neglected and the maximum of C_3 occurs at $x = 0,5$ the slopes at $x = 0$ also at $x = 1$ are the same.

$$\Delta = \left(\frac{dA}{dx}\right)_{x=0} = -\left(\frac{dA}{dx}\right)_{x=1}$$

Then the absorbance (A) observed is related to the molar absorptivity index of the complex formed

$$A = a_{m3} \left[\frac{C+1/K}{2} - \left(\left[\frac{C+1/K}{2} \right]^2 - C^2 x(1-x) \right)^{1/2} \right] \quad \text{..... 12}$$

as shown by Schwarzenbach (18) and Treadwell (20).

When $x = 0$ the expression of A becomes

$$\Delta = a_{m3} \frac{Kc^2}{Kc+1} \quad \text{..... 13}$$

Where C is the molar concentration of B at a given x value solving equations 12, and 13 simultaneously, we obtain the value of K - the equilibrium constant and also the value of the molar absorptivity index for the complex.

In this study of thallos dithizonates, it was decided to analyze only the organic phase. On account of this the computed value of the equilibrium constant of thallos dithizonate in the chloroform solution is only proportional to the dissociation constant K_m as evaluated from the following equation:

$$\frac{[MeHOZ]_0}{[Me^+]_w} = \frac{P_m [H_2OZ]_0}{K_m F [H^+]_w}$$

In determination of the absorbances and the relative amounts of thalious ion in thalious dithizonates the following apparatus has been used:

Beckman Model D U Photoelectric Quartz Spectrophotometer, serial no. 3210; during the absorbance measurements sensitivity was kept constant, only the slit opening was varied.

Electropode manufactured by Fisher Scientific Co. Roller-Smith, precision balance.

The materials used were:

Dithizone purchased from Eastman Kodak Co., original purity 84%. After purification according to the procedure described by Sandell (16) purity of dithizone was about 93-94%. The actual dithizone concentration was determined by measuring the absorbance of a chloroform solution of dithizone at the wave lengths 440, 505, 606 $m\mu$ and using Cooper & Sullivan (2) values of molar absorbancy indices as illustrated by the table I, Graph 1.

TABLE I

Mg of (H ₂ DZ)	Dissolved in ml CHCl ₃	Absorbance					Av. Conc. of (H ₂ DZ)	Purity %
		440	505	510	520	605		
2.00	100	1.030	.401				6.55×10^{-5}	84.6
6.70	250	1.450	.539	.610	.790		9.88×10^{-5}	93.9(Av)
6.41	250	1.220	.469	.493	.629		7.90x	
5.18	250	0.990	.369	.387	.489	2.1	6.3x	
3.35	250	0.780	.300	.314	.394	1.96	4.9x	
1.34	250	0.310	.116	.124	.157	0.822	1.97x	
0.67	250	0.152	.058	.062	.078	.419	0.38x	

Molar absorbaney indices a_m for dithizone


440	m μ	16×10^3
505	m μ	6×10^3
605	m μ	41.5×10^3

Remaining impurities were probably the oxidation product of dithizone and some moisture. Since dithizone solutions decompose in light and above room temperature they were kept in the dark and in the refrigerator. Usually only the amount necessary for immediate use was prepared.

Dithizone is soluble in polar organic solvents: for the purposes of this work chloroform (CHCl₃) was employed. USP and CP grades of chloroform were used without further purification.

Thalious carbonate (Tl₂CO₃) (CP) purchased from Fairmount Chemical Co. was found by the gravimetric method described by Scott (17) to be 99.92% pure. This analysis was confirmed by the volumetric determination of thalious ion by the titration with standard ceric sulfate as recommended by Willard and co-workers (23, 24). The foregoing is illustrated

GRAPH No 1
ABSORBANCE OF DITHIZONE VERSUS
CONCENTRATION



in the data below:

	I sample	II sample
Weight of sample Tl_2CO_3	0.2344 gm.	0.2344 gm
Weight of Tl_2CrO_4	0.2609 gm.	0.2608 g
Weight of Tl in Tl_2CrO_3 samples	0.2043 gm.	0.2043 gm
Weight of Tl in Tl_2CrO_4	0.2037 gm.	0.2036 gm
Total volume of washings and solution	75 ml	75 ml

The amount of Tl in Tl_2CrO_4 must be increased by approximately 0.0004 gm. due to the loss in washing.

Therefore, the purity of thallos carbonate is 99.92% --
-- 99.91%

In volumetric determinations solutions were prepared in the following way:

Hydrochloric acid concentrated	15.25 ml
Iodine chloride (catalyst)	10.00 ml
Water	74.75 ml

Standardization of ceric sulfate with sodium oxalate:

Weight of $\text{Na}_2\text{C}_2\text{O}_4$	Ml of $\text{Ce}(\text{SO}_4)_2$	Normality of $\text{Ce}(\text{SO}_4)_2$
6.01 mg	7.94	
6.01 mg	7.92	
6.01 mg	7.95	Av. 0.1124 N

Titration of thallos carbonate:

Weight of Tl_2CO_3	Ml of $\text{Ce}(\text{SO}_4)_2$
0.2344 g ^N	8.85 ml
0.2344 g ^N	8.88 ml

Purity of the sample is 99.71%.

The buffer solutions were prepared according to the directions of Clark & Lubs (22).

The water was redistilled for the solutions used in Job's continuous variation method.

2, 3 dimercaptopropanol* was used in the recursion method. This is also known as "British anti-Lewisite" abbreviated (BAL)

The mixed color method was applied, while studying absorption of thallos dithizonates. The volumes of two phases: Chloroform and aqueous phase were kept constant.

* Obtained through the courtesy of J. E. Fagel, Jr. General Electric Research Laboratories.

TABLE II

Absorption spectra of dithizone, thalious dithizonates, thalious dithizonates with (BAL) and the decolorized (in the sun light) thalious dithizonate are illustrated in 2 following graphs.

2A and 2B.

<u>Wave Length</u> m μ	(a)	(b)	<u>Absorbance</u> (c)	(d)	(e)	(f)	(g)
250	.319	.213					.440
260	.351	.233					.460
270	.339	.226			.266	.712	.390
280	.305	.204			.232	.627	.325
290	.239	.163			.212	.482	.282
300	.196	.133			.195	.401	.257
310	.165	.113			.165	.335	.243
320	.138	.098			.167	.270	.225
330	.118	.084			.163	.216	.195
340	.108	.077			.137	.198	.150
350	.105	.075			.110	.191	.137
360	.102	.074			.100	.181	.125
370	.096	.071			.107	.157	.113

<u>Wave Length</u> m μ	<u>Absorbancy</u>						
	(a)	(b)	(c)	(d)	(e)	(f)	(g)
380	.090	.067			.124	.126 .23	.107
390	.087	.065			.156	.106	.099
400	.089	.067			.200	.100	.093
410	.098	.078			.244	.111	.087
420	.111	.083			.285	.142	.085
430	.131	.094		.500	.325	.193	.077
440	.157	.110	.548	.530	.328	.262	.075
450	.188	.128	.737	.560	.320	.350	.069
460	.225	.148	.980	.550	.294	.458	.063
470	.267	.172	1.220	.530	.261	.588	.054
480	.311	.195	1.360	.510	.218	.700	.048
490	.349	.217	1.580	.480	.172	.800	.091
500	.384	.239	1.680	.458	.147	.802	.034
505			1.710	.468			
508			1.720				
510	.399	.247	1.721		.147	.878	.030
512			1.680				

<u>Wave Length</u> m μ	<u>Absorbance</u>						
	(a)	(b)	(c)	(d)	(e)	(f)	(g)
520	.389	.245	1.610	.490	.178	.810	.026
530	.358	.230	1.350	.500	.222	.662	
540	.296	.199		.508	.275	.473	
550	.239	.168	.634	.548	.357	.302	
560	.187	.138		.622	.440	.182	.018
570	.141	.114	.257		.518	.106	
580	.116	.102	.187	.810	.590	.080	
590	.110	.098	.165	.930	.678	.063	
600	.106	.096	.158	1.020	.741	.060	
605			.156	1.050			
610	.101	.088	.150	1.000	.735	.055	
620	.086	.079	.120	.630	.050		
625				.740			
630	.068	.062			.305	.048	
640					.128	.043	
650	.038	.031	.030	.264	.075	.040	
660					.052	.038	
670							

<u>Wave Length</u> m μ	(a)	(b)	<u>Absorbance</u> (c)	(d)	(e)	(f)	(g)
675	.015	.016	.010	.058			
680					.040		
690							
700	.011	.011			.032	.028	


Absorption spectra of:

- a) Unknown thallos dithizonate 1:1
- b) " " " 1:2
- c) 1.4 mg of unknown thallos dithizonate per 50 cc of chloroform
- d) 1.4 mg of unknown thallos dithizonate per 50 cc of chloroform with (BAL)
- e) Dithizone in chloroform
- f) Unknown dithizonate in chloroform
- g) Decolorized thallos dithizonate.

GRAPH No 2A

ABSORPTION SPECTRUM OF UNKNOWN THALLOUS
DITHIZONATE

- a) reagents in 1:1 ratio
- b) Deficiency of Dithizone (1:2)
- c) 1.4 mg in 50 ml of chloroform
- d) 1.4 mg in 50 ml of chloroform
after addition of (BAL)



GRAPH No 2B

ABSORPTION SPECTRA OF

- a) Unknown thallos dithizonate
- f) Dithizone in chloroform
- g) Decolorized Thallos dithizonate




TABLE III

Graph 5

Job's Continuous variation method

pH 9.4

T 20°C

 $(1-x) = 8.5 \times 10^{-7} M$ when $x = 0$

Solutions were kept at 20° for 30-35 minutes before measuring absorbance.

Serial No.	Moles of H_2DZ	$4.25 \times 10^{-5} M$	ml $CHCl_3$	Moles of Tl^+	$5.0 \times 10^{-4} N$	ml Buffer	ml Water	Resulting Mol. Of H_2DZ in $CHCl_3$
1	6.4×10^{-7}	15.0	-	2.1×10^{-7}	0.42	8.0	1.58	4.25×10^{-5}
2	6.0x	14.1	0.9	2.5x	0.50	8.0	1.50	4.00x
3	5.5x	13.0	2.0	3.0x	0.60	8.0	1.40	3.77x
4	5.5x	13.0	2.0	3.0x	0.60	8.0	1.40	3.77x
5	5.1x	12.0	3.0	3.4x	0.68	8.0	1.32	3.40x
6	4.5x	10.6	4.4	4.0x	0.80	8.0	1.20	3.00x
7	4.25x	10.0	5.0	4.25x	0.81	8.0	1.19	2.84x
8	4.00x	9.6	5.4	4.5x	0.90	8.0	1.10	2.66x
9	3.40x	8.0	7.0	5.1x	1.02	8.0	1.08	2.26x
10	2.95x	7.0	8.0	5.55x	1.11	8.0	0.89	1.98x
11	2.55x	6.0	9.0	5.95x	1.19	8.0	0.81	1.70x
12	2.10x	5.0	10.0	6.40x	1.28	8.0	0.72	1.31×10^{-5}
13	1.26x	3.0	12.0	7.24x	1.48	8.0	0.52	8.00×10^{-6}

No.	Absorbance at 510	$\mu m C_1$	Absorbance (A) at 510 with (BAL)	Incriment
1	.648	.268 .380	.560	.088
2	.658	.250 .408		
3	.688	.237 .451	.510	.178
4	.715	.237 .477		

No.	Absorbance at 510	$a_m c_l$	γ	Absorbance (A) at 510 with (BAL)	Incriment (b)
5	.740	.213	.527	.490	.250
6	.761	.188	.573		
7	.770	.179	.591	.430	.340
8	.742	.167	.575		
9	.635	.143	.492	.371	.264
10	.568	.123	.443	.317	.251
11	.497	.108	.389	.264	.233
12	.394	.081	.313	.198	.196
13	.245	.050	.195	.092	.153

TABLE IV (Graph 5)

Job's continuous variation method

pH 9.6 ; T 20° C ; $(1-x) = 8.5 \times 10^{-7}$ when $x = 0$

Solutions were kept at 20° for 30-35 minutes before measuring the absorbance (A).

No.	Moles of H_2D_2 ml H_2D_2 ml $4.25 \times 10^{-5} M$	$CHCl_3$ ml	Moles Tl^+ of Tl^+ $5.0 \times 10^{-4} M$	Buffer ml	Water ml	Resulting mol H_2D_2 in $CHCl_3$
1	6.5×10^7 14.95	0.05	2.0×10^7 0.40	8.9	1.60	4.28×10^{-5}
2	5.5 x 12.95	2.05	3.0x 0.60	8.0	1.40	3.77x
3	4.50x 10.35	4.65	4.0x 0.80	8.0	1.20	3.00x
4	4.00x 9.40	5.60	4.5x 0.90	8.0	1.10	2.66x
5	3.50x 8.35	6.65	5.0x 1.00	8.0	1.00	2.36x
6	3.00x 7.05	7.95	5.5x 1.10	8.0	0.90	1.98x
7	2.50x 5.88	9.12	6.0x 1.20	8.0	0.80	1.66x

No.	Moles of H_2O_2	H_2O_2 4.25×10^{-5}	$CHCl_3$ ml	Moles Te^{+}	Te^{+} ml	Buffer ml	Water ml	Resulting mol. (H_2O_2)
8	1.00×10^{-7}	2.35	12.65	$7.5 \times$	1.50	8.0	0.50	0.67×10^{-5}

	Absorbance at 510 m μ	(A)	a_{mC_1}	γ
1	.620		.268	.352
2	.737		.236	.501
3	.790		.190	.600
4	.755		.168	.587
5	.688		.148	.540
6	.608		.122	.486
7	.445		.102	.343
8	.220		.041	.161

TABLE V (Graph 5)

Job's Continuous variation method

pH 9.6 ; T = 20°C ; $(1-x) = 8.5 \times 10^{-7}M$ when $x = 0$

Solutions were kept at 20°C for 30-35 minutes before measuring the absorbance (A).

No.	Moles of H_2O_2	H_2O_2 4.25×10^{-5} ml	$CHCl_3$ ml	Moles Te^{+} Te	Te^{+} 5.0×10^{-4}	Buffer ml	Water ml	Resulting mol (H_2O_2)
1	6.5×10^{-7}	14.95	0.05	2.0×10^{-7}	0.40	8.0	1.6	4.25×10^{-5}
2	$6.0x$	14.10	0.90	$2.5x$	0.50	8.0	1.5	$4.0x$
3	$5.5x$	12.95	2.05	$3.0x$	0.60	8.0	1.4	$3.77x$
4	$5.0x$	11.80	3.20	$3.5x$	0.70	8.0	1.3	$3.30x$
5	$4.50x$	10.35	4.65	$4.0x$	0.80	8.0	1.2	$2.92x$
6	$4.00x$	9.40	5.60	$4.5x$	0.90	8.0	1.1	$2.66x$
7	$3.50x$	8.35	6.65	$5.0x$	1.00	8.0	1.0	$2.34x$
8	$3.00x$	7.05	7.95	$5.5x$	1.10	8.0	0.9	$2.00x$
9	$2.50x$	5.88	9.12	$6.0x$	1.20	8.0	0.8	$1.66x$
10	$2.00x$	4.70	10.30	$6.5x$	1.30	8.0	0.7	$1.33x$
11	$1.50x$	3.55	11.45	$7.0x$	1.40	8.0	0.6	$1.00x$
12	$1.00x$	2.35	12.65	$7.5x$	1.50	8.0	0.5	$0.65x$

No.	Absorbance (A) at 510	a_{mCl}	
1	.603	.268	.335
2	.625	.250	.375
3	.727	.236	.491
4	.758	.208	.550

No.	Absorbance (A) at 510	$a_m C_1$	γ
5	.810	.190	.620
6	.764	.168	.596
7	.695	.148	.547
8	.603	.122	.481
9	.453	.102	.351
10	.383	.081	.302
11	.302	.062	.240
12	.204	.041	.163

TABLE VI (Graph 3,4)

Job's continuous variation method.

pH 10.0 T = 20°C ; $1-x = 10. x 10^{-7} M$ when $x = 0$

Solutions were kept at 20°C for 30-35 minutes before measuring absorbance (A).

No.	Moles of H_2O_2	H_2O_2 (ml) 5.9×10^{-5}	$CHCl_3$ ml	Moles Fe^{2+}	Fe^{2+} (ml) 5×10^{-4}	Buffer ml	Water ml	Resulting mol. H_2O_2 .
1	7.5×10^{-7}	12.75	1.25	2.5×10^{-7}	0.50	8.0	1.50	5.0×10^{-5}
2	6.5x	11.00	4.00	3.5x	0.70	8.0	1.30	4.34x
3	6.0x	10.20	4.80	4.0x	0.80	8.0	1.20	4.00x
4	5.5x	9.30	5.70	4.5x	0.90	8.0	1.10	3.66x
5	5.5x	9.30	5.70	4.5x	0.90	8.0	1.10	3.66x
6	5.0x	8.50	6.50	5.0x	1.00	8.0	1.00	3.35x
7	4.5x	7.60	7.40	5.5x	1.10	8.0	0.90	3.00x
8	4.5x	7.60	5.5x	1.10	1.10	8.0	0.90	5.00x

No.	Moles of H_2O_2	H_2O_2 (ml) 5.9×10^5	$CHCl_3$ ml	Moles Te^{+}	Te^{+} (ml)	Buffer ml	Water ml	Resulting Mol H_2O_2
9	4.0x	6.80	8.20	6.0x	1.20	8.0	0.80	2.66x
10	4.0x	6.80	8.20	6.0x	1.20	8.0	0.80	2.66x
11	3.0x	5.10	9.90	7.0x	1.40	8.0	0.60	2.00x
12	2.0x	3.40	11.60	8.0x	1.60	8.0	0.40	1.33x

No.	(A) 440 m μ	$a_m C_1$	Δ	(A) 505 m μ	$a_m C_1$	Δ	(A) 510 m μ	$a_m C_1$	Δ
1	.660	.770	-.110	.825	.395	.430	.757	.312	.445
2	.506	.666	-.160	.876	.255	.621	.884	.270	.614
3	.418	.612	-.194	.974	.234	.740	.980	.250	.730
4	.344	.561	-.217	.990	.217	.773	1.000	.230	.770
5	.355	.561	-.225	.982	.217	.765	.983	.230	.753
6	.330	.520	-.190	.950	.198	.752	.980	.211	.769
7	.290	.460	-.170	.840	.175	.665	.850	.188	.662
8	.298	.460	-.162	.857	.175	.682	.865	.188	.677
9	.255	.410	-.155	.810	.153	.657	.815	.167	.648
10	.249	.410	-.161	.798	.153	.645	.803	.167	.636
11	.202	.307	-.105	.690	.115	.575	.695	.125	.570
12	.132	.205	-.073	.438	.078	.360	.455	.083	.377

* (A) is absorbance observed.

(D) is the difference

No.	(A) <i>520 mμ</i>	$a_m G_1$	ψ
1	.760	.395	.365
2	.845	.343	.502
3	.936	.315	.621
4	.950	.290	.660
5	.943	.290	.653
6	.948	.265	.683
7	.820	.235	.585
8	.827	.235	.592
9	.788	.210	.578
10	.776	.210	.566
11	.675	.158	.517
12	.445	.106	.339
13			


(BAL) added ; TABLE VII (Graph 4)

	(A) <i>440 mμ</i>	D	(A) <i>505 mμ</i>	D	(A) <i>510 mμ</i>	D	(A) <i>520 mμ</i>	D
	2	.622	.316-	.390	.486			
2	.662	.116	.390	.486	.402	.482	.461	.384
3	.580	.162	.418	.556	.430	.550	.468	.468
4	.565	.220	.410	.580	.420	.580	.455	.495
6	.518	.188	.343	.607	.355	.625	.390	.558
7	.440	.150	.405	.435	.408	.442	.408	.412
9	.435	.180	.278	.532	.283	.532	.315	.473
11	.363	.161	.265	.425	.264	.431	.285	.473
12	.220	.088	.150	.288	.156	.299	.175	.270

GRAPH No 3

JOB'S CONTINUOUS VARIATION METHOD
pH 10

(Y) versus (X)
(A) versus (X)




a)	(A)	at 510 mμ
b)	(Y)	510 "
c)	(Y)	505 "
d)	(Y)	520 "
e)	(Y)	440 "

GRAPH No 4

DIFFERENCE BETWEEN ABSORBANCE
WITHOUT (BAL) AND WITH (BAL)
VERSUS (X)

- a) at 510 $m\mu$
b) 505 "
c) 520 "
d) 440 "




GRAPH No 5

JOB'S CONTINUOUS VARIATION METHOD
pH 9.4 and pH 9.6

(Y) Versus (X)
(A) Versus (X)
(D) Versus (X)

- a) (A) at 510 ~~mm~~ pH 9.6
 b) (A) at 510 " pH 9.4
 c) (Y) at 510 " pH 9.6
 d) (Y) at 510 " pH 9.4
 e) (D) at 510 " pH 9.4 (B.L.)
 Added



(D) Is difference between absorbances

TABLE VIII (Graph 6)

H₂DZ constant varying amount of thalious ion added

H₂DZ = approximately 4.25×10^{-7} moles

pH 9.4

Chloroform phase 15 ml ; aqueous phase 10 ml

	Tl Moles	Tl ml 5×10^{-4}	Absorbance at 510 <i>mμ</i>	at 440 <i>mμ</i>
1	1×10^{-7}	0.20	.338	.425
2	2x	0.43	.475	.334
3	3x	0.60	.600	.310
4	4.0	0.80	.742	.291
5	6.0x	1.20	.804	.277
6	8.0x	1.40	.804	.278
7	10.0x	2.00	.810	.275
8	9.0x	1.80	.810	
9	12.0x	2.40	.815	

H₂DZ = approximately 4.20×10^{-7} moles

pH 9.8

Chloroform phase 15 ml ; aqueous phase 10 ml

1	1×10^{-7}	0.20	.432	.372
2	2x	0.40	.519	.344
3	3x	0.60	.620	.317
4	4x	0.80	.740	.297
5	5x	1.00	.775	.298
6	6x	1.20	.788	.292
7	8x	1.60	.796	.291
8	10x	2.00	.800	.290

GRAPH No 6

ABSORBANCE VERSUS THALLOUS ION CONCENTRATION
DITHIZONE CONCENTRATION
CONSTANT

- a) pH 9.4 (A) at 510 m μ
 b) pH 9.4 (A) at 440 " "
 c) pH 9.6 (A) at 510 " "
 d) pH 9.6 (A) at 440 " "




TABLE IX (Graph 7)

Effect of pH on absorbance

Dithizone and thalious ion concentrations constant.

H ₂ DZ moles	H ₂ DZ (ml)	Tl ⁺ moles	Tl ⁺ (5x10 ⁻⁴) ml	Buffer ml	pH	(A) at* 510 mμ
6.5x10 ⁻⁷	11.0	6.5x10 ⁻⁷	1.30	12.7	6.0	0.545
6.5x	11.0	13.0x	2.60	11.4	6.0	0.555
6.5x10 ⁻⁷	11.0	6.5x10 ⁻⁷	1.30	12.7	8.0	0.575
6.5x	11.0	13.0x	2.60	11.4	8.0	0.700
6.5x10 ⁻⁷	11.0	6.5x10 ⁻⁷	1.30	12.7	9.0	1.100
6.5x	11.0	13.0x	2.60	11.4	9.0	1.240
6.5x10 ⁻⁷	11.0	6.5x10 ⁻⁷	1.30	12.7	9.6	1.430
6.5x	11.0	13.0x	2.60	11.4	9.6	1.550
6.5x10 ⁻⁷	11.0	6.5x10 ⁻⁷	1.30	12.7	10.0	1.640
6.5x	11.0	13.0x	2.60	11.4	10.0	1.750

* (A) --- absorbance

TABLE X (Graph 8)


Weight of (Tl ₂ Dz)	Chloroform (ml)	Absorbance at 510 mμ	Molar (Tl ₂ H ₂ Dz)	a _m *
1.40 mg	50.0	1.730	3.04x10 ⁻⁵	5.69x10 ⁺⁵
1.12 mg	50.0	1.413	2.43x	6.14x
0.84 mg	50.0	1.060	1.82x	5.82x
0.56 mg	50.0	0.703	1.21x	5.80x
0.39 mg	50.0	0.500	0.85x	5.82x
0.28 mg	50.0	0.330	0.61x	5.41x
				Av. 5.78x10 ⁻⁵

a_m is computed assuming that thalious dithizonate is of keto form having molecular weight of 459.7.

GRAPH No 7


ABSORBANCE VERSUS PH

- a) Thallous ion excess
b) Equivalent amount of thallous ion



GRAPH No 8

ABSORBANCE VERSUS THALLOUS DITHIZONATE
mg/50 ml IN
CHLOROFORM AT 510 $m\mu$



Solutions for polarographic work were:

0.1 M in potassium chloride (KCl)

0.002% w/v in gelatin

Purified nitrogen was bubbled through the solution for 15 min.

Two compartment cell with agar-gel salt bridge and S.C.E.

reference cell on one side--was used.

TABLE XI (Graph 9, 10, 11)
Deflections


Volts (-)	(A) $S=2$	(B) $S=2$	(C) $S=2$	(D) $S=2$	(E) $S=2$	(F) $S=2$	(G) $S=2$
0.00	-3.5	-2.5	-2.0	-2.2	-2.5	0.0	0.0
0.05	0.9	2.0	1.0	2.0	1.0	4.0	4.0
0.10	7.1	8.5	6.8	12.0	7.2	9.9	9.9
0.15	9.1	11.1	9.5	16.0	10.5	13.0	13.0
0.20	10.6	13.0	11.0	18.1	12.1	16.0	15.9
0.25	12.0	14.5	12.0		13.5	17.4	17.5
0.30	13.2	15.5	12.6	21.5	14.7	19.0	18/8
0.35	16.0	17.0	14.0	22.7	15.5	21.0	20.0
0.375	19.2	19.8	16.0	23.6			
0.40	25.7	23.5	20.0	25.1	16.0	23.0	21.3
0.425	45.0	53.2	26.9		25.0	23.2	
0.450	75.0	48.6	38.4	35.7	16.8	28.0	25.0
0.475	$S=5$ 50.5	73.0	56.7	45.6		32.0	29.0
0.500	65.0	92.0	73.0	54.1	17.8	36.5	33.0
0.525	74.5	$S=5$ 42.5	82.7	59.5			
0.55	78.6	44.8	87.5	62.0	18.5	41.0	37.5
0.60	81.2	46.4	90.3	64.0	19.0	43.5	40.0
0.65	82.2	47.0	91.2	65.3	19.8	44.2	41.5

0.70	89.8	47.5	92.3	67.0	20.9	45.6	43.0
0.75	83.0	48.0	93.3	68.8	22.3	47.0	44.5
0.80	83.2	48.5	94.4	70.5	23.8	49.0	45.8
0.85	83.8	49.6	95.4	72.6	25.3	51.0	47.8
0.90	84.5	50.0	96.4	75.0	26.8	53.0	49.5
0.95	85.1	50.5	97.4	78.5	28.9		
1.00	85.5	51.0	98.2	80.5	30.2		
1.05					32.0		
1.10					33.3		
1.15					34.5		

- a) Tl ion concentration $9.80 \times 10^{-4} \text{ M}$
- b) " " " $5.00 \times 10^{-4} \text{ M}$
- c) " " " $4.00 \times 10^{-4} \text{ M}$
- d) " " " $2 \times 10^{-4} \text{ M}$
- e) 0.1M KCl--supporting electrolyte.
- f) 2.24 mg of unknown thallos dithizonate
- g) 1.86 mg of unknown thallos dithizonate.

GRAPH No 9
GALVANOMETER DEFLECTIONS VERSUS
POTENTIAL


a) THALLOUS ION CONCENTRATION 9.8×10^{-4}
 b) " " " $5.0 \times$
 c) " " " $4.0 \times$
 d) " " " $2.0 \times$
 e) 0.1M KCl SUPPORTING ELECTROLYTE




GRAPH No 10

UNKNOWN THALLOUS DITHIZONATE
GALVANOMETER DEFLECTIONS VERSUS
POTENTIAL APPLIED

f) 2.24 mg of Thallous dithizonate
g) 1.86 mg " " "



GRAPH No 11

DIFFUSION CURRENT VERSUS THALLOUS ION
CONCENTRATION

From graph 11 we can evaluate the concentration of thalious ion. Two samples of unknown thalious dithizonate prepared as will be described were decomposed with hot sulfuric acid and hydrogen peroxide. In order to reduce thallium III back to thallium I hydroxyl ammonium chloride was used. The completeness of reduction was checked by measuring the absorbance of aqueous solution at $215\text{ m}\mu$ and $240\text{ m}\mu$ as recommended by Gritsen (6).

By polarographic work the following results were obtained:

Weight of sample mg	Galv. Deflections	Tl M/l observed	Tl M/l computed
2.24	16.5	0.94×10^{-4}	0.96×10^{-4}
1.86	13.7	0.79×10^{-4}	0.80×10^{-4}

From graphs 2A and 2B, curves a and b, it is evident that thalious dithizonate in chloroform solution has a maximum absorption at $510\text{ m}\mu$ regardless of the ratio of reagents mixed. The curve c confirms that thalious dithizonate prepared as will be described later in chloroform solution gives a maximum absorption at $510\text{ m}\mu$. After addition of (BAL) to thalious dithizonate the absorption curve d becomes almost identical to that of the dithizone in chloroform solution (curve f). Thalious dithizonate in chloroform solution upon being exposed to strong sun light fades to a yellowish solution. This absorption spectrum (curve g) resembles that of the oxidation product

of dithizone (diphenylthiocarbondiazone) (14). Besides, it is worth mentioning that the maximum absorption for thallos dithizonate $510 \text{ m}\mu$ almost coincide with a minimum for dithizone which is at $505 \text{ m}\mu$.

In the application of Job's continuous variation method it is seen in graph 3 curves b, c, d, e; that minima or maxima in (Y) occur almost at the same value of (x) . This confirms the point of view that only one complex is formed where the "n" value is unity. The slightly higher values of (Y) and (A) when $x = 0.5$ are probably due to the greater solubility of keto-thallos dithizonate (TiH_4Dz). But from previous observations on the effect of relative ratios of reagents it is impossible to conclude that the enol complex might be formed.

The maximum in (A) occurs when $\frac{dA}{dx} = 0$ which is seen from the graph to fall at the point where $x = 0.48$ and

$$n = \frac{x}{1-x} = 1.$$

From graph 4 it is evident that the difference between absorbances observed without (BAL) and after the addition of (BAL) gradually increases up to $x = 0.5$ and then reaches a minimum at approximately $x = 0.6$ and again returns to the expected level. This behavior, probably is due to the formation of enol thallos dithizonate, but there is no further substantiating evidence to prove it.

In graph 5 curves a, b, give maximum at $x = 0.5$, which confirms that there is only one complex present in chloroform solution at pH 9.4 and pH 9.6. Also the plot ^{of Y} against (x) shows a maximum at $x = 0.5$.

In order to establish the effect of one reagent on the formation of the colored product the measurements of absorbance may be used as recommended by Mellon (10). Keeping dithizone concentration constant and varying the amount of thalious ion the curves a, b, c, and d, in graph 6 are obtained. It is evident that in the beginning absorbance increases considerably until it reaches the point where the reagents are in 1:1 ratios. On further addition of thalious ion the absorbance increases very slightly which is probably due to the greater solubility of thalious dithizonate in chloroform. If there were another complex formed the slope would change at another equivalence point where the reagents would be in ratio (1:2) but this is not noticed.

The pH of the solution is of great importance in the study of metal dithizonates with respect to both: their formation and their solubility in the organic phase. By varying the pH and keeping dithizone and thalious ion concentrations and volumes constant curves a and b in graph 7 were obtained. Between pH 6--8 there is only a small amount of thalious dithizonate in chloroform phase, above pH 8 it increases rapidly. Besides, the excess thalious ion gives higher absorbance probably due to the increased solubility of thalious dithizonate in chloroform.

For determination of molar absorbandy index solid thalious dithizonate was prepared as follows:

Dithizone chloroform solution was shaken with thalious carbonate buffered at pH 9.6. After a few minutes the chloroform solution was separated from the aqueous phase and chloroform was evaporated slowly in the dark at room temperature, until a precipitate formed and only a few ml. of chloroform were left. Then it was filtered and the precipitate washed with cold alcohol and chloroform mixtures. The product was dried in a desiccator and was used for determinations, of molar absorbandy index as shown in graph 8, and polarographic studies.

On account of time limitations work could not be carried to the extent planned. One of the interestes of this work was to establish the equilibrium constant

$$K_m = \frac{[M^+]_w [HD_2]_w}{(M \ell HD_2)_w}$$

for thalious dithizonates between two phases: chloroform and water. As already mentioned in previous discussion K_m could be computed from the relationship

$$\frac{(M \ell HD_2)_o}{[M^+]_w} = \frac{P_m}{K_m} \frac{[H_2 D_2]_o}{F [H^+]_w}$$

if the partition coefficient (P_m), the amount of metal dithizonate in chloroform, the concentration of metal ion in water and the undissociated dithizone concentration in chloroform were known.

Having prepared kete-thalious dithizonate (TLEDZ) it is possible to determine the partition coefficient by absorbance measurements of chloroform solution or by using a radioactive isotope of thallium and determining by tracer methods.

Thalious ion concentration (Tl^+) in the aqueous phase might be determined by measuring the absorbance at 215 *m μ* if there are interfering substances present.

The amount of keto-thalious dithizonate ($TlHDZ$)_o may be determined by measuring the absorbance at 510 *m μ* . Since the molar absorbancy index for dithizone is about 100 times less than that for keto thalious dithizonate, the correction of observed absorbance is unnecessary, unless dithizone is in excess.

The amount of free dithizone (H_2DZ)_o in the organic phase might be determined by extraction of thalious dithizonate with dilute acid, and measuring the absorbance of chloroform solution at 610 *m μ* .

The (H^+) concentration of the solution may be measured by a pH meter. The "F" value is computed by Irving and co-workers as already mentioned.

From this data K_m may be evaluated.

- (1) K. Buch and F. Koroleff, *Finska Kemistsamfundets Medd.*
54, 98, (1945)
- (2) S. S. Cooper; M. L. Sullivan, *Analyt. Chem.*, 23, 613, (1951)
- (3) E. Fischer, *Ann.*, 190, 118 (1878)
- (4) H. Fischer, *Angew. Chem.*, 47, 685, (1934)
- (5) H. Fischer, *Angew. Chem.*, 50, 919, (1937)
- (6) J. Gritmon, Unpublished work, Union College, (1952)
- (7) H. Irving; S. J. H. Cooke; S. G. Woodger; R. J. P. Williams;
J. Chem. Soc., 1347, 1949
- (8) P. Job, *Compt. rend.*, 282, 1621-3
- (9) P. Job, *Ann. chim.*, 2, 113-203, (1928)
- (10) M. G. Mellon, *Analytical Absorption Spectroscopy*, p. 48-9, (1951)
- (12) E. B. Sandell, *Colorimetric Determination of Traces of Metals*,
p. 89, V. 3, 2nd Ed., (1950)
- (13) E. B. Sandell, *Colorimetric Determination of Traces of Metals*,
p. 90, V. 3, 2nd Ed., (1950)
- (14) E. B. Sandell, *Colorimetric Determination of Traces of Metals*,
p. 102, V. 3, 2nd Ed., (1950)
- (15) E. B. Sandell, *Colorimetric Determination of Traces of Metals*,
p. 106, V. 3, 2nd Ed., (1950)
- (16) E. B. Sandell, *Colorimetric Determination of Traces of Metals*,
p. 107, V. 3, 2nd Ed., (1950)
- (17) W. W. Scott, *Standard Methods of Chemical Analysis*, p. 944.
- (18) G. Schwarzenbach and A. Willi, *Helv. Chim. Acta*, 34, 532, (1951)
- (19) A. D. Snyder A Thesis presented to Department of Chemistry,
Union College, (1950)
- (20) W. D. Treadwell, *Helv. Chim. Acta*, 31, 577, (1948)
- (21) W. C. Vosburgh; G. R. Cooper, *J. Am. Chem. Soc.*, 63, 437, (1941)
- (22) Willard; Merrit; Dean; *Instrumental Methods of Analysis*, p. 183
- (23) H. H. Willard; P. Young, *J. Am. Chem. Soc.*, 50, 1322 (1928)
- (24) H. H. Willard; P. Young, *J. Am. Chem. Soc.*, 52, 36 (1930)