



## Original Article

## Predictors of emotional exhaustion among physicians from Iraq - a descriptive cross-sectional multicentre study

Saad Ahmed Ali Jadoo<sup>1\*</sup>, Ilker Dastan<sup>2</sup>, Mustafa Ali Mustafa Al Samarrai<sup>3</sup>, Shukur Mahmood Yaseen<sup>4</sup>, Perihan Torun<sup>1</sup>

### Abstract

**Background:** Doctors and paramedics in countries suffering from long acting conflicts including Iraq are working in serious and exceptional conditions, putting them under severe physical and psychological pressure, therefore examining burnout is important when dealing with quality of care and working conditions. The aim of this study was to assess the point prevalence and to explore factors associated with emotional exhaustion (EE) among medical doctors in Iraq.

**Methods:** Descriptive and cross sectional study was conducted (January to June 2014) among a randomly selected sample of medical doctors (n=576, 87.3% response rate) working in twenty large general hospitals and medical centers. In additions to EE, the self-administered questionnaire used consisting of questions on sociodemographic, work-related characteristics, conflict-related variables and job satisfaction. EE was measured using the emotional exhaustion subscale of the Maslach Burnout Inventory (MBI).

**Results:** The prevalence of EE was reported by 60.0% of the respondents. In multiple linear regression analysis the emotional burnout was higher among doctors who were married, female, bearing children, being threatened, displaced internally, non-specialist doctors, working more than 40 hours per week, experienced unsafe medical practice, disagreed with the way manger handle the staff and those who reported that the doctor-patient relationship as not excellent.

**Conclusion:** Our findings suggest that job dissatisfaction, conflict and violence related factors were significantly associated with high level of emotional exhaustion among Iraqi physician.

**Keywords:** Emotional exhaustion, burnout, workplace, doctors, job satisfaction, conflict, Iraq

### Background

The burnout syndrome has been widely discussed since its first time emerged in its three dimensions; emotional exhaustion, depersonalization, and low personal accomplishment [1]. Work-related burnout is becoming increasingly recognized as a serious problem affecting many people working in human services, especially healthcare workers [2]. It has been shown to be directly related to a considerable list of adverse outcomes, including absenteeism from work, increased turnover and poor job performance [3]. Burnout has also been shown to have effects on physicians and their mental and physical health [4,5],

in addition to the clinical performance and the quality of care [6,7]. These features have been regularly reviewed and recognized over the past fifty years [8]. Literature suggest that burnout is a common health problem among doctors in different countries globally [9-12]. However, despite such wide recognition in Western societies, the subject of burnout has not received the required attention from the Middle East researchers until the beginning of 21st century, and to our knowledge, there is a lack of national studies on burnout among Iraqi health workers. This is a particular problem in Iraq where the health care system has suffered a long standing catastrophic collapse since 1991 and has been overloaded since that time [13]. Furthermore, after the US-led invasion of Iraq in 2003, health system exposed to deliberated destruction of its infrastructure accompanied with new exodus of brain drain including the medical doctors [14]. In fact, the high level of violence in

\*Correspondence: [drsaadalezzi@gmail.com](mailto:drsaadalezzi@gmail.com)

<sup>1</sup>Department of Public Health, Faculty of Medicine, Bezmialem Vakif University, Istanbul, Turkey

Full list of author information is available at the end of the article



conflict zones such as Iraq produced chronic stress among health personnel, so that continuity in work became almost impossible [15]. Moreover, lack of services and the attempt of government to spoil the doctor-patient relationship by directing the media to focus mainly on the performance of health care providers made them victims of unfortunate and unjustified incidents of different type of abuse [15,16]. Thus, the combination of lack of security, work and violence related and political factors has significantly contributed to raise the level of turnover intention and migration among Iraqi doctors. Consequently, the staying doctors, who are still working in Iraq, were not able to cope with the resultant weakness in the level of health service presented to the patients who already feel dissatisfied and carry negative impression against this service [17-19]. As a results, the doctors are more likely to pass into a state of an inevitable physical and emotional exhaustion (EE) ending to an exaggerated motivation to leave job and looking for a better opportunity outside their country [14-19].

This study aimed to predict the main factors related to emotional burnout among Iraqi doctors working in different health care centers during the eruption of armed conflict in 2014. The study focused on Job satisfaction, violence-related variables, work-related variables, and socio-demographic factors to investigate their interrelationships with EE.

## Methods

### Study design and subjects

This cross-sectional study was conducted among Iraqi medical doctors to test the impact of violence-related and workplace-related variables on the Overall EE. The present study is part of a larger research initiative [14,15], in which we recruited a multistage sampling technique to draw a random sample of 660 physicians from twenty large general hospitals and medical centers covering the main five geographically regions in Iraq (north, west, south, central, and the capital city). Design, sampling, and data collection have been reported in detail previously [14,15]. In current analysis, responses were received from 576 physicians (response rate 87.3%). At the time of survey, all Iraqi physicians working at the selected hospitals were included and received a copy of the self-administered questionnaire manually with contact number and email of the data collector. The exclusion criteria included the chief medical officers (CMO), hospital managers and their deputies.

### Outcome variable

Emotional burnout was assessed by nine items of EE subscale of the validated Maslach Burnout Inventory (MBI) [20-22]. EE were answered in terms of frequency on a 7-point Likert scale ranging from 0 (never) to 6 (every day). The overall EE was measured by summing the scores of the 9 items (in range of 0 to 54). A higher score indicated greater emotional exhaustion [3] and, accordingly, a higher emotional burnout. The Cronbach's alpha coefficient of the EE subscale reported in this study was 0.89.

### Independent variables

Job satisfaction was measured with the 10-item Warr-Cook-Wall (WCW) job satisfaction (seven-point Likert-type) scale

ranging from 1 = "very dissatisfied" to 7 = "very satisfied". The overall job satisfaction was measured by summing the scores of the 10 items (in range of 10 to 70).

The socio-demographic variables were collapsed and coded as follows: age (more or less than 40 years old); gender (male or female), marital status (married or single), presence of children (yes or no), residency (Urban or rural). Variables of conflicts or war-related violence were collapsed and coded as either (1) "Yes" or (0) "No" in response to the following questions: "Because of war violence in Iraq, have you been lost a family member?"; "Have you been threatened?"; "Have you been displaced internally?"; "Do you think that medical practice is safe?" and "The doctor-patient-relationship is excellent?".

Individual work-related variables were categorized as follows: the current professional level (specialist or not); the working hours per week (more or less than 40 hours / week); the number of years spent at their work or the same facility (more or less than 10 years); the type of employment (government only or dual in government and private); the training and educational opportunities (yes or no); The way the senior manager handles the staff is effective (agreed or disagreed).

### Statistical analysis

The Statistical Package for Social Sciences (SPSS) version 16.0 was used to analyze data in this study. Descriptive analysis on sociodemographic, conflict-related variables, work-related characteristics and job satisfaction was performed. The 9 items of emotional exhaustion subscale were summed to obtain the total score (0 to 54). A high degree emotional burnout was based on the cut-off point of the emotional exhaustion subscale in the MBI ( $\geq 27$ ) [3]. Data were presented as mean and standard deviation (SD). Test of normality of the total score of emotional exhaustion was conducted.

Student's *t*-test was used to compare the mean of emotional burnout score across demographic variables, work-related characteristics, conflict-related variables and job satisfaction. Categorical data were presented as numbers and percentages and the chi-squared test was used for statistical analysis. Multivariate linear regression using "Backward" technique was employed to obtain factors associated significantly with emotional burnout score. Variables that were significantly associated with burnout in the bivariate analysis were included in the multivariate analysis. The accepted level of significance was set below 0.05 ( $P < 0.05$ ).

## Results

### Descriptive analyses

Table 1 shows descriptive characteristics of the socio-demographic variables. The mean age ( $\pm$ SD) was 40.43 years ( $\pm 8.59$ ) (in range of 27-56). More than half of respondent (53.8%) were females, married (64.2%), living in urban region (63.4%) and bearing children (51.2%). There was significant difference on overall emotional exhaustion. Overall emotional exhaustion was higher among female doctors (95% CI = 0.79-3.16,  $p=0.001$ ), married (95% CI = 0.99-3.46,  $p<0.000$ ), and those who are bearing children (95% CI = 0.59-2.96,  $p=0.003$ ).

**Table 1** Socio-demographic variables on overall emotional exhaustion (n=576)

Variable	Category	N%	Mean $\pm$ SD	t-test	P value	95% CI Upper-Lower
Age	> or=40 years old	291(50.5)	28.73+7.46	1.355	0.176	0.37-2.02
	<40 years old	285(49.5)	27.90+7.12			
Gender	Female	310 (53.8)	29.23+7.81	3.262	0.001	0.79-3.16
	Male	266 (46.2)	27.26+6.51			
Marital status	Married	370 (64.2)	29.11+7.32	3.542	0.000	0.99-3.46
	Single	206(35.8)	26.89+7.05			
Presence of children	Yes	295(51.2)	29.18+6.96	2.935	0.003	0.59-2.96
	No	281(48.8)	27.41+7.55			
Residency	Urban	365 (63.4)	28.49+7.34	0.747	0.456	0.77-1.71
	Rural	211 (36.6)	28.02+7.24			

**Emotional exhaustion (Burnout)**

The mean ( $\pm$ SD) value on the total emotional exhaustion score was 28.72 (SD = 7.30). Three hundred and forty five respondents (60.0%) experienced high level of emotional burnout (Table 2).

**Table 2** Descriptive statistics of the 9 items and overall emotional exhaustion scale

No.	Emotional Exhaustion (Burnout)	Mean	S.D.	Min.	Max.
1	I feel emotionally drained from my work	3.18	1.63	0	6
2	I feel used up at the end of the workday	3.26	1.69	0	6
3	I feel fatigued when I get up in the morning and have to face another day on the job	3.10	1.71	0	6
4	Working with people all day is really a strain for me	3.12	1.66	0	6
5	I feel burned out from my work	3.07	1.70	0	6
6	I feel frustrated by my job	3.25	1.69	0	6
7	I feel I'm working too hard on my job	3.00	1.72	0	6
8	Working with people directly puts too much stress on me	3.43	1.72	0	6
9	I feel like I'm at the end of my rope	2.90	1.68	0	6
10	Overall scale emotional exhaustion	28.72	7.30	0	54

**Conflict-related variables on overall emotional exhaustion**

In table 3, about one-fourth (26.6%) had lost one or more of their close relatives, 54.3% have been threatened, 39.1% have been internally displaced at least once, 66.8% experienced unsafe medical practice, however 70.7% of them described their relation with patient as excellent. Most of the conflict or war-

related variables showed significant difference on overall emotional exhaustion. Overall emotional exhaustion was higher among doctors who have been threatened or kidnapped (95% CI = 0.75-3.13,  $p=0.000$ ), internally displaced (95% CI = 1.50-3.91,  $p<0.000$ ), experienced unsafe medical practice (95% CI = 0.106-3.57,  $p<0.000$ ), and those who described the doctors-patients relationship was unwell (95% CI = 1.40-3.99,  $p<0.000$ ).

**Work - related variables on overall emotional exhaustion**

In table 4, the vast majority (73.3%) of respondents spent more than 10 years in the same health facility, and were not being specialist yet (60.2%), however more than half of them (55.7%) have dual job (government and private) and were satisfied with the available training and educational opportunities. Overall emotional exhaustion was higher among doctors who were not specialist (95% CI = 0.72-3.14,  $p=0.002$ ), disagreed with the way manger handles the staff (95% CI = 0.72-3.14,  $p=0.003$ ) and those who are working more than 40 hours per week (95% CI = 0.59-2.97,  $p=0.003$ ).

**Predictors of burnout**

Table 5 shows the results of multiple linear regression analysis to identify the associated variables with emotional burnout. In backward elimination (or backward deletion) the multivariate linear regression analysis (after excluding of non-contributing variables ) was statistically significant, and overall, explained 24.8% of the variance in the overall emotional exhaustion,  $F(11, 16.845) = 680.670$ ,  $P < .0005$ . The "Internally displaced" and the "doctors-patient relationship" appeared to be the strongest factors predicting the EE (Table 5). Doctors who were internally displaced were more likely to have high EE ( $B = 0.269$ ,  $P < .0005$ ). Doctors who considered the doctor-Patient relationship is not excellent were more likely to have high EE ( $B = 0.267$ ,  $P < .0005$ ). In general, the emotional burnout was higher among doctors who were married, female, bearing children, being threatened, displaced internally, non-specialist doctors, working more than 40 hours per week, experienced unsafe medical practice, disagreed with the way manger handle the staff and considered the doctor-patient relationship as not excellent.

**Table 3** Conflict and war-related variables on overall emotional exhaustion (n=576)

Variable	Category	N%	Mean+ SD	t-test	P value	95%CI Upper-Lower
Loss of family member	No	423(73.4)	28.39+7.31	0.395	0.693	1.08-1.63
	Yes	153 (26.6)	28.12+7.29			
Exposure to threat or kidnapped	Yes	313(54.3)	29.20+7.33	3.207	0.001	0.75-3.13
	No	263(45.7)	27.26+7.13			
Internally displaced	Yes	225(39.1)	29.96+7.73	4.404	0.000	1.50-3.91
	No	351(60.9)	27.26+6.82			
Medical practice in Iraq is safe	No	385 (66.8)	29.09+7.48	3.623	0.000	1.06-3.57
	Yes	191(33.2)	26.77+7.48			
Doctor-patient relationship is excellent	no	169(29.3)	30.22+7.02	4.096	0.000	1.40-3.99
	Yes	407(70.7)	27.53+7.27			

**Table 4** Work - related variables on overall emotional exhaustion (n=576)

Variable	Category	N%	Mean+ SD	t-test	P value	95%CI Upper-Lower
Current professional level	Non-specialist	347(60.2)	29.08+6.97	3.124	0.002	0.72-3.14
	Specialist	229(39.8)	27.16+7.63			
Way managers handle staff	Agreed (yes)	343(59.5)	27.56+7.37	3.034	0.003	0.66-3.07
	Disagreed (no)	233(40.5)	29.43+7.07			
Training and educational opportunities	Yes	316(54.9)	28.84+7.33	1.881	0.060	0.05-2.34
	No	260(45.1)	27.79+7.23			
Years of service	>10 years	422(73.3)	28.55+7.37	1.283	0.216	0.50-2.20
	< or = 10 years	154(26.7)	27.69+7.07			
Hours of work/week	>40 h	267(46.4)	29.27+7.21	2.941	0.003	0.59-2.97
	<or=40 h	309(53.6)	27.49+7.17			
Type of employment	Government and private	321(55.7)	28.58+7.43	0.977	0.329	0.60-1.80
	Government only	255(44.3)	27.98+7.13			

**Table 5** Factors associated with emotional exhaustion in multiple linear regressions (n=576)

Variables	B	S.E.	Beta	t-test	Sig.	95% CI Lower-Upper	Tolerance	VIF
Constant	39.048	1.302	-	29.994	0.000	36.49-41.61	-	-
Overall job satisfaction	0.048	0.023	0.098	2.115	0.035	0.03-0.93	0.626	1.598
Working more than 40 h/week	1.426	0.570	0.098	2.501	0.013	0.31-2.55	0.872	1.147
40 h/week or less						Reference		
Female	1.261	0.552	0.078	2.283	0.023	0.18-2.35	0.928	1.078
Male						Reference		
Non -Specialist	1.602	0.657	0.108	2.438	0.015	0.31-2.89	0.677	1.477
Specialist						Reference		
Not agree with way manger handle staff	1.482	0.549	0.100	2.700	0.007	0.41-2.56	0.970	1.031
Agree						Reference		
Doctor -patient relationship is not excellent	4.244	0.688	<b>0.267</b>	6.165	0.000	2.89-5.60	0.715	1.398
Excellent						Reference		
Displaced internally	4.398	0.624	<b>0.296</b>	7.052	0.000	3.17-5.62	0.759	1.317
Not						Reference		
Presence of Children	1.929	0.545	0.133	3.538	0.000	0.86-3.00	0.948	1.055
Not						Reference		
Married	2.924	0.682	0.193	4.287	0.000	1.58-4.26	0.661	1.514
Single, divorced, widow						Reference		
Have been threatened	1.928	0.582	0.132	3.314	0.001	0.79-3.07	0.838	1.193
Not						Reference		
Medical practice is Not safe	4.036	0.715	0.261	5.642	0.000	2.63-5.44	0.625	1.601
Medical practice is Safe						Reference		

## Discussion

It was an interesting exercise to test several factors that may predict the EE among Iraqi doctors. However, it was not possible to capture all possible factors. Only 24.8% of the variation was explained; 75.2% of the variation remained unexplained. Clearly, other factors may affect the EE. In this study, the mean value of the total burnout score among Iraqi physicians was found to be 28.72 (SD=7.3, range 0-54). Around 60% of the study participants revealed experiencing high level of emotional burnout. This finding is consistent with other international studies. Elsewhere in the Middle East, burnout has been found to be quite common among health professionals in Tunisia [23]. In Saudi Arabia, a high prevalence of burnout among multinational health professionals has been reported [24]. A quarter of the health professionals had suffered from high levels of burnout in Egypt [25]. In contrast, a study conducted in Qatar found that only 12.6% of responding primary care physicians experienced burnout [26].

The first step in coping with burnout threats is to identify the most common causes. The results from previous studies show that among the doctors, burnout is the result of exposure to factors associated with both personal and work environments. Taking into consideration the conclusions from previous studies and the different health system and culture in Iraq, in the current study, the following factors were investigated for their association with burnout, with the aim of reducing burnout among Iraqi doctors: socio-demographic variables (age, gender, marital status, presence of children, residency), work-related variables (job satisfaction, being specialist, years of service, less than 40 hours of work per week, working in both government and private sector, managers handle staff well, poor doctor-patient relationship), and workplace violence variables (loss of family member, exposure to threat or kidnapping, internally displaced, unsafe medical practice). Eleven of these factors maintained in the backward regression analysis. Thus, burnout is potentially multi-determined, therefore, any type of prevention or intervention will require multi-faceted approaches.

In our study, multivariate regression analysis of variables affecting burnout revealed that among the demographic factors, being female, having children or being married are significant predictors of burnout among the study population in Iraq. Regarding to the role of gender in burnout, some papers found female doctors to be at a higher risk of burnout [27,28]. This is in agreement with the findings of the present study. However, only few studies reported burnout by sex, and a study conducted across Europe found burnout to be more prevalent amongst males [29]. In our study, married physicians and physicians having children are expected to have higher level of burnout than their counterparts. In a study in Turkey, it was reported that problems with childcare were significantly associated with burnout [30]. Disequilibrium between family and work demands was found to be one of the most important factors contributing to burnout among Iraqi physicians [31]. In contrast, unmarried physicians were found to have a higher level of burnout and more desensitization in a few other studies [32,33]. Physicians are exposed to a variety of work-related factors that are important in creating burnout.

This study showed that the prevalence of burnout is higher in physicians reporting lower job satisfaction. The entire literature evaluating the correlation between job satisfaction and burnout has yielded the same result [34-38]. However, the majority of the literature is based on cross-sectional surveys as it does in this study, which makes it difficult to verify a causality. While the researchers suggest that low job satisfaction was a risk factor for burnout, it can be asserted that burnout is a risk factor for low job satisfaction, or poor job satisfaction may itself be a symptom of burnout. Further research is needed to clarify the negative relationship.

Among the work-related factors, work time and the doctor-patient relationship are two other factors that have been shown to be related to burnout, in both this and previous studies [39-42]. Physicians in the current study who worked more than 40 hours per week or reported dissatisfaction with doctor-patient relationship had significantly higher burnout rates. Iraqi healthcare system has faced a catastrophic collapse since 1991. Iraqi doctors have left the country due to falling wages and worsening security conditions during the sanction years and after the 2003 invasion [43]. Even though the migration rates have slowed more recently, it was estimated that about half of the doctors have already left Iraq after the 2003 invasion, and most healthcare providers have significant turnover intentions or to leave the country [44]. In addition, health care system has been rapidly changing in recent years, which have increased many doctors' responsibilities, as well as demands of their patients. As a result, doctors are working longer hours and taking less time to communicate with patients, and medical disputes often occur caused by patient dissatisfaction. These combined effects have created an increasing conflict between patients and doctors and damaged the doctor-patient relationship [14]. Consequently, organizational interventions such as reducing long working hours and developing doctor-patient relationships should be considered in new health care reforms. The doctor-patient relationship needs be improved through the implementation of various measures, including malpractice insurance for physicians and improving doctor-patient communication [45].

There is broad acceptance that work satisfaction and burnout levels are strongly affected by the leadership skills of senior managers. Previous studies have shown that relationship with managers is an important factor that affect burnout [46,47]. Poor relationships with managers can lead to feelings of exclusion and loneliness among physicians that can easily turn into burnout [20]. Our study reported similar results; physicians who reported low managerial skills of their immediate supervisors have higher rates of burnout. The findings in this study contribute to the understanding that high levels of managerial support and skills have positive effects on fighting with burnout. The reduction of occupational stress, development of managerial skills of supervisors and strengthening relationship with managers can decrease burnout and improve quality of life of physicians, as seen in other studies [48] [50].

Further, non-specialist physicians are found to be at high risk of burnout in our study group. This finding is similar to the results of other studies. For example, in a Finnish study, the

highest burnout scores were noted in general practitioners and non-specialists working in health centers. This was due to heavy patient loads, long hours of work, low salaries, and professional identity issues [49]. Conflict-related factors and workplace violence have not been systematically examined as a predictor of physician's burnout except a few studies. The present study results demonstrated that physicians who were threatened, who were displaced internally, or who believed medical practice was not safe were more likely to have higher burnout rates in our studied sample. Previous studies in Egypt, Poland, Turkey and a meta-sample of seven studies from different regions also demonstrated that workplace violence, conflicts, personal threats, or incivility were positively associated with the psychological conditions of physicians, including burnout [25,50,51].

There are some limitations to this study. First of all, because the study design was cross-sectional, a causal relationship could not be established. Second, because we had no information about the non-respondents, response bias is a possible bias. Third, this study was conducted during a peak crises of conflict in 2014 which affected the accessibility to the hospitals and districts and consequently a selection bias is another possible bias. Fourth, "Although we have test piloted the English version of the questionnaire, the language barrier could be a limitation because the native language of the respondent is Arabic".

### Policy implication

The above findings may can be applied to the design and implementation of burnout reduction programs in Iraq and facilitate the development of rational strategies. For example, the results showed burnout in physicians to be higher in females, married, or individuals with children. Therefore, we suggest that the target group of burnout reduction programs should primarily focus on these physicians. Intervention strategies may cover communication skill training, stress and time management. Further, burnout was found to be associated with work-related factors: job satisfaction, working hours, being non-specialist, poor relation with managers and patients. Improving working conditions of Iraqi doctors should be an important goal for their job satisfaction and low burnout levels. Policy reforms can address these factors contributing to burnout by reducing workload, reasonable welfare, a healthier working environment and senior-junior support groups. Increasing compensation can be an effective method to help attracting and retaining good doctors. A large proportion of patients in hospitals are affecting the quality of care and the doctor-patient relationship and therefore the burnout of doctors. A comprehensive health care reform is needed to strengthen primary health care intended to address the overutilization of hospitals. This may involve the government hiring and training more staff. Conflict and violence related factors were found to be significantly associated with burnout of Iraqi physicians in current study. Measures to prevent violence in workplace against doctors are needed to reduce risk and negative consequences. Although health institutions in general have made considerable progress in the development of programs to prevent violence in Iraq, there are still shortcomings in the

legislation prohibiting violence at work, and it is essential that laws are enacted to improve the medical environment and prevent violence at work. In order to effectively carry out these policies, government and health care organizations must develop, implement, and encourage a strong safety and security environment. [51].

### Conclusion

High level of emotional exhaustion has been detected in about two third (60.0%) of participants. Findings of this study show that various socio-demographic, occupational and conflict-related factors are important determinants of the burnout of Iraqi doctors. The results of this study would therefore be of great interest to Iraqi policy makers and health care managers seeking to reduce physician burnout levels. The importance of burnout in physicians should not be underestimated and many of the associated factors identified may potentially be reversed. This study may serve as a guide for setting effective interventions and treatment. According to the results, the focus should not only be on providing financial incentives but also on special efforts to create a safe and acceptable work environment.

### Abbreviations

EE: Emotional Exhaustion CMO: Chief Medical Officers WCW: Warr-Cook-Wall SD: Standard Deviation CI: Confidence Interval OR: Odds ratio WHO: World Health Organization GDP: Gross Domestic Product

### Declarations

#### Acknowledgement

We render our special thanks to all directors of hospitals and medical centers we visited. We are also grateful to all the Iraqi doctors and the paramedical staff for their working every day to serve their public in the face of violence and for their time and openness during the data collection.

#### Funding

The author (s) received no financial support for the research, authorship, and/or publication of this article.

#### Availability of data and materials

Data will be available by emailing drsaadalezzi@gmail.com

#### Authors' contributions

SAAJ is the principal investigator of the study who designed the study and coordinated all aspects of the research including all steps of the manuscript preparation. He is responsible for the study concept, design, writing, reviewing, editing and approving the manuscript in its final form. ID, MA, SY and PT contributed in the study design, analysis and interpretation of data, drafting the work, writing the manuscript and reviewed and approved the manuscript. All authors read and approved the final manuscript.

#### Ethics approval and consent to participate

We conducted the research following the Declaration of Helsinki, and the protocol was approved by the Ethical Committee of the Izmir University of Economics (Ref: B.30.2.IEU.0.05.05-020-014). Confidentiality was assured with signed informed consent.

#### Consent for publication

Not applicable

#### Competing interest

The authors declare that they have no competing interests.

#### Open Access

This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits

unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated

#### Author details

<sup>1</sup>Department of Public Health, Faculty of Medicine, Bezmialem Vakif University, Istanbul, Turkey. <sup>2</sup>Department of Economics, Izmir University of Economics, Izmir, Turkey. <sup>3</sup>Department of Public Health, Faculty of Medicine, Anbar University, Anbar, Iraq. <sup>4</sup>Department of Medical Biology and Anatomy, Faculty of Medicine, Diyala University, Diyala, Iraq.

#### Article Info

Received: 23 May 2018

Accepted: 8 November 2018

Published: 8 December 2018

#### References

- Maslach C, Jackson SE. The measurement of experienced burnout. *Journal of Occupational Behaviour* 1981;2:99-113. .
- Sørgaard KW, Ryan P, Hill R, Dawson I; OSCAR Group. Sources of stress and burnout in acute psychiatric care: inpatient vs. community staff. *Soc Psychiatry Psychiatr Epidemiol* 2007;42(10):794–802.
- Maslach CC, Schaufeli WB, Leiter MP. Job burnout. *Annu Rev Psychol* 2001;52:397–422.
- Deary IJ, Blenkin H, Agius RM, Endler NS, Zealley H, Wood R. Models of job-related stress and personal achievement among consultant doctors. *British Journal of Psychology* 1996; 87(1):3–29.
- Deckard G, Meterko M, Field D. Physician burnout: An examination of personal, professional, and organizational relationships. *Medical Care* 1994; 32: 745– 754.
- McCue JD. The effects of stress on physicians and their medical practice. *N Engl J Med* 1982;306:458–463.
- Shanafelt TD, Bradley KA, Wipf JE, Back AL. Burnout and self-reported patient care in an internal medicine residency program. *Ann Intern Med* 2002;136:358–367.
- Felton JS. Burnout as a clinical entity- its importance in health care workers. *J Occup Health* 1998; 48: 237–50.
- Balch CM, Freischlag JA, Shanafelt TD. Stress and burnout among surgeons: Understanding and managing the syndrome and avoiding the adverse consequences. *Archives of Surgery* 2009; 144:371–376.
- Houkes I, Winants Y, Twellaar M. Specific determinants of burnout among male and female general practitioners: A cross-lagged panel analysis. *Journal of Occupational and Organizational Psychology* 2008; 81: 249–276.
- Schaufeli W, Enzmann D. *The burnout companion to study and practice: A critical analysis*. London: Taylor & Francis, 1998.
- Wu S, Zhu W, Li H, Wang Z, Wang M. Relationship between job burnout and occupational stress among doctors in China. *Stress and Health* 2008; 24: 143–149.
- Squires A, Sindi A, Fennie K. Health system reconstruction: perspectives of Iraqi physicians. *Global Public Health* 2010; 5:561–77.
- Ali Jadoo SA, Aljunid SM, Dastan I, Tawfeeq RS, Mustafa MA, Ganasegeran K, Aldubai SA. Job satisfaction and turnover intention among Iraqi doctors-a descriptive cross-sectional multicenter study. *Hum Resour Health* 2015; 13:21.
- Ali Jadoo SA, Torun P, Dastan I, Al-Samarrai M. Impact of conflict related and workplace related violence on job satisfaction among physicians from Iraq –a descriptive cross-sectional multi centre study. *Journal of Ideas in Health* 2018; 1(1): 14-22.
- Al-Samarrai M, Ali Jadoo SA. Iraqi medical students, are still planning to leave after graduation. *Journal of Ideas in Health* 2018;1(1):23-28.
- Sa'adoon A A, Hussien AH, Museher TR. Patients' satisfaction for health care services at Thi-qar province, Iraq. *Thi-Qar Medical Journal (TQMJ)* 2008; 2(1):39–45.
- Shabila NP, Al-Tawil NG, Al-Hadithi TS, Sondorp E: A qualitative assessment of the Iraqi primary health care system. *World Health Popul* 2012; 13(3): 18-27.
- Webster P. Reconstruction efforts in Iraq failing health care. *Lancet* 2009; 373: 617-620.
- Ozyurt A, Hayran O, Sur H. Predictors of burnout and job satisfaction among Turkish physicians. *QJM* 2006; 99 (3):161–169.
- Beckstead JW. Confirmatory factor analysis of the Maslach Burnout Inventory among Florida nurses. *International Journal of Nursing Studies* 2002; 39 (8): 785–792.
- Al-Dubai SAR, Ganasegeran K, Perianayagam W, Rampal KG. Emotional burnout, perceived sources of job stress, professional fulfilment, and engagement among medical residents in Malaysia. *Scientific World Journal* 2013; 2013: 137620.
- Halayem-Dhouib S, Zaghoudi L, Zremdini R, Maalej I, Béchir MB, Labbène R. Burnout en psychiatrie: une expérience tunisienne [Burnout among mental health professionals: a Tunisian experience]. *Rev Epidemiol Sante Publique* 2010;58(6):403–408.
- Al-Turki HA, Al-Turki RA, Al-Dardas HA, Al-Gazal MR, Al-Maghribi GH, Al-Enizi NH, et al. Burnout syndrome among multinational nurses working in Saudi Arabia. *Ann Afr Med* 2010;9(4):226–229.
- Abdo SAM, El-Sallamy RM, El-Sherbiny AAM, Kabbash IA. Burnout among physicians and nursing staff working in the emergency hospital of Tanta University, Egypt. *East Mediterr Health J*. 2016 Mar 15;21(12):906-915.
- Abdulla L, Al-Qahtani DM, Al-Kuwari MG. Prevalence and determinants of burnout syndrome among primary healthcare physicians in Qatar. *South African Family Practice Journal*. 2011;53(4):380–383.
- Dyrbye LN, Shanafelt TD, Balch CM, Satele D, Sloan J, Freischlag J. Relationship between work-home conflicts and burnout among American surgeons: a comparison by sex. *Arch Surg* 2011;146:211–217.
- Shanafelt TD, Oreskovich MR, Dyrbye LN, Satele DV, Hanks JB, Sloan JA, et al. Avoiding burnout: the personal health habits and wellness practices of US surgeons. *Ann Surg* 2012;255(4):625–633.
- Soler JK, Yaman H, Esteva M, Dobbs F, Asenova RS, Katic M, et al. Burnout in European family doctors: the EGPRN study. *Fam Pract* 2008;25:245–265.
- Demir A, Ulusoy M, Ulusoy MF. Investigation of factors influencing burnout levels in the professional and private lives of nurses. *Int J Nurs Stud* 2003;40(8):807–27.
- Lagerström M, Josephson M, Arsalani N, Fallahi-Khoshknab M. Striving for balance between family and work demands among Iranian nurses. *Nurs Sci Q* 2010;23(2):166–172.
- Prins JT, van der Heijden FM, Hoekstra-Weebers JE, Bakker AB, van de Wiel HB, Jacobs B, et al. Burnout, engagement and resident physicians' self-reported errors. *Psychol Health Med* 2009; 14: 654–66.
- Wang, Zhihui, Zhenyu Xie, Junming Dai, Liqian Zhang, Yunbiao Huang, and Bo Chen. "Physician burnout and its associated factors: a cross-sectional study in Shanghai." *Journal of occupational health* 56, no. 1 (2014): 73-83.
- McManus IC, Winder BC, Gordon D. The causal links between stress and burnout in a longitudinal study of UK doctors. *Lancet* 2002; 359: 2089–90.
- Visser MRM, Smets EMA, Oort FJ, de Haes HCJM. Stress, satisfaction and burnout among Dutch medical specialists. *CMAJ* 2003; 168(3): 271–75.
- Elit L, Trim K, Mand-Bains IH, Sussman J, Grunfeld E, and in conjunction with the Society of Gynecologic Oncology Canada. Job

- satisfaction, stress and burnout among Canadian gynecologic oncologists. *Gynecol Oncol* 2004; 94(1):134–139.
37. Thomsen S, Soares J, Nolan P, Dallender J, Arnetz B. Feelings of professional fulfilment and exhaustion in mental health personnel: the importance of organizational and individual factors. *Psychother Psychosom* 1999; 68:157–64.
38. Renzi C, Tabolli S, Ianni A, Di Pietro C, Puddu P. Burnout and job satisfaction comparing healthcare staff of a dermatological hospital and a general hospital. *J Eur Acad Dermatol Venereol* 2005; 19:153–157.
39. Maccacaro G, Di Tommaso F, Ferrai P, Bonatti D, Bombana S, Merseburger A. The effort of being male: a survey on gender and burnout. *Med Lav* 2011; 102:286–296.
40. Piko BF. Burnout, role conflict, job satisfaction and psychosocial health among Hungarian health care staff: a questionnaire survey. *Int J Nurs Stud* 2006; 43:311–318.
41. Wu H, Liu L, Wang Y, Gao F, Zhao X, Wang L. Factors associated with burnout among Chinese hospital doctors: a cross-sectional study. *BMC public health* 2013; 13(1): 786.
42. Al-Dubai SA, Rampal KG. Prevalence and associated factors of burnout among doctors in Yemen. *J Occup Health* 2010;52:58–65.
43. Al-Dewachi O. *The Professionalization of the Iraqi Medical Doctors In Britain*. Ann Arbor, United States: Harvard University; 2008.
44. Webster PC. Iraq's health system yet to heal from ravages of war. *Lancet*. 2011;378(9794):863–6.
45. Gong Y, Han T, Chen W, Dib HH, Yang G, Zhuang R, Lu Z. Prevalence of anxiety and depressive symptoms and related risk factors among physicians in China: a cross-sectional study. *PloS one* 2014; 9(7): e103242.
46. Escribà-Agüir V, Martín-Baena D, Pérez-Hoyos S: Psychosocial work environment and burnout among emergency medical and nursing staff. *Int Arch Occup Environ Health* 2006; 80:127–133.
47. Karasek R, Theorell T. *Healthy work: stress, productivity, and the reconstruction of working life*. New York: Basic Books; 1990.
48. Wang H, Zhang N: Analysis on the status of occupational burnout among doctors. *Zhong Guo Jian Kang Xin Li Xue Za Zhi* 2008; 16:397–399.
49. Olkinuora M, Asp S, Juntunen J, Kauttu K, Strid L, Aärimaa M, et al. Stress symptoms, burnout and suicidal thoughts in Finnish physicians. *Soc Psychiatr Epidemiol* 1990; 25(2):81–86.
50. Merez D, Drabek M, Moscicka A. Aggression at the workplace—psychological consequences of abusive encounter with coworkers and clients. *International Journal of Occupational Medicine and Environmental Health* 2009; 22: 243–260.
51. Aytac S, Dursun S. The effect on employees of violence climate in the workplace. *Work* 41 Suppl 1:2012: 3026–3031.