

2017


Exploring the association between FGM/C and fistula: A review of the evidence

Pooja Sripad
Population Council

Charity Ndwiga
Population Council

Kaji Tamanna Keya
Population Council

Follow this and additional works at: https://knowledgecommons.popcouncil.org/departments_sbsr-rh

 Part of the [Demography, Population, and Ecology Commons](#), [Family, Life Course, and Society Commons](#), [International Public Health Commons](#), [Maternal and Child Health Commons](#), [Sociology of Culture Commons](#), and the [Women's Health Commons](#)

Recommended Citation

Sripad, Pooja, Charity Ndwiga, and Kaji Tamanna Keya. 2017. "Exploring the association between FGM/C and fistula: A review of the evidence," Evidence to End FGM/C: Research to Help Girls and Women Thrive. New York: Population Council.

This Report is brought to you for free and open access by the Population Council.



Research to Help Girls and Women Thrive

EXPLORING THE ASSOCIATION BETWEEN FGM/C AND FISTULA: A REVIEW OF THE EVIDENCE

May 2017



EXPLORING THE ASSOCIATION BETWEEN FGM/C AND FISTULA: A REVIEW OF THE EVIDENCE

POOJA SRIPAD
POPULATION COUNCIL

CHARITY NDWIGA
POPULATION COUNCIL

KAJI TAMANNA KEYA
POPULATION COUNCIL

MAY 2017

The Evidence to End FGM/C programme consortium generates evidence to inform and influence investments, policies, and programs for ending female genital mutilation/cutting in different contexts. Evidence to End FGM/C is led by the Population Council in partnership with the Africa Coordination Centre for Abandonment of Female Genital Mutilation/Cutting, Kenya (ACCAF); Gender and Reproductive Health & Rights Resource Centre, Sudan (GRACE); Global Research and Advocacy Group, Senegal (GRAG); MannionDaniels, Ltd. (MD); Population Reference Bureau (PRB); University of California, San Diego (Prof. Bettina Shell-Duncan); and University of Washington (Dr. Gerry Mackie). Evidence to End FGM/C is funded by UK aid by the UK Government.



The Population Council confronts critical health and development issues—from stopping the spread of HIV to improving reproductive health and ensuring that young people lead full and productive lives. Through biomedical, social science, and public health research in 50 countries, we work with our partners to deliver solutions that lead to more effective policies, programs, and technologies that improve lives around the world. Established in 1952 and headquartered in New York, the Council is a nongovernmental, non-profit organization governed by an international board of trustees. www.popcouncil.org

Suggested citation: Pooja Sripad, Charity Ndwiga, Tamanna Keya. May 2017. “Exploring the association between FGM/C and fistula: A review of the evidence.” Evidence to End FGM/C: Research to Help Women Thrive. New York: Population Council.

Please address any inquiries about the Evidence to End FGM/C programme consortium to:

Dr. Jacinta Muteshi, Project Director, jmuteshi@popcouncil.org

© 2017 The Population Council, Inc.

Funded by:



This document is an output from a project funded by the UK aid by the UK Government for the benefit of developing countries. However, the views expressed and information contained in it are not necessarily those of, or endorsed by, the UK Government, which can accept no responsibility for such views or information or for any reliance placed on them.

Acknowledgements

This report was co-authored by Pooja Sripad (Associate), Charity Ndwiga (Program Officer), and Kaji Tamanna Keya (Intern) of the Population Council. The report is nested within the *Evidence to End FGM.C: Research to Help Girls and Women Thrive* programme, directed by Jacinta Muteshi of the Population Council. The authors of this report are thankful for the thoughtful internal reviews provided by Jacinta Muteshi, Carolyne Njue, Charlotte Warren, and Benjamin Bellows of the Population Council.

The authors extend their gratitude to Vandana Tripathi, Director of Fistula Care*Plus* at EngenderHealth and Susan Igras, Senior Technical Advisor at Institute for Reproductive Health, Georgetown University, who provided critical external reviews. The input provided by all these individuals collectively strengthened the quality and coverage of the report, ensuring that the findings presented reflect the state of the evidence on the association between FGM/C and fistula. The authors also appreciate the editorial support of Janet Munyasya and Robert Pursley, Population Council.

Table of Contents

Acknowledgements	ii
Acronyms	iv
Executive Summary	v
Introduction	1
Methods.....	2
Findings.....	5
Article description.....	5
Association between FGM/C and Fistula	5
Formal assessment of association of FGM and fistula.....	7
Speculative association of FGM and fistula.....	7
Health and Social Consequences of FGM/C and Fistula	12
Discussion.....	16
Summary and conceptual mapping.....	16
Limitations.....	17
Implications for research, programming, and policy	17
References	19
Appendix	23

Acronyms

ANC	Antenatal care
DfID	Department for International Development (UK)
EmONC	Emergency obstetric and newborn care
FGM/C	Female genital mutilation/cutting
HIC	High income country
LMIC	Low and middle income country
OF	Obstetric Fistula
OR	Odds Ratio
PC	Population Council
PPH	Postpartum hemorrhage
PNC	Postnatal care
RR	Relative risk
RVF	Recto-vaginal fistula
SSA	Sub-Saharan Africa
UNFPA	United Nations Population Fund
VVF	Vesico-vaginal fistula
WHO	World Health Organisation

Executive Summary

Female genital mutilation/cutting (FGM/C) and fistula are both conditions that have a range of health, socioeconomic and lifestyle causes and consequences for women globally. There has been sparse empirical and conceptual effort to look at how these two conditions manifest and relate to one another. This rigorous review of the literature aims to fill this research gap by assessing the state of evidence on the association of FGM/C and fistula and conceptually mapping this association within broader social, political, and health systems contexts.

We applied a systematic approach to reviewing English-language peer-reviewed and grey literature and assessed the quality of the evidence using DfID's *Assessing the Strength of Evidence* guide (DfID 2014) to capture domains such as transparency, appropriateness, reliability, validity, and cogency. We applied a non-time bound search strategy through multiple databases using a combination of terms that captured the range of names for FGM/C and fistula. In addition, we drew on key references and global reports. A four-person reviewer team identified the final list of articles from which information was extracted using a data form with open and closed-ended questions related to FGM/C, fistula, other health and social consequences, context, population sample, study methodology, and quality indicators.

The review identified 30 full text documents of mixed quality to include in our final analysis. Eleven studies formally assessed the association of FGM/C with the occurrence of fistula, out of which eight described the conditions as positively related and three described no association. Risks of fistula development increase with cut severity. Nineteen studies speculate that these two conditions are positively and indirectly related with varying degrees of evidence. Findings also covered a range of health consequences of FGM/C and fistula, including maternal and newborn, gynecological, and psychosocial consequences; quality of care and health systems capacity moderating influences; as well as underlying factors that affect both conditions. In some studies, the proposed indirect relationship was physiologically reasoned—namely, that FGM/C-induced scarring may lead to the development of fistula by the tightening of the vaginal wall, prolonged labour, or deinfibulation following delivery or perineal tears. Others describe key determinants of both conditions that lead to a speculative positive association.

Our review also found that maternal and newborn consequences during pregnancy, labour, delivery, and postpartum are the most commonly cited as factors affecting the FGM/C and fistula relationship. Beyond obstructed or prolonged labour and fetal distress, particularly in women who have undergone FGM/C Type III which predisposes women to developing fistula, maternal complications of FGM/C include postpartum hemorrhage, perineal tears and trauma and episiotomies; adverse newborn outcomes include infant resuscitation, neonatal death, and low birth weight. Gynecological consequences of FGM/C—a subset of which are also consequences of fistula—include urogenital outcomes (e.g. scarring, keloids, abscesses, fistula, damaged tissue, and cysts), infertility, and reduced sexual functioning and satisfaction. Moreover, women who undergo FGM/C and those who develop fistula independently suffer from psychosocial consequences such as trauma, fear, depression, and divorce—all of which affect their association.

Our review additionally found that quality of care and the capacity of the health system to prevent, detect, and treat consequences of FGM/C affects the likelihood of a woman developing a fistula. Access to instrumental delivery (cesarean sections, episiotomies, and deinfibulation)—particularly in severe forms of FGM, is essential to enable women to deliver safely. Studies suggest that the efficacy of longer hospital stays for delivery among women with FGM/C is debatable in preventing fistula, but the need for sufficient human and material resource capacity for emergency obstetric care and newborn resuscitation is clear. Provider knowledge of how to counsel and treat women who have undergone FGM/C—in both native and migrant populations—is variable, but overall lacking across high, middle and low income contexts. In some cases, this is due to inadequate legal and professional guidance.

Finally, all the studies describe underlying socio- economic and contextual factors, including gender and socio-cultural norms, that affect both fistula and FGM/C. Poverty, an inability to work, living in a rural area, limited access to health services, and malnutrition (stunting) increase the likelihood of women undergoing FGM/C and developing fistula. The level of education influences a family's choice to continue or abandon the FGM/C practice for a daughter and also has implications on their awareness around the importance of antenatal care and facility delivery to prevent fistula. The underlying intersectionality of gender with socio-cultural norms around controlling one's sexuality and marriageability (FGM/C) as well as early marriage and pregnancy (increases women's risk for fistula development) all affect the social praise and sanctioning of women (cut and uncut/living without and with fistula).

We have developed a conceptual mapping framework to situate the evidence of an association between FGM/C and fistula, including intermediate and underlying factors that relate to both FGM/C and fistula high burden settings. High burden settings include high-income countries (HIC) and low and middle income countries (LMICs) where migrants or underserved populations lack access to proper pregnancy and delivery care. They also include contexts in which women have a higher exposure to sexual violence. We recommend that researchers, as well as policy and program implementers, think through and utilise the framework developed in their work to consider the intersectional influences on both conditions. Qualitatively, it is critical to explore multiple perspectives around FGM/C and fistula to better develop contextually sound interventions. Quantitatively, we recommend the use of latent variables, scales and indices to investigate the influence of positive and negative social norms, health systems, and poverty on both FGM/C and fistula in high FGM/C exposure areas. Implementation research around psychosocial factors is critical to both conditions as is girl-centered holistic programming. Finally, more research is needed on the influence of laws, policies and professional guidelines for health workers on FGM/C (e.g. related to re-infibulation and deinfibulation) and how these affect the experiences of women in HIC and LMIC contexts.

Introduction

Female genital mutilation/cutting (FGM/C) is the partial or total removal of exterior female genitalia or any sort of injury in female genitalia without medical reasons (WHO 2006). WHO categorizes FGM into four different types: Type I, or clitoridectomy, refers to the removal of total or part of the clitoris and prepuce; Type II, or excision, refers to the removal of total or part of the clitoris and labia minora; Type III, also known as infibulation, is the narrowing of the external genitalia and stitching together the edges of the vulva; Type IV, known in some contexts as *Gishiri*, is any kind of non-therapeutic procedure including pricking, piercing, incising, and scraping (WHO 2006). An estimated 200 million girls and women are currently living with FGM/C with the majority residing in 30 countries in low and middle income countries (LMICs) in Africa, Asia and the Middle East (UNICEF 2016). FGM/C in high income countries (HIC) with migrant populations, such as Australia, the United States, the European Union, Norway and Switzerland, is a recognized concern as well (Reisel and Creighton 2015).

Genital fistula is a condition in which a hole between the vagina and the rectum or bladder causes a woman to continuously leak urine, feces or both. Though fistulas can be obstetric, traumatic, or iatrogenic, the most common type is obstetric fistula (OF). Of the approximate one to two million women living with fistula globally, the majority reside in South Asia and sub-Saharan Africa (Adler et al 2013). Obstetric fistula affects about 1.57 per 1,000 women in sub-Saharan Africa (SSA). Obstetric fistula results from inadequately managed prolonged obstructed labour. Error during surgical procedures such as caesarean sections and hysterectomies can lead to iatrogenic fistulas (Barageine et al 2104, Sori et al 2016, Raassen et al 2014). Injury, including through sexual violence, can lead to traumatic fistula, although this is less common than obstetric and iatrogenic fistula in most settings (Longombe 2008). Obstetric fistula is indicative of a health system that has failed to provide accessible, timely and appropriate intrapartum care (Tunçalp 2015). The condition primarily affects those of lower socioeconomic classes, who are underprivileged, un- or under-employed, and have limited access to safe delivery attended by qualified health personnel (Wall 2012, Tebeu 2012, Suzan 2013). Women living with fistula not only suffer from a physically debilitating morbidity, but also a loss of healthy life years. The disability adjusted life years weights for vesico-vaginal (VVF) and rectovaginal fistula (RVF) are 0.342 and 0.501, respectively compared to the 0.187 associated with blindness (Salomon 2015). Women with fistula often exhibit poor health indicators including other co-morbidities of obstructed labour and may experience negative social consequences including lifelong isolation, stigma, shame, and rejection by their husband, family, and community (WHO 2006, Jones 2007, Roush 2009, Mselle 2012).

FGM/C and fistula represent complex conditions that reflect socio-economic, cultural, legal, and health systems determinants of women's health (Kimani et al 2016). There is a lack of literature on these determinants as well as the factors that are associated with prevention, identification, management, and counseling for either condition. Despite the existence of independent literature on FGM/C (Almroth-Berggren 2001, WHO 2006, WHO 2013) and fistula (Bellows et al 2015, Ahmed 2015), there is limited research on the association of the two conditions. Moreover, there is little concurrence on the intensity of the health risks associated with different types of FGM/C and how cut severity relates to various types of fistula (Slanger et al 2002). It is plausible, given the proximity and relationship of the FGM/C site to genital fistula, implications of pregnancy and childbirth, informal cutting procedures and the context of FGM/C, that FGM/C could lead to the development of fistula (obstetric, traumatic, iatrogenic). To date, to the best of our knowledge, there has been no conceptual work mapping the association between FGM/C and fistula.

This rigorous review seeks to fill this research gap and aims to 1) assess the state of evidence on the association of FGM/C and fistula, 2) conceptually map this association within broader social, political, and health systems contexts, 3) identify evidence gaps and areas for further research, and 4) develop recommendations for policy and programming.


Methods

We conducted a rigorous review of English-language peer-reviewed and grey literature, drawing upon processes identified in DfID's *Assessing the Strength of Evidence* guide (2014). We applied a search strategy using the following terms in combination with each other and as Medical Subject Headings (MeSH) terms: 'female genital mutilation', 'FGM', 'female genital cutting', 'FGC', 'female circumcision', 'infibulations' AND 'fistula', 'obstetric complications', 'obstetric consequences', 'types of tears', 'excision', and 'delivery outcomes'. The databases and search engines utilised included PubMed, Google Scholar, Scopus, JSTOR, and Brandeis Scholar as well as organizational databases including Population Council, UNFPA, and EngenderHealth. Our search did not apply time nor geographic restrictions. In addition to the database searches; we drew upon key references cited in review articles and global reports.

Our search strategy (Figure 1) yielded a total of 512 records. Upon screening of titles and abstract review, 422 records were excluded and 61 articles, reports, and academic theses were identified as eligible for full text review. A four-person team read and reviewed the full text documents identified. Each full text document was reviewed by two independent readers and classified as "to include", "to exclude", "not sure". Any discrepancy between two reviewers was deliberated and jointly reviewed by the research team. Thirty full text documents were selected for inclusion in the body of evidence exploring the association between FGM/C and fistula. Articles were excluded if FGM/C and fistula were not mentioned in the articles' findings.

Data were collected through a Google survey form jointly developed by the research team after critical reflection of an initial set of peer-reviewed articles and grey literature. Descriptive measures extracted include authors' names, publication year, country or countries (region) in which the study was conducted, research type (primary, secondary), sample size and population, study design (observational, systematic review, other review), analytic methods used (qualitative, quantitative, mixed), data sources (community, facility, other), and data collection time-frame. Reviews without a specific indication of sample size (number of studies reviewed) were labeled "NI" (not indicated); similarly, facility and community data sources were classified as not applicable (N/A). Research questions, objectives, and key findings were extracted through open-ended responses.

Figure 1. Search strategy flow diagram



Both formal and speculative associations of FGM/C and fistula were abstracted from each article—formal associations refer to those that were explicitly assessed in the results’ section and speculative relationships that may be implicit or explicitly mentioned in the results and discussion with or without data and reference support. Direct, indirect, and inconclusive associations were also extracted. Formal associations were categorically deemed positive (increase in FGM/C, increase in fistula), indirect/did not describe (which included indirect associations and those that weren’t explicitly described), or none (i.e. ‘no association’). Speculative associations were classified as positive (if increase in FGM/C, increase in fistula is supported by data or reference), indirect/did not describe (indirect association through prolonged labour, but lacked explicit descriptive link), and none (i.e. ‘no association’). Negative associations between FGM/C and fistula were not seen nor extracted. Review 1 and Review 2 findings were jointly analysed and reconciled to obtain findings on the association of FGM and fistula.

Reviewers additionally assessed the health and social causes and consequences, including contextual factors that influence FGM/C and fistula using open- and close-ended questions. Obstetric and newborn complications—including postpartum hemorrhage (PPH), pre-term births, sepsis, prolonged labour, maternal death, newborn death and newborn complications – and mode of delivery (e.g. cesarean section, episiotomy, vaginal birth) were explicitly extracted. Open-ended questions about any other emerging associations with FGM/C and fistula allowed reviewers to extract data on a range of other health and social factors including physical (gynecological, sexual functioning, other) and psychological factors, health systems capacity, underlying poverty, gender dynamics, education, socio-cultural norms and legal contexts.

Finally, the quality of evidence was assessed using 19 questions, informed by DFID’s ‘Assessing strength of the evidence’ report, as listed in Table 1. Questions were evaluated as both binary and 3-point responses resulting in a total potential score of 29 points. Each article’s score was calculated and quality level classified in its relation to the maximum score as ‘high’ (>75% score), ‘moderate’ (50-75% score) or ‘low’ (<50% score).

Data were analysed using mixed methods. Closed-ended questions were tabulated and analysed quantitatively using Microsoft Excel. Textual analysis of open-ended questions around causes and consequences of FGM/C and fistula was conducted by the research team through deliberation and corroborating reviewer responses.

Table 1. Assessing quality of evidence

Quality Area	Specific question	Responses
Conceptual framing	Does the study acknowledge existing research?	{0: no, 1:yes}
	Does the study construct a conceptual framework?	{0:no, 1:yes}
	Does the study pose a research question or outline a hypothesis?	{0:no, 1:yes}
Transparency	Does the study present or link to the raw data it analyses?	{0:no, 1:yes}
	Did the authors describe the geography/context in which the study was conducted?	{0:no, 1:partially, 2: in a detailed way}
	Does the study declare sources of support/funding?	{0:no, 1:yes}
Appropriateness	Does the study identify a research design?	{0:no, 1:yes}
	Does the study identify a research method?	{0:no, 1:yes}
	Does the study demonstrate why the chosen design and method are well suited to the research question?	{0: not at all, 1: describes partially, 2: strongly describes}
Cultural sensitivity	Does the study explicitly consider any context-specific cultural factors that may bias the analysis/findings?	{0:no, 1:yes}
Validity	To what extent does the study demonstrate measurement validity?	{0:low, 1: moderate, 2: high}
	To what extent is the study internally valid?	{0:low, 1: moderate, 2: high}
	To what extent is the study externally valid?	{0: not, 1: somewhat, 2: high}

Reliability	To what extent are the measures used in the study stable?	{0: <i>not very</i> , 1: <i>moderately</i> , 2: <i>very</i> }
	To what extent are the measures used in the study internally reliable?	{0: <i>not vary</i> , 1: <i>moderately</i> , 2: <i>very</i> }
	To what extent are the findings likely to be sensitive/changeable depending on the analytical technique used?	{0: <i>highly</i> , 1: <i>moderately</i> , 2: <i>not very</i> }
Cogency	Does the author 'signpost' the reader throughout?	{0: <i>no</i> , 1: <i>yes</i> }
	To what extent does the author consider the study's limitations and/or alternative interpretations of the analysis?	{0: <i>does not</i> , 1: <i>moderately</i> , 2: <i>strongly</i> }
	Are the conclusions clearly based on the study's results?	{0: <i>no</i> , 1: <i>somewhat</i> , 2: <i>yes</i> }

Findings

Article descriptions

Descriptive characteristics of the 30 final documents (detailed in Appendix 1, Table 2) show that more than half (23 of 30) were published between 2000 and 2016 and comprise both grey (8) and peer-reviewed (22) literature. Documents were classified as primary research studies (n=18, 60%) and reviews (n=12) and applied qualitative (13), quantitative (13), and mixed methods (4). With the exception of one high income country investigation, the majority of studies were in Africa, in low- and medium income countries, including Nigeria (5), Kenya, (3), Ethiopia (3), Egypt (2), and Sudan (1). The 14 multi-country studies covered both LMIC and HIC settings. Six of the primary studies focused on women following their fistula repair surgeries; eight explicitly looked at women attending antenatal (ANC), maternity, family planning (FP), and postnatal (PNC) services; and four explored health service provider and other stakeholder perspectives; the latter included nurses, midwives, surgeons, and obstetricians and gynecologists as well as policymakers, opinion leaders and community members. Although the majority of data sources were collected in facilities through record review, clinical observation, and self-reporting, two qualitative studies explicitly drew on community-based sources, and one quantitative secondary analysis of household surveys. Timeframes for primary studies' data collection ranged from under six months (n=6), six months to one year (n=3), and over a year (n=6); the longest were those focused on fistula patients, typically between three and 11 years. Sample sizes ranged from eight to 35,037; eight studies had fewer than 100 respondents, three studies had between 101 and 200, and nine had over 200. Nine studies, all of which were secondary reviews, did not explicitly describe the number of articles reviewed.

An array of research questions and objectives were asked in the studies, focusing either on FGM/C, fistula, or both. Only two studies presented a conceptual framework (Berg et al 2014, Fantu 2007). With the exception of one study that explicitly set out to study the direct association of FGM/C and fistula (Browning et al 2010), rarely did the studies focus on this association. Rather, the reviewed articles sought to explore broader confounding, modifying, and mediating factors relating to both conditions. Research questions and objectives were under the following aims:

- To study the prevalence and the social and health consequences of fistula (n=5)
- To study the association between FGM/C and obstetric complications (n=7)
- To study the social and health consequence of FGM/C (n=13)
- To understand the type, context, and management of FGM (n=4)
- To study health center readiness to treat FGM/C or fistula (n=2)

Association between FGM/C and Fistula

Of the 30 articles, 11 articles formally assessed the association of FGM/C with the occurrence of fistula, out of which eight described the conditions as positively related and three described no association. Eighteen studies speculated that these two conditions were indirectly positively related with varying degrees of evidence (Table 3).

Table 3. Associations between FGM/C and Fistula

Authors	Year Published	Type of FGM/C*	Type of association between FGM/C and fistula (formally assessed)	Type of association between FGM/C and fistula (speculative/assumed)
Jaldesa G, Askew I, Njue C, Wanjiru M	2005	NI; (e.g. III)	Positive	
Jones H, Diop N, Askew I, Kabore I	1999	I, II, III	Positive	
Mabeya HM	2004	III	Positive	
Saracoglu M, Zengin T, Ozturk H, Genc M	2014	IV (Gishiri)	Positive	
Sharfi AR, Elmeqboul MA, Abdella AA	2013	I, II, III, IV	Positive	
Tahzib F	1983	IV (Gishiri)	Positive	
Tukur J, Jido TA, Uzoho CC	2006	IV	Positive	
WHOa	2000	I, II, III, IV	Positive	
Browning A, Allsworth JE, Wall LL	2010	I, II	None	
Peterman A & Johnson K	2009	I, II, III	None	
Slanger TE, Snow RC, Okonofua FE	2002	I, II, III, IV	None	
Al-Hussaini TK	2003	I, II		Indirect /did not describe
Berg RC, Underland V, Odgaard-Jensen J, Fretheim A, Vist GE	2014	NI, review		Indirect /did not describe
Creighton SM	2015	NI, review		Indirect /did not describe
Fantu T	2007	Type II		Indirect /did not describe
Ibekwe PC, Onoh RC, Onyebuchi AK, Ezeonu PO, Ibekwe RO	2012	NI		Indirect /did not describe
Kasim K, Shaaban S, Sadak AEAE, Hassan H	2012	NI		Indirect /did not describe
Khisa M, Nyamongo IK	2012	NI		Indirect /did not describe
Kimani S, Mutesi J, Njue C	2016	NI, review		Indirect /did not describe
Magoha GAO, Magoha OB	2000	NI, review		Positive
Mumtaz A, Mohammed R	2007	NI, review		Indirect /did not describe
Onuh, SO, Igberase GO, Umeora JOU, Okogbenin SA, Otiodo VO, Gharoro EP	2006	NI		Indirect /did not describe
Reisel D, Creighton SM	2015	NI, review		Positive
Ruiz JI, Martinez AP, Bravo PMM, Roche PF	2012	NI, review		Positive
Rushwan H	2000	NI, review		Positive
Ryan F	2012	NI		Indirect /did not describe
Teufel K, Dorfler DM	2013	NI, review		Positive
Toubia N	1994	NI, review		Positive
UNFPA and Engender Health	2003	NI		Indirect /did not describe
WHOb	2006	I, II, III, IV		Indirect /did not describe

*NI: not indicated explicit

Formal assessment of association of FGM and fistula

Eleven studies, including primary and secondary designs, in LMIC settings with high FGM/C prevalence (56% to 95%), formally assessed the association of fistula and FGM/C, with mixed findings reflecting both direct and indirect relationships (Table 4). Among mostly obstetric fistula patients in Kenya, 80 percent had undergone Type III FGM/C (Mabeya 2004). In northern Nigeria, where Type IV FGM/C (*gishiri*) is common, nine to 18 percent of fistulae were attributed to trauma (i.e. cutting process or sexual violence) in addition to 60 percent attributed to obstetric situations (Tahzib 1985, Tukur et al 2006). Among women in Mali and Burkina Faso clinics who had undergone FGM/C and had developed obstetric complications, two and one percent, respectively, reported developing obstetric fistula (Jones et al 1999). Among Sudanese women attending clinics who had undergone FGM/C, 9.2 percent and 14.6 percent developed vesico-vaginal fistula (VVF) and recto-vaginal fistula (RVF), respectively. Two studies in Ethiopia and Nigeria found no association between FGM/C (Type I and II) and the development of obstetric fistula (RR/OR ~1.0).

The studies demonstrate variability and a distinct pattern of FGM/C and fistula association by FGM/C types. Positive associations are reported in studies investigating Type III (infibulation) and Type IV (*gishiri*) FGM/C only (Jaldesa et al 2005, Mabeya 2004, Saracoglu et al 2014, Tahzib 1985, Tukur et al 2006), or in studies of all FGM/C types, individually or in aggregate (Jones et al 1999, Sharf et al 2003, WHOa 2000). Studies with no evidence of association report on fistula effects in populations with greater exposure to Types I and II FGM/C and lack sufficient data to assess Type III's contribution to obstructed labour or fistula (Browning et al 2010, Peterman and Johnson 2009, Slanger et al 2002). One study (Jones et al 1999) that considered FGM/C Types I, II and III, demonstrated that the risk of developing fistula increases with the severity of FGM/C cut.

Quite a few of the formally assessed studies, including those finding no direct statistical association between FGM/C and fistula, describe indirect associations (e.g. via prolonged labour, scarring, tears) and mention contextual, personal, and health systems factors that may influence both conditions and confound the association (Mabeya 2004, Tukur et al 2006, Slanger et al 2002).

Speculative association

Out of the 30 studies, 19 suggested, either explicitly or through presentation and discussion of intermediate factors, association between FGM/C and fistula (Table 5): These studies present indirect mechanisms through which FGM/C affects fistula through related health consequences, including those that might occur at childbirth, and moderated by levels clinical management. Six multi-country reviews (Table 5) explicitly speculated that fistula is an outcome of FGM/C (mainly Type III); five indicated that fistula as sequelae to long-term obstetric complications (prolonged labour, perineal or paraurethral tears) and poor clinical management (of episiotomy or instrumental delivery), while one suggested that the transformation of an infibulation-induced infection into an abscess propels fistula development (Ruiz et al 2012). Some studies speculated that at the time of, and immediately following, cutting procedures (particularly infibulation), women are at risk for urinary and traumatic fistula, as late complications due to injury while being cut (Magoha and Magoha 2000, Muntaz and Mohammed 2007, Reisel and Creighton 2015, Ruiz et al 2012).

The other 13 studies elaborated long-term health consequences incurred during childbirth (Table 5) but implied only indirect relationships to fistula, reporting that fistula can be caused by vaginal wall tightening resulting in FGM/C-induced scarring, prolonged labour, or deinfibulation during delivery or perineal tears that may result in both RVF or VVF (Tal-Hassaini 2003, Reisel and Creighton 2015, Teufel and Dorfler, Rushwan 2000, Slanger et al 2002). Although these articles do not directly causally link FGM/C and fistula, they mention key intermediate conditions comparable to those in the articles explicitly offering speculative positive associations.

Table 4. Studies in which associations were formally assessed

Author	Methods and sample	Prevalence of FGM/C or fistula	Study setting	Type of FGM studied	Association
POSITIVE ASSOCIATIONS					
Jaldesa et. al., 2005	Mixed methods exploratory: In-depth Interview-FGDs, Facility assessments, Client exit interviews attending ANC, & service provider interviews	FGM/C: 97.5% Fistula: 2-3% (reported by nurse-midwife)	Two districts in Kenya (<i>one urban and other rural</i>) with women of the Somali ethnic community and providers	NI; (e.g. III)	Qualitative data from providers suggest FGM indirectly leads to fistula (VVF and RVF) via difficult delivery due to tightening of the vagina, prolonged labour, and perineal tears.
Jones et.al. 1999	Retrospective facility and patient survey (clinical observation & self-report) 5337 women attending clinic for range of RH and maternal services	FGM/C: 93%-94% 477 (of 5337 women at clinic) women developed gynecological complications due to FGM	Mali (rural and urban) and Burkina Faso (rural)	I, II, III	Eight out of 477 women who had delivery complication had fistula. 1% of women who had undergone FGM and developed obstetric complications in Burkina Faso and Mali reported VVF; in Mali an additional 1% reported RVF.
Mabeya 2004	Retrospective facility-based record review 66 women who received fistula repair surgeries	Fistula: 0.1 %	Two rural districts in Kenya	III	80% of repaired women had undergone severe FGM/infibulation
Sharfi et. al., 2003	Retrospective patient survey 2000 women (including university students and women attending outpatient clinic)	FGM/C: 73.4% Type III: 63%	A Sudanese university tertiary hospital in Khartoum	I, II, III, IV (majority: type III)	9.2% (n=57) of women developed VVF 14.6% (n=90) developed rectal vagina fistula. VVF occurred as a result of prolonged obstructed labour.
Saracoglu M. et al., 2014	Review Not Indicated	NI	Multicounty	IV (Gishiri)	<i>Gishiri</i> cut is common among Hausa's living in Northern Nigeria and Southern Niger. 5.68 to 18% of all the fistulae cases in the region result from <i>Gishiri</i> .
Tahzib F, 1985	Facility record reviews 1443 fistula repair patients, out of which 80 were pediatric cases (<13 years)	NI	University hospital in, Northern Nigeria	IV (Gishiri)	60%, 15%, and 9% of pediatric fistula cases resulted from prolonged labour, <i>Gishiri</i> cut, and trauma (sexual violence), respectively.
Tukur et al, 2006	Facility record reviews 50 consecutive fistula patients	Fistula incidence: 150,000-2000,000 cases/year	Jigawa state, Northern Nigeria	IV	After prolonged obstructed labour, the most common cause of VVF was the <i>Gishiri</i> cut (18%) administered by local barbers.
WHOa, 2000	Review 422 articles	NI	Multicounty	I, II, III, IV	7 studies looked at the association, 4 of which found that Type III and IV lead to the development of fistula through obstructed labour, vulval scarring, anatomical distortion and incorrectly performed episiotomy (Damas 1972, Shandall 1967, De Villeneuve 1937 and Pieters 1972). Indirect positive association through obstructed labour was speculative (Tahzib, 1985; Harrison, 1983).

NO ASSOCIATION					
Browning. A et al 2010	Facility record/ case notes review – comparative study 492 fistula patients who were cut (255) and uncut (237)	FGM/C: 81%	Fistula center, Amhara, Ethiopia	I, II	There is no independent association between Type I and II FGM/C the development of obstetric fistula from obstructed labour (RR = 0.93, 95% CI 0.85, 1.02). There was no association between FGC and urinary incontinence (RR = 0.92, 95% CI 0.82, 1.03).
Peterman A & Johnson K, 2009	DHS secondary analysis of population-level surveys 35037 women of reproductive age (15-49 years) across Malawi (11,698), Rwanda (11,321); Uganda (8524) and Ethiopia (4,066)	NI	Community – households in Malawi, Rwanda, Uganda, and Ethiopia	I, II, III	No statistically significant evidence that FGM/C contributes to the prevalence of incontinence (OR = 0.819, 95% CI -0.299, 0.137). In this study, incontinence was used as a proxy for fistula –both traumatic and obstetric.
Slanger et al., 2002	Facility-based comparative study (self-reported on first delivery) 1107 women attending FP or antenatal services (with at least one child)	FGM/C: 56% (n=621 in study); 25% (country)	Edo, Nigeria	I, II, III, IV	No difference in complications and procedures was found between those who had undergone types I and II FGM/C and those who were uncut. Both obstructed labour and perineal tears (major contributing factors to fistula) – odds were not significantly associated with FGM.

Table 5. Speculative associations between FGM/C and fistula

	Authors	Key findings (including proposed relationships)	Data sources
Speculative positive	Magoha and Magoha, 2000	FGM/C may cause fistula in the long term through injury of urological organs such as the uterus and bladder.	Review; cited two articles as evidence (Baker et al. 1993; Dirie and Landmark 1992)
	Reisel and Creighton 2015	FGM/C may lead to damage to the urethra and urethral strictures and lead to fistula formation.	Review; cited Amin et al. 2013 as evidence
	Toubia 1994	Infibulated (Type III FGM/C) women: during labour, the fetal head may get obstructed and lead to VVF.	Review; cited Warsame 1989 as evidence
	Ruiz et.al., 2012	Cut women may have infection that evolves into an abscess and progresses to fistula development.	Review; did not present data or refer to an article
	Teufel and Dorfler, 2013	FGM/C leads to prolonged delivery resulting to VVF in some cases; in other instances, necrosis and perineal tear from FGM/C lead to fistulae. Very tight circumcision scar may lead to what is normally referred to as dribbling incontinence.	Review; did not present data or refer to an article
	Rushwan. H, 2000	Infibulation could increase the risk of fistula through prolonged labour and constant pressure of the fetal head on adjacent organs.	Review; did not present data or refer to an article
Indirect/ did not describe	Al Hussain 2003 Berg et al. 2014 Creighton 2015 Fantu 2007 Ibekwe et al. 2012 Kasim 2012 Khisa 2012 Kimani 2016 Mumtaz & Muhammed 2007 Onuh et al. 2006 Ryan 2012 UNFPA & EngenderHealth 2003 WHO, 2006	Described long term complication of FGM/C such as prolonged labour, perineal tear, episiotomy, postpartum hemorrhage, perineal trauma and caesarean section, difficult delivery, episiotomies.	Data includes primary data, secondary data and review

Health and Social Consequences of FGM/C and Fistula

Nearly all the studies report that FGM/C has consequences for a woman's physical and mental health, from the moment she is cut as a child, throughout her adult life (Berg et al 2014). The most common immediate or short-term consequences described include bleeding, sepsis, urine retention, urethral injury, and urinary fistula (Sharfi et al 2013, Toubia 1994). Consequences due to fistula include serious bad odor due to continuous dripping of urine or feces and stillbirths (UNFPA and EngenderHealth 2003). Long-term health consequences of FGM/C, described in 26 of the 30 articles, include maternal, newborn, and gynecological complications. The psychosocial consequences of FGM/C overlap with those of fistula. In some cases, consequences of FGM/C are the causes of fistula.

Maternal and newborn consequences

Maternal and newborn consequences and outcomes during pregnancy, labour, delivery and postpartum are the most commonly cited intermediate factors in the FGM/C and fistula relationship. Obstructed or prolonged labour is the largest risk factor for obstetric fistula. Sixteen studies describe scenarios where FGM/C scarring, and in cases of Types III and IV, leads to a narrowing of the vaginal orifice and fetal distress. This combined with prolonged labour predisposes women to developing fistula (Mabeya et al 2004, Jaldesa et al 2005, Teufel and Dorfler 2013, Fantu 2007, Tukur et al 2006, Sharfi et al 2013, Toubia 1994, Khisa and Nyamongo 2012, Slanger et al 2002, Kasim et al 2012, Magoha and Magoha 2000). Infibulated women are 2.5 times more likely to develop any kind of obstetric complication than uncut women (Jones et al 1999). Given the nature and severity of Type III FGM/C, these women in particular risk perineal tears and need de-infibulation during ANC or delivery care to safely give birth (Toubia 1994 and Rushwan 2000).

Thirteen articles described PPH as a consequence of FGM/C or hemorrhage at the time of the FGM/C procedure (Sadak et al, Jaldesa et al 2005, Kimani et al 2016, Onuh et al 2016, Kasim et al 2012, Mumtaz and Mohammed 2007, Reisel and Creighton 2015, Ruiz et al 2012, Sharfi et al 2013, Toubia 1994, Teufel and Dorfler 2013, WHO 2006). Seven articles cite perineal tears, trauma, and episiotomy as long-term consequences of FGM/C (Creighton et al 2015, Reisel and Creighton 2015, Kimani et al 2016, Magoha and Magoha 2000, Slanger et al 2002, WHO 2006, WHOa 2000). At least six articles mention maternal death as a consequence of FGM/C (WHOa 2000, Kimani et al 2016, Tukur et al 2006, Ruiz et al 2012, Teufel et al 2013) and 11 report newborn deaths or health complications due to FGM/C (WHOa 2000, Tukur et al 2006, Jones et al 1999, Toubia 1994, Ruiz et al 2012, Teufel and Dorfler 2013, Fantu 2007). Similarly, stillbirths often co-occur with obstetric fistula, given the precursor of prolonged labour (UNFPA and EngenderHealth, 2003). According to one systematic review (WHO 2006), despite unclear explanatory mechanisms, women who have undergone FGM/C are more likely to suffer adverse newborn outcomes including infant resuscitation, neonatal death, and low infant birth weights.

Gynecological consequences

The gynecological consequences of FGM/C revealed in the studies include urogenital outcomes, infertility, and sexual functioning and satisfaction. Urogenital outcomes such as scarring, keloids, abscesses, fistula, damaged tissue (perineum, anal sphincter), disfigurement, vaginal obstruction, and cysts are mentioned in the background sections of studies but are not frequently reported in the research findings (Kimani et al 2016). Five of the 30 articles describe urogenital infection, urine retention, and other moderating factors after FGM/C that could lead to fistula development (Creighton et al 2015, Magoha and Magoha 2000, Sharfi et al 2013, Tukur et al 2006). To address these compounding factors in clinics, some studies stress the need for specialised provider training

and to involve urologists in fistula surgery teams for successful repair (Creighton et al 2015, Magoha and Magoha 2000). Although it may not have implications on obstetric fistula, infertility, reported as a late complication of FGM/C (Tahzib 1983), is more common (2.9 times increased likelihood) among cut women (Kasim et al 2012). Quite a few articles describe women who had undergone FGM/C as more likely to report painful intercourse, no sexual desire, less sexual satisfaction, and less experience of orgasm compared to uncut counterparts (Khisa and Nyamongo 2012, Kimani et al 2016, Saracoglu et al 2014, Sharfi et al 2013).

Psychosocial consequences

Women who undergo FGM/C as well as those who develop fistula independently suffer a number of psychosocial consequences. Understanding how these consequences interrelate may have implications for the associations between the two conditions. At least five articles found that women who had undergone FGM/C suffered from psychosocial consequences such as trauma, fear, depression; in some cases, these led to or were compounded by divorce (Tahzib et al 1983, Creighton et al 2015, Browning 2010, Jaldesa et al 2005, Saracoglu et al 2014, Teufel and Deufel 2013). In cases where women migrate to new contexts where FGM/C is less prevalent, the shock around its non-normative practice also has psychosocial consequences for a woman (Teufel and Deufel 2013). Psychosocial consequences suffered by women living with fistula include isolation, prejudice, depression, and anxiety, many of which are mitigated by divorce or living separately from one's husband or family (Khisa 2015, Tahzib 2015).

Quality of care and health system capacity

Quality of care and the capacity of the health system emerged as a potential mediator in the association between FGM/C and fistula. Qualitative and quantitative findings suggest that improving the capacity of the health system to prevent, detect, and treat consequences of FGM/C affects the progression to fistula (Sharfi 2013, Mabeya 2004, WHO 2006, Onuh et al 2006). Twelve out of the 30 articles describe that FGM/C and delivery type, including access to instrumental delivery (cesarean sections, episiotomies, de-infibulation), is essential to enabling women to deliver safely (Slanger et al 2002, Teufel et al 2013, Onuh 2006, Creighton 2015, Kimani et al 2016). In contexts where a needed episiotomy was not performed, women who have undergone FGM/C have 1.31 to 3.19 times higher relative risk of a perineal tear compared to uncut women (WHO 2006). This risk increases with cut severity: 1.31 greater for those with Type I, 1.92 times greater for those with Type II, and 3.19 for those with Type III. Instrumental delivery is critical, especially for women living with Types III and IV. Some studies observed that women with FGM/C require extended hospital stays (WHO 2006), while others suggest no difference (Creighton 2015, Browning 2010). Many studies advocate that hospitals need human and material resource capacity for newborn resuscitation and emergency obstetric care (EmOC) to avoid fistula, mitigate other maternal consequences (potentially intensified by FGM/C), and avoid newborn death or morbidity (Browning 2010, Kimani et al 2016, WHO 2006). The provision of timely and personalized ANC for women with FGM/C where women are counseled on the importance of skilled birth attendance and deinfibulation in preparation of a vaginal delivery also emerged as critical in HIC and LMIC contexts (Jones et al 1999, Rushwan 2000, UNFPA and Engender Health 2003).

Provider knowledge of how to counsel and treat women who have undergone FGM/C is variable by context. There was less understanding in HICs as FGM/C cases were concentrated among migrant women (Creighton 2015, Rushwan 2000, Reisel and Creighton 2015). In these settings, inadequate or a complete lack of legal and professional guidance constrains providers from managing FGM/C clients appropriately—including policies on re-infibulation and de-infibulation (Reisel and Creighton 2015). De-infibulation by traditional birth attendants (or unskilled attendants)

can cause fistula in some cases (Browning 2010, Toubia 1994, Onuh et al 2006). One study in Kenya showed five in eight women with FGM/C developed fistula during their first delivery due to the unskilled birth attendant with whom they delivered (Khisa and Nyamongo 2013). Capacity building of skilled assistants in many LMICs where women have less access is critical (Jaldesa et al 2005).

Underlying contextual factors influencing both fistula and FGM

Nearly all studies in either the results or interpretation sections, touch on one or more underlying socio-economic and contextual factors that affect both fistula and FGM/C, including poverty, place of residence, and malnutrition. For example, women living in poverty, who are unable or not allowed to work for pay, have limited access health services and have a higher likelihood of continuing FGM/C, are at greater risk of developing fistula. These women have the lesser means to deal with the health and social consequences of either condition (Ryan 2012, Jones et al 1999, Fantu 2007). Women living with fistula are further unable to work given the constant smell they emit, which plays into their cycle of poverty (UNFPA and EngenderHealth 2003). In high prevalence FGM/C areas, malnutrition, which stunts pelvic growth predisposing some women to prolonged labour and elevates their risk of fistula, may further contribute to FGM/C-fistula indirect association (Mabeya 2004). The level of education may influence a family's choice to continue or abandon the FGM/C practice for a daughter. Mothers and fathers with more schooling tend to discontinue the practice over generations (Fantu 2007, Ibekwe et al 2012). Education also affects awareness around the importance of antenatal care and facility delivery, both of which help women with FGM/C to identify prolonged labour and seek skilled care to prevent fistula.

Studies also demonstrate that gender and socio-cultural norms with respect to women's relational roles to spouses and families affect FGM/C and fistula. While early marriage and pregnancy places women at risk for fistula (Mabeya 2004, UNFPA and EngenderHealth 2003), FGM/C is often described as way to control women's sexuality (Ibekwe et al 2012). Moreover, the reduced sexual satisfaction and strained marital relationships due to FGM/C and fistula described by many studies (UNFPA and EngenderHealth 2003, Jaldesa 2005, WHO 2006) have implications for gender-dynamics in households. In both cases, the intersectionality of gender and socio-cultural preferences around women are at stake. Socio-cultural, including religious, norms around marriageability, virginity, sexuality and social status influence FGM/C (Magoha and Magoha 2000, Kimani et al 2016, Ruiz et al 2012, Saracoglu et al 2014). Social sanctioning for uncut women (un-marriageability, promiscuity, dishonor) tends to sustain the FGM/C practice (Ibekwe et al 2012, Browning et al 2010), though these normative claims are rarely assessed in quantitative assessments. Social sanctioning of women living with fistula in many cases relates to their isolation from family and community spaces as well as their inability to remain employed due to their leaking condition (Khisa and Nyamongo 2012, Fantu 2007, Onuh et al 2006). Less information exists on sexual violence and/or rape though it is described as influencing fistula formation (Peterman and Johnson 2009), there is insufficient evidence on how it affects the relationship of FGM/C's with fistula (Tahzib 1985, Fantu 2007).

Quality of evidence

The quality of evidence is mixed on the association of FGM/C and fistula (Table 2). Of the 30 final documents, 12 (40%) are of high quality, 13 (43%) of medium quality, and five (17%) are classified as low quality studies. This mixed level of evidence on a relatively under-studied issue reflects manuscript transparency, cogency, and moderate reliability and transferability of findings, as opposed to an inability to draw significant causal inferences from the limited study designs. Most were facility-based record reviews, of both primary and secondary designs, that although provide


clinical evidence, do not enable us to ascribe a population-based assessment of the association of FGM/C and fistula. Comparative studies that looked at uncut and cut women were helpful, although rarely was fistula a focused outcome. Primary qualitative and quantitative studies were strongest when they insight into the broader context that influenced both FGM/C and fistula.

Discussion

Summary and conceptual mapping

This review sought to describe the existing evidence on the association of FGM/C and fistula and found mixed evidence that reflects both direct and indirect relationships. We found 30 peer-reviewed and grey literature studies that commented on both conditions; while 11 documents formally demonstrated positive and no association between FGM/C and fistula, many others speculate about positive relationships. The dissonance between speculative and evidence-based associations demonstrates a lack of formal rigorous quantitative measures (relative risks and odds ratios) investigating the FGM/C and fistula relationship. Interestingly, the three documents that confirmed no association were carried out in contexts of predominantly FGM/C Types I and II, while those that saw positive associations covered all or focused on FGM/C Types III and IV, indicating a potential dose-response of severity, i.e. the more severe the cut, the more likely a woman is to develop pregnancy and other complications that lead to fistula development. Within these higher severity types, Type III was often associated with obstetric fistula, while Type IV was additionally linked to traumatic fistula caused by cutting procedures themselves. We also found that the indirect association of FGM/C and fistula, including via intermediate health and social consequences of FGM/C, mediating health systems challenges, and underlying socio-economic and context factors, are critical to consider in conceptually framing the two conditions in a way that helps frame implications for research and practice (Figure 2).

Figure 2. Conceptual mapping of associations between FGM/C and fistula



EmONC refers to emergency obstetric and newborn care and ANC refers to antenatal care

This conceptual framework demonstrates the complexity of factors identified through our review that affect both FGM/C and fistula. In addition to the direct and mediated pathway through which FGM/C leads to the development of fistula, there are health consequences of both conditions that may influence the association. Though the data on severity is mixed and insufficiently identified in articles (Table 3), the conceptual framework maps our current understanding. Moreover, the similarities in sociodemographic, economic and cultural context of women exposed to both FGM/C and fistula is reflected in their lack of access to health care services broadly and in the specific context of pregnancy and childbirth; this moderates the association between the two conditions.

Quality of antenatal and delivery care, including the facility material and human resource mix, adequate legal and professional guidance, as well as the competency and sensitivity required of providers tasked with counseling women exposed to both conditions plays a mitigating role. Finally, FGM/C and fistula are situated in settings with shared underlying societal factors that affect our ability to capture a measurable association between the conditions. The intersectional influence of socio-economic (wealth, education) and poverty statuses—including women’s abilities to maintain employment, gender dynamics including women’s autonomy to make health care-seeking decisions, socio-cultural norms and sanctions related to marriageability, cleanliness, and women’s sexuality—determine both FGM/C and fistula. These underlying and health systems causes and consequences of both conditions demand more attention in terms of research, programming, and policy in LMICs and HICs.

Limitations

We faced several challenges in carrying out this review. First, given the complexity of both conditions, there were only a few studies that captured a diverse range of perspectives (qualitative) and reached adequate samples to comment statistically significantly on associations and their strength (quantitative). Secondary analyses were predominantly clinical cases, or other reviews, that did not adequately define replicable methodologies or provide clear evidence of assessing an association. A number of studies took place in facility settings where women received surgical repair or instrumental delivery to safely give birth. By contrast, there were relatively few studies that drew on population-based surveys or targeted community samples; our review had insufficient data from women and men living in communities with high burden of FGM/C and fistula. This selection bias likely lead to an underreporting of evidence on the association of FGM/C and fistula, given that these groups are less likely have access to medical services overall and more likely to experience a co-occurrence of both conditions. Also, a paucity of longitudinal quantitative designs limits our ability to detect causal inference; however, conducting such studies is not pragmatic in most settings given the long data collection time-frame required to collect relevant information, the relative rarity of fistula, and the high rates of loss-to-follow up (WHO 2006).

Implications for research, programming, and policy

Despite these limitations, this rigorous review of the literature points to the lack of an in-depth understanding around the contexts in which associations between various types of FGM/C and different types of fistula exist and require more deliberate programming efforts. The conceptual mapping presented in this report may serve as a particularly useful tool for those involved in programming in that it may help decision makers think through the various consequences, positive and negative, of intervening at various levels. First contexts in which FGM/C is a norm ought to more systematically take the associated risks of fistula and other childbirth outcomes into account. For example, the type of provider counseling intervention may differ between HIC and LMIC contexts based on the types of FGM/C and fistula women typically present with. A social norms targeting program strategy may require a variety of inputs including increasing male and female awareness around the risks of FGM/C on maternal and newborn outcomes and soliciting buy-in of community leaders. The quality of care and underlying factors that affect both FGM/C and fistula point to the importance of advocating for multifaceted girl-centered policy and programming.

There is a need to build the evidence base around understanding and addressing the association of FGM/C and fistula. A second phase of this study is planned and will undertake limited multivariate analyses of the DHS/MICS datasets from multiple African countries that have the relevant data to investigate the associations, causative factors and relevant correlates for FGM/C and fistula. Comparative research across high and low burden FGM/C settings, types of cutting,

socio-economic status, and country policy environments would be useful for shared learning on this ambiguous area in both LMIC and HIC contexts. We recommend that researchers focus on studying confounders, mediators, and underlying factors as related to both FGM/C and fistula at the community level and in high burden settings where both are prevalent. High burden FGM/C settings may be in HICs and LMICs, including among migrants or other underserved populations that lack access to proper pregnancy and delivery care. The association of FGM/C and traumatic fistula may be directly explored in contexts where women are exposed to sexual violence, for example in conflict and post-conflict settings. Traumatic fistula should also be further studied in contexts where unskilled persons are performing FGM/C procedures and unintended injury directly results in fistula formation.

We recommend utilising the conceptual mapping as a guide to look at different complex relationships between factors through both qualitative and quantitative means. Qualitatively, it is critical to explore multiple perspectives around FGM/C and fistula to better develop contextually sound interventions. Quantitatively, we recommend the use of latent variables, scales and indices to investigate the influence of positive and negative social norms, health systems, and poverty on both FGM/C and fistula in high FGM/C exposure areas. Implementation research around psychosocial interventions that addresses depression, anxiety, social and cultural challenges associated with both conditions is a key investigative space to pursue—especially in LMIC contexts where the double burden of FGM/C and fistula is more pronounced. Finally, it is important to conduct further nuanced research on the influence of laws, policies and professional guidelines for health workers around FGM/C (e.g. related to re-infibulation and deinfibulation) and how these affect the experience of women in HIC and LMIC contexts.

References

- Adler AJ, Ronsmans C, Calvert C, Filippi V. 2013. Estimating the prevalence of obstetric fistula: a systematic review and meta-analysis. *BMC Pregnancy and Childbirth* 13(1): 246
- Ahmed S, Tunçalp Ö. 2015. Burden of obstetric fistula: from measurement to action. *Lancet Glob Health* 3(5): e243-4. [http://dx.doi.org/10.1016/S2214-109X\(15\)70105-1](http://dx.doi.org/10.1016/S2214-109X(15)70105-1)
- Al-Hussaini, T.K. 2003. Female genital cutting: Types, motives and perineal damage in labouring Egyptian women. *Medical Principles and Practice* 12(2): 123–128. <http://doi.org/10.1159/000069119>
- Almroth-Berggren LA. 2001. Reinfibulation among Women in a Rural Area in Central Sudan. *Health Care Women Int* 22(8): 711-721
- Barageine JK, Tumwesigye NM, Byamugisha JK, Almroth L, Faxelid E. 2014. Risk Factors for Obstetric Fistula in Western Uganda: A Case Control Study. *PLoS ONE* 9(11): e112299. doi:10.1371/journal.pone.0112299
- Bellows, B., R. Bach, Z. Baker, C. Warren. 2015. Barriers to Obstetric Fistula Treatment in Low-income Countries: A Systematic Review. Nairobi: Population Council.
- Berg, R.C., Underland, V., Odgaard-Jensen, J., Fretheim, A., Vist, G.E. 2014. Effects of female genital cutting on physical health outcomes: A systematic review and meta-analysis. *BMJ Open* 4(11): e006316–e006316. <http://doi.org/10.1136/bmjopen-2014-006316>
- Berg, R.C., Denison, E., Fretheim, A. 2010. Psychological, social and sexual consequences of female genital mutilation/cutting (FGM/C): A systematic review of quantitative studies. Report from Kunnskapssenteret nr 13–2010. Oslo: Nasjonalt kunnskapssenter for helsetjenesten
- Browning, A., Allsworth, J.E., Wall, L.L. 2010. The relationship between female genital cutting and obstetric fistulas. *Obstet Gynecol* 115(3): 578-583. <http://doi.org/10.1038/jid.2014.371>
- Creighton, S.M. 2015. Female genital mutilation (FGM) and the lower urinary tract. *International Journal of Urological Nursing* 9(2): 69-73. <http://doi.org/10.1111/ijun.12075>
- DfID. 2014. How to Note: Assessing strength of evidence. www.gov.uk/government/uploads/system/uploads/attachment_data/file/291982/HTN-strength-evidence-march2014.pdf
- Elmusharaf, S. 2013. *Female genital mutilation: determinants and consequences among girls and women in Sudan*. Karolinska Institutet, Universitetservice AB ISBN 978-91-7549-341-1
- Engender Health. 2012. Living with Obstetric Fistula: Qualitative Research Findings from Bangladesh and the Democratic Republic of Congo. *Fistula Care*, EngenderHealth. www.engenderhealth.org/files/pubs/fistula-care-digital-archive/1/1.2/living-with-obstetric-qualitative-brief-english.pdf
- Fantu, T. 2007. Socio-cultural causes and consequences of fistula on Ethiopian women: case study in Addis Ababa fistula hospital, Master's Thesis
- Ibekwe, P.C., Onoh, R.C., Onyebuchi, A.K., Ezeonu, P.O., Ibekwe, R.O. 2012. Female genital mutilation in Southeast Nigeria: A survey on the current knowledge and practice. *Journal of Public Health and Epidemiology* 4(5): 117-122. <http://doi.org/10.5897/JPHE11.194>
- Jaldesa, G.W., Askew, I., Njue, C., Wanjiru, M. 2005. Female genital cutting among the Somali of Kenya and management of its complications.

www.popcouncil.net/pdfs/frontiers/FR_FinalReports/Kenya_Somali.pdf
http://www.popcouncil.net/pdfs/frontiers/FR%5C_FinalReports/Kenya%5C_Somali.pdf

- Jones, D. 2007. Living testimony: Obstetric fistula and inequities in maternal health. New York: Family Care International. www.msh.org/sites/msh.org/files/living_testimony_2007.pdf
- Jones, H., Diop, N., Askew, I., Kabore, I. 1999. Female genital cutting practices in Burkina Faso and Mali and their negative health outcomes. *Studies in Family Planning*. <http://onlinelibrary.wiley.com/doi/10.1111/j.1728-4465.1999.00219.x/abstract>
- Kasim, K., Shaaban, S., Sadak, A.E.-A. El, Hassan, H. 2012. Impacts of Female Genital Mutilation on Women's Reproductive Health. *Journal of Community Medicine & Health Education* 2(3): 2-5. <http://doi.org/10.4172/jcmhe.1000137>
- Khisa, A.M. and Nyamongo, I.K. 2012. Still living with fistula: An exploratory study of the experience of women with obstetric fistula following corrective surgery in West Pokot, Kenya. *Reproductive Health Matters* 20(40): 59-66. [http://doi.org/10.1016/S0968-8080\(12\)40661-9](http://doi.org/10.1016/S0968-8080(12)40661-9)
- Kimani, S, Mutesi J, Njue C. 2016. Health impacts of FGM/C: A synthesis of the Evidence. Population Council
- Longombe AO, Claude KM, Ruminjo J. 2008. Fistula and traumatic genital injury from sexual violence in a conflict setting in Eastern Congo: case studies. *Reproductive Health Matters* 16(31): 132-141. doi: 10.1016/S0968-8080(08)31350-0
- Mabeya, HM. 2004. Characteristics of women admitted with obstetric fistula in the rural hospitals in West Pokot, Kenya
- Magoha, G.A.O. and Magoha, O.B. 2000. Current global status of female genital mutilation: a review. *East African Medical Journal* 77(5): 268-272
- Mselle LT, Evjen-Olsen B, Moland KM, Mvungi A, Kohi TW. 2012. Hoping for a Normal Life Again: Reintegration after Fistula Repair in Rural Tanzania. *Journal of Obstetrics and Gynaecology* 34(10): 927-938
- Mumtaz, A. and Mohammed, R. 2007. Review Obstetric management of women with female genital mutilation. *The Obstetrician & Gynaecologist* 9: 95-101
- Onuh, S.O., Igberase, G.O., Umeora, J.O.U., Okogbenin, S.A., Otiode, V.O., Gharoro, E.P. 2006. Female genital mutilation: Knowledge, Attitude and Practice among Nurses. *Journal of the National Medical Association* 98(3): 409-414
- Peterman A and Johnson. 2009. Incontinence and trauma: Sexual violence, female genital cutting and proxy measures of gynecological fistula. *Social Science and Medicine* 68(5): 971-979. <http://doi.org/10.1016/j.socscimed.2008.12.006>
- Raassen T, Ngongo CJ, Mahendeka MM. 2014. Latrogenic genitourinary fistula: an 18-year retrospective review of 805 injuries. *Int Urogynecol J* 25(12): 1699-1706
- Reisel, D. and Creighton, S.M. 2015. Long term health consequences of Female Genital Mutilation (FGM). *Maturitas* 80(1): 48-51. <http://doi.org/10.1016/j.maturitas.2014.10.009>
- Rodriguez, M.I., Seuc, A., Say, L., Hindin, M.J. 2016. Episiotomy and obstetric outcomes among women living with type 3 female genital mutilation: a secondary analysis. *Reproductive Health* 13(1): 131. <http://doi.org/10.1186/s12978-016-0242-9>
- Roush KM. 2009. Social implications of obstetric fistula: an integrative review. *J Midwifery Women's Health* 54(2): e21-33. doi: 10.1016/j.jmwh.2008.09.005

- Ruiz Jiménez et al. 2012. Approximation to the Ablation/Female Genital Mutilation (A /FGM) from the Transcultural Nursing. A bibliographical revision. *Enfermeria Global* 28: 411-424
- Rushwan, H. 2000. Female genital mutilation (FGM) management during pregnancy, childbirth and the postpartum period. *International Journal of Gynaecology and Obstetrics* 70: 99-104
- Ryan, F. 2012. The 'typical story' of obstetric fistula: the need to enhance awareness, action and funds, Doctoral Dissertation (working paper)
- Salomon JA, Haagsma JA, Davis A, de Noordhout CM, Polinder S, Havelaar AH, Cassini A, Devleeschauwer B, Kretzschmar M, Speybroeck N, Murray CJ, Vos T. 2015. Disability weights for the Global Burden of Disease 2013 study. *Lancet Glob Health* 3(11): e712-23
- Saracoglu, M. 2014. *Female Genital Mutilation/Cutting Type 4*.
- Sharfi, A.R., Elmeqboul, M.A., Abdella, A.A. 2013. The continuing challenge of female genital mutilation in Sudan. *African Journal of Urology* 19(3): 136–140.
<http://doi.org/10.1016/j.afju.2013.06.002>
- Slanger, T.E., Snow, R.C., Okonofua, F.E. 2002. The impact of female genital cutting on first delivery in southwest Nigeria. *Studies in Family Planning*. <http://doi.org/10.1111/j.1728-4465.2002.00173.x>
- Sori D.A, Azale A.W., Gemed, D.H. 2016. Characteristics and repair outcome of patients with Vesicovaginal fistula managed in Jimma University teaching Hospital, Ethiopia. *BMC Urology* 16: 41 DOI: 10.1186/s12894-016-0152-8©
- Tahzib, F. 1985. Vesico-vaginal fistula in Nigerian Children. *Lancet* : 1291-1293
- Tebeu PM, Fomulu JN, Khaddaj S, de Bernis L, Delvaux T, Rochat CH. 2012. Risk factors for obstetric fistula: a clinical review. *Int Urogynecol J* 23(4): 387-394.
- Teufel, K. and Dörfler, D.M. 2013. Female genital circumcision/mutilation: Implications for female urogynaecological health. *International Urogynecology Journal and Pelvic Floor Dysfunction* 24(12): 2021–2027. <http://doi.org/10.1007/s00192-013-2160-5>
- Toubia, N. 1994. *Female circumcision as a public health issue*. *Nejm*, 712–716.
- Tukur, J., Jido, T.A., Uzoho, C.C. 2006. The contribution of gishiri cut to vesico-vaginal fistula in Birnin Kudu, Northern Nigeria. *African Journal of Urology* 12(3): 121-125
- Tunçalp O, Tripathi V, Landry E, Stanton CK, Ahmed S. 2015. Measuring the incidence and prevalence of obstetric fistula: approaches, needs and recommendations. *Bulletin of the World Health Organisation* 93(1): 60-62
- UNFPA, EngenderHealth. 2003. Obstetric fistula: Needs assessment report: Findings from Nine African Countries. www.endfistula.org/sites/endpointfistula.org/files/pub-pdf/Nine%20Country%20Needs%20Assessment.pdf
- UNICEF. 2016. Female Genital Mutilation/Cutting: A global concern. Fact Sheet. New York. www.unicef.org/media/files/FGMC_2016_brochure_final_UNICEF_SPREAD.pdf
- Varol, N. et al. 2016. Obstetric outcomes for women with female genital mutilation at an Australian hospital, 2006-2012: A descriptive study. *BMC Pregnancy and Childbirth* 16(1): 328. <http://doi.org/10.1186/s12884-016-1123-5>
- Wall LL. 2012. Obstetric Fistula Is a “Neglected Tropical Disease.” *PLoS Negl Trop Dis* 6(8).
- Warren, C. and Mwangi, A. 2008. *Obstetric Fistula: Can Community Midwives make a difference?* Population Council.

- WHO. 2013. Female genital mutilation. www.who.int/mediacentre/factsheets/fs241/en/index.html
- WHO. 2017. 10 Facts on Obstetric Fistula. www.who.int/features/factfiles/obstetric_fistula/en
- WHO, FCH, WMH. 2000. A systematic review of the health complications of female genital mutilation including sequelae in childbirth
- WHO. 2006. Female genital mutilation and obstetric outcome: WHO collaborative prospective study in six African countries. www.global-sisterhood-network.org/content/view/993/76

Appendix 1

Table 2. Descriptive characteristics of final documents

Authors	Year published	Country/ies (region)	Research type	Sample size	Study design	Methods	Data sources	Collection time-frame	Quality
Al-Hussaini TK	2003	Egypt	Primary	306 primigravida maternity admissions	Observational	Quantitative	Facility	21 months	medium
Berg RC, Underland V, Odgaard Jensen J, Fretheim A, Vist GE	2014	Multi-country	Secondary	57 studies	Systematic review	Qualitative	NA	NA	high
Browning A, Allsworth JE, Wall LL	2010	Ethiopia (Amhara)	Primary	492 fistula patients (cut:255, uncut:237)	Observational	Quantitative	Hospital	3 years	high
Creighton SM	2015	UK	Secondary	NI	Other review	Mixed	NA	NA	medium
Fantu T	2007	Ethiopia (Addis Ababa)	Primary	422 fistula patients	Observational	Mixed	Health center	2 months	high
Ibekwe PC, Onoh RC, Onyebuchi AK, Ezeonu PO, Ibekwe RO	2012	Nigeria (Southeast)	Primary	320 antenatal and postnatal women	Observational	Quantitative	Facility	4 months	medium
Jaldesa G, Askew I, Njue C, Wanjiru M	2005	Kenya	Primary	29 influential persons, 14 groups with community members, 15 facility assessments, 101 antenatal clients, 44 service providers	Observational	Qualitative	Community	3-5 months	high
Jones H, Diop N, Askew I, Kabore I	1999	Burkina Faso & Mali	Primary	7257 women attending clinic for antenatal, FP, ObGyn services (Burkina Faso: 1920; Mali: 5337)	Observational	Quantitative	Facility	4 months	high
Kasim K, Shaaban S, Sadak AEAE, Hassan H	2012	Egypt (Alexandria)	Primary	200 circumcised & 200 uncircumcised women attending maternal clinics	Observational	Mixed	Facility	7 months	high

Khisa M, Nyamongo IK	2012	Kenya (West Pokot)	Primary	8 women with fistula, 2 FGDs: fistula survivors and persons attending awareness seminar	Observational	Qualitative	Community	2 months	medium
Kimani S, Mutesi J, Njue C	2016	Multi-country	Secondary	44 reference materials related to FGM/C	Systematic review	Qualitative	NA	NA	high
Mabeya HM	2004	Kenya (West Pokot)	Primary	NI (women treated over 5 year period)	Observational	Quantitative	Health center	5 years	medium
Magoha GAO, Magoha OB	2000	Multi-country (Africa)	Secondary	NI	Other review	Qualitative	NA	NA	medium
Mumtaz A, Mohammed R	2007	Multi-country (UK and HICs)	Secondary	NI	Systematic review	Quantitative	NA	NA	low
Onuh, SO, Igberase GO, Umeora JOU, Okogbenin SA, Otiode VO, Gharoro EP	2006	Nigeria (Benin)	Primary	193 nurses	Observational	Quantitative	Health center	3 months	medium
Peterman A & Johnson K	2009	Malawi, Rwanda, Uganda, Ethiopia	Secondary	35037 women of reproductive age (15-49 years)	Observational	Quantitative	Community	~1 year (DHS household surveys)	high
Reisel D, Creighton SM	2015	Multi-country	Review	NI	Other review	Qualitative	NA	NA	medium
Ruiz JI, Martinez AP, Bravo PMM, Roche PF	2012	Multi-country	Secondary	NI	Other review	Qualitative	NA	NA	medium
Rushwan H	2000	Multi-country	Review	NI	Other review	Qualitative	NA	NA	low
Ryan F	2012	Ethiopia	Primary	15 interviews with & participant observation of missionaries, local medical staff and fistula patients	Observational	Qualitative	NA	NA	medium
Saracoglu M, Zengin T, Ozturk H, Genc M	2014	Multi-country	Secondary	NI	Other review	Qualitative	NA	NA	low
Sharfi AR, Elmeqboul MA, Abdella AA	2013	Sudan (Khartoum focus)	Primary	2000 women of reproductive age (attending clinic & not)	Observational	Quantitative	Health center & university	5 years	medium

Slanger TE, Snow RC, Okonofua FE	2002	Nigeria (Edo)	Primary	1107 women attending FP or antenatal services	Observational	Quantitative	Health center	8 months	high
Tahzib F	1985	Nigeria (Northern)	Primary	80 fistula patients (<13 years of age)	Observational	Quantitative	Health center	11 years	medium
Teufel K, Dorfler DM	2013	Multi-country (Africa)	Primary	NI	Other review	Qualitative	NA	NA	low
Toubia N	1994	N/A	Review	NI	Other review	Qualitative	NA	NA	low
Tukur J, Jido TA, Uzoho CC	2006	Nigeria	Primary	50 fistula patients	Observational	Quantitative	Health center	4 years	medium
UNFPA and Engender Health	2003	Benin, Chad, Mali, Malawi, Mozambique, Niger, Nigeria, Uganda, Zambia	Primary	>35 facilities; government and UNFPA representatives & relevant stakeholders	Observational	Mixed	Health center	6 months	high
WHOa	2000	Multi-country	Secondary	422 articles	Systematic review	Qualitative	NA	NA	high
WHOb	2006	Burkina Faso, Ghana, Kenya, Nigeria, Senegal and Sudan	Primary	28393 women attending maternity services	Observational	Quantitative	Health center	17 months	high