

## $\beta$ -carotene in sunflower oil oxidation

By N. V. Yanishlieva, V. G. Raneva and E. M. Marinova

Institute of Organic Chemistry, Bulgarian Academy of Sciences,  
Kv. Geo Milev, Acad. G. Bontchev Str., Bl. 9, 1113 Sofia, Bulgaria  
E-mail: nelly@orgchm.bas.bg

### RESUMEN

#### $\beta$ -caroteno en la oxidación de aceite de girasol.

Se estudió la cinética de oxidación de aceite de girasol (SO), así como la de triacilglicerol puros de aceite de girasol (TGSO) en presencia de diferentes concentraciones (0.001-0.02 %) de  $\beta$ -caroteno. El proceso se llevó a cabo a altas (régimen cinético) y bajas (régimen de difusión) concentraciones de oxígeno a temperatura ambiente con luz natural y en la oscuridad, y se presentan también los resultados de la oxidación de SO y TGSO a 100°C en presencia de  $\beta$ -caroteno. Los resultados indicaron que en los sistemas lipídicos libres de antioxidantes, el  $\beta$ -caroteno no dió protección antioxidante. Funcionó como prooxidante durante la oxidación a temperatura ambiente y a concentraciones de oxígeno suficientemente elevadas, siendo el efecto más pronunciado en la oscuridad que con luz natural. El  $\beta$ -caroteno aumentó la estabilidad del SO conteniendo tocoferol durante su oxidación a temperatura ambiente y con luz natural. Este efecto es expresado más intensamente en un régimen cinético de oxidación. El sinergismo del  $\beta$ -caroteno con los tocoferoles se caracterizó por el factor de estabilidad F y la actividad A. En el régimen cinético de oxidación, F y A variaron en el intervalo F= 2.0-6.3, y A= 2.7-21.0. En el régimen de difusión F = 1.3-1.5, y A= 1.5-2.8.

**PALABRAS-CLAVE:** Aceite de girasol -  $\beta$ -caroteno - Oxidación.

### SUMMARY

#### $\beta$ -carotene in sunflower oil oxidation.

The oxidation kinetics of sunflower oil (SO), as well as of pure triacylglycerols of sunflower oil (TGSO) in the presence of different concentrations (0.001-0.02 %)  $\beta$ -carotene was studied. The process was performed at high (kinetic regime) and low (diffusion regime) oxygen concentrations at room temperature in the dark and under daylight. The results from the oxidation of SO and TGSO at 100°C in the presence of  $\beta$ -carotene were also presented. It was established that in the antioxidant-free lipid system, the  $\beta$ -carotene did not give any antioxidative protection. It worked as a prooxidant during the oxidation at room temperature and at sufficiently high oxygen concentration, the effect being more pronounced in the dark than under daylight.  $\beta$ -carotene increased the stability of tocopherol-containing SO during its oxidation at room temperature and under daylight. This effect is more strongly expressed in a kinetic regime of oxidation. The synergism of  $\beta$ -carotene with the tocopherols was characterized by the stabilization factor F and the activity A. In the kinetic regime of oxidation F and A varied in the interval F=2.0-6.3, and A =2.7-21.0. In the diffusion regime F=1.3-1.5, and A=1.5-2.8.

**KEY-WORDS:**  $\beta$ -Carotene - Oxidation - Sunflower oil.

### 1. INTRODUCTION

Carotenoids are present in a wide variety of food ingredients. They are mainly used as natural colorants in foods and in dietary fats (Bhagavan and Nair, 1992), and they have also antioxidative properties (Rajalakshi and Narasimhan, 1996).

Carotene research has progressed rapidly in the past few years.  $\beta$ -Carotene has the highest provitamin A activity of all carotenoids (Kotareddy and Devi, 1997). Burri (1997) recently discussed the important new developments in  $\beta$ -carotene research, and provided an interpretation of how these results reflect the impact of  $\beta$ -carotene on human health.  $\beta$ -Carotene reduces the risk of certain cancer (Peto *et al.*, 1981; Appel and Woutersen, 1996; Ziegler *et al.*, 1996; Hughes, 1998) and is high effective in preventing coronary cardiac disease (Seelert, 1991; Kohlmeier and Hastings, 1995).

The role of  $\beta$ -carotene in lipid oxidation was discussed many years ago. It was established that  $\beta$ -carotene increased the rate of oxygen absorption by fats (Olcott, 1934; Henk, 1941; Werner, 1942; Thompson and Steenbock, 1944; Holman, 1949). Wodsak (1951) reported that in the light  $\beta$ -carotene accelerated the fat oxidation, whereas in the dark it had an antioxidative effect. Herriset (1948) also found that  $\beta$ -carotene and vitamin A were inhibitors of spontaneous fat and oil oxidation.

Recently, however, contradictory results on the role of carotenoids, and in particular, of  $\beta$ -carotene, in lipid oxidation are also reported - antioxidative (Kiritsakis and Dugan, 1985; Fakourelis *et al.*, 1987; Lee and Min, 1990; Jung and Min, 1991), as well as prooxidative (Terao *et al.*, 1980; Faria and Mukai, 1983; Warner and Frankel, 1987; Haila and Heinonen, 1994) effects are observed.

The numerous investigations demonstrate that the behaviour of  $\beta$ -carotene in the oxidation process depends strongly on its concentration (Krinsky, 1968), on the environment (Pryor *et al.*, 1993), as well as on whether the process proceeds in the dark or in the light (Yanishlieva *et al.*, 1998). The anti-or prooxidative activity of  $\beta$ -carotene is also related closely to both oxygen concentration and to the

presence of other antioxidants (Krinsky, 1993; Palozza and Krinsky, 1993). Numerous reviews discuss the ability of  $\beta$ -carotene to act as a chain-breaking antioxidant and as a singlet oxygen quencher (Burton, 1989; Krinsky, 1989; Palozza and Krinsky, 1992a; Handelman, 1996).

There are no data in the literature concerning the role of  $\beta$ -carotene in sunflower oil oxidation at storage conditions. As known, this oil is one of the most commonly used valuable oils containing 60-70 % of the essential 9-cis,12-cis-octadecadienoic (linoleic) acid.

The objective of this study is to throw light on the influence of  $\beta$ -carotene on the kinetics of sunflower oil oxidation at room temperature.

The experiments were performed with commercially available sunflower oil (SO), as well as with pure triacylglycerols of sunflower oil (TGSO) at different storage conditions - in the dark and under daylight, at high (kinetic regime) and low (diffusion regime) oxygen concentrations. Different  $\beta$ -carotene concentrations (0.001-0.02 %) were investigated. Kinetic results from the autoxidation of SO and TGSO at 100°C (the most frequently used temperature for accelerated lipid stability determination) in the presence of  $\beta$ -carotene were also presented.

## 2. EXPERIMENTAL

### 2.1. Materials

$\beta$ -Carotene was from Merck (Darmstadt, Germany).

A commercially available sample of sunflower oil (SO) was used.

Pure triacylglycerols of sunflower oil (TGSO) were obtained by cleaning the sunflower oil from pro- and antioxidants and trace metals by adsorption chromatography (Popov *et al.*, 1968): 50 g lipid substrate in 500 ml distilled hexane were passed through a column (i.d. 2 cm) filled with 35g alumina (type 507C, neutral, Fluka AG, Buchs, Switzerland) activated for 4 h at 180°C, and collected in nitrogen in the dark. The solvent was removed in a rotary evaporator at 30°C in the dark. The product obtained was stored in an inert atmosphere at -20°C no more than 10 days.

Lipid samples containing 0.001-0.02 % ( $1.86 \times 10^{-5}$  -  $3.70 \times 10^{-4}$  M)  $\beta$ -carotene were prepared by adding aliquots of a solution of  $\beta$ -carotene in purified acetone to a weighted lipid sample followed by the removal of the solvent in nitrogen.

### 2.2. Methods

The fatty acid composition of SO was determined by gas chromatography of its methyl esters using a Pye Unicam instrument, model 304, equipped with a

dual flame-ionization detector and a glass capillary column (30 m x 0.2 mm i.d.) coated with SILAR 10C (Supelco Inc., Bellefonte, PA). The carrier gas was nitrogen at a flow rate of 14 ml min<sup>-1</sup>. The temperature was maintained at 165°C for 5 min, then increased to 220°C at 2°C/min.

The tocopherol content of the sunflower oil was determined by normal-phase HPLC (Ivanov and Aitzetmüller, 1995) using a Merck-Hitachi apparatus equipped with a pump L-6000 and a fluorescence detector Merck - Hitachi F-1050. A column with Nucleosil SI 50-5, 250x4 mm (Macherey - Nagel), and an elution system hexane: dioxane = 96:4 with a rate of 1 ml/min were used.

Oxidation at room temperature, 22°C ( $\pm 2^\circ\text{C}$ ) in the dark and under daylight (southern window) was carried out at oxygen sufficient conditions (kinetic regime), as well as at oxygen - insufficient conditions (diffusion regime). The kinetic regime was realized by storage of the samples in a 1 mm layers in Petri dishes having a diameter of 52 mm. The diffusion controlled oxidation was performed by storage of 10 g of the samples in glass jars with a diameter of 32 mm. Oil samples were thoroughly stirred before sampling.

Oxidation at 100°C ( $\pm 0.2^\circ\text{C}$ ) was carried out by blowing air through the samples (2 g) in the dark at a rate of 100 ml min<sup>-1</sup>.

The process was followed by withdrawing samples at measured time intervals and subjecting them to iodometric determination of the primary product (peroxide) concentration, i.e. the peroxide value (PV) (Yanishlieva *et al.*, 1978). As established, one of the most reliable methods for investigating the role of the carotenoids in lipid oxidation is the study of the peroxide accumulation (Terao *et al.*, 1992; Haila and Heinonen, 1994; Kasaikina, 1995; Tsuchihashi *et al.*, 1995; Servili *et al.*, 1996; Haila *et al.*, 1996). All kinetic curves are the mean result of two independent experiments.

The influence of  $\beta$ -carotene on lipid oxidation was estimated on the basis of the induction periods (IP) determined by the method of the tangents to the two parts of the kinetic curves (Yanishlieva and Popov, 1971; Le Tutour and Guedon, 1992). The oxidation rate of the control sample ( $W_0$ ) and of the samples with  $\beta$ -carotene ( $W_{\beta\text{-Car}}$ ) were found from the tangents to the initial phase of the kinetic curves of peroxide accumulation and expressed as M s<sup>-1</sup>. Recalculation of the rate from meq kg<sup>-1</sup> h<sup>-1</sup> into M s<sup>-1</sup> was performed according to the following formula (Marinova and Yanishlieva, 1992):

$$1 \text{ meq kg}^{-1} \text{ h}^{-1} = 1.4 \times 10^{-7} \text{ M s}^{-1}$$

## 3. RESULTS AND DISCUSSION

The fatty acid composition of TGSO was as follows: palmitate 6%, stearate 5%, oleate 25%,

linoleate 64%. The oil contained 0.062% tocopherols (Toc):  $\alpha$ -Toc 88.4%,  $\beta$ -Toc 3.9%,  $\gamma$ -Toc 5.1%,  $\delta$ -Toc 1.6%, and  $\alpha$ -tocotrienol ( $\alpha$ -Toc-3) 1.0%. The initial PV of the commercial sunflower oil was 13.0 meq  $\text{kg}^{-1}$ .

### 3.1. Triacylglycerols of sunflower oil (TGSO)

Figures 1 and 2 represent the kinetic curves of peroxide accumulation during oxidation of pure triacylglycerols of sunflower oil (TGSO) at room temperature at a sufficiently high oxygen concentration in the dark (Figure 1) and under daylight (Figure 2). Figures 1 and 2 show that  $\beta$ -carotene works as a prooxidant.

These results are in agreement with the data reported by other authors. Haila and Heinonen (1994) established that  $\beta$ -carotene concentrations between 0.005 and 0.2% accelerated hydroperoxide formation during autoxidation of purified rapeseed oil under light at 25°C. At the same concentrations Heinonen et al. (1996) found a prooxidative effect of  $\beta$ -carotene in a 10% purified rapeseed oil-in-water emulsion. The acceleration of autoxidation of different lipid systems by  $\beta$ -carotene at high oxygen pressure was observed also by Burton and Ingold (1984); Pryor *et al.* (1988), Kennedy and Liebler (1992), Jorgensen and Skibsted (1993), Tsuchihashi et al. (1995), Palozza *et al.*, (1995).

The prooxidative effect of  $\beta$ -carotene may be due to the oxidation of  $\beta$ -carotene alone leading to the formation of chain propagating radicals, and to the formation of products which act as initiators of the process (Kennedy and Liebler, 1992).




Figure 1

Kinetic curves of peroxide accumulation during oxidation of TGSO at room temperature in the dark at a sufficiently high oxygen concentration (kinetic regime) in the presence of different  $\beta$ -carotene concentrations: 0 - 0%; 1 - 0.001% ( $1.86 \times 10^{-5} \text{M}$ ); 2 - 0.005% ( $9.30 \times 10^{-5} \text{M}$ ); 3 - 0.01% ( $1.86 \times 10^{-4} \text{M}$ ); 4 - 0.02% ( $3.72 \times 10^{-4} \text{M}$ )




Figure 2

Kinetic curves of peroxide accumulation during oxidation of TGSO at room temperature under daylight at a sufficiently high oxygen concentration (kinetic regime) in the presence of different  $\beta$ -carotene concentrations: 0 - 0%; 1 - 0.001% ( $1.86 \times 10^{-5} \text{M}$ ); 2 - 0.005% ( $9.30 \times 10^{-5} \text{M}$ ); 3 - 0.01% ( $1.86 \times 10^{-4} \text{M}$ ); 4 - 0.02% ( $3.72 \times 10^{-4} \text{M}$ )




Figure 3

Dependence of the decrease of the induction period in the presence of  $\beta$ -carotene ( $IP_0/IP_{\beta\text{-Car}}$ ) on  $\beta$ -carotene concentration during oxidation of TGSO in a kinetic regime at room temperature in the dark (1) and under daylight (2).

After treatment of the curves in Figures 1 and 2 the dependence of the decrease of the induction period (IP) in the presence of  $\beta$ -carotene ( $IP_0/IP_{\beta\text{-Car}}$ ) on its concentration was obtained (Figure 3).

The induction period of TGSO without additive IPO in the kinetic regime was 38 h under daylight, and 7.7 days in the dark. It could be seen from Figure 3 that the prooxidative effect of  $\beta$ -carotene is much more pronounced in the dark than under daylight.

The results obtained during oxidation of TGSO at unsufficiently oxygen concentration (diffusion regime) showed that at both conditions, in the dark ( $IP_0 = 18$  days), as well as under daylight ( $IP_0 = 4.7$  days),  $\beta$ -carotene did not change the kinetics of the process (data not shown). The same was true when TGSO were oxidized at 100°C in the dark ( $IP_0 = 0.5$  h).

The data presented above show that  $\beta$ -carotene does not give any antioxidative protection during the oxidation of pure triacylglycerols of sunflower oil, e.g. of the antioxidant free lipid system.

### 3.3. Sunflower oil (SO)

Figures 4 and 5 illustrate the kinetic curves of peroxide accumulation during oxidation of SO at room temperature under daylight in a kinetic (Figure 4) and in a diffusion regime (Figure 5), respectively. One can see that at these conditions  $\beta$ -carotene retards the oxidation of SO, increasing the induction period  $IP$  and decreasing the rate of the process  $W$  in the initial phase.

For quantification of the retardation effect of  $\beta$ -carotene on SO oxidation we determined the following kinetic parameters characterizing the lipid oxidation during its initial stage: stabilization factor  $F$ , oxidation rate ratio  $ORR$  and activity  $A$  (Yanishlieva and Marinova, 1992).  $F$  is a measure of the effectiveness:

$$F = IP_{\beta\text{-Car}} / IP_0$$

where  $IP_{\beta\text{-Car}}$  is the induction period in the presence of  $\beta$ -carotene, and  $IP_0$  is the induction period of SO without additive.

$ORR$  is an inverse measure of the strength:

$$ORR = W_{\beta\text{-Car}} / W_0$$

where  $W_{\beta\text{-Car}}$  is the oxidation rate in the presence of  $\beta$ -carotene, and  $W_0$  is the initial oxidation rate of the control.




Figure 4

Kinetic curves of peroxide accumulation during oxidation of SO at room temperature under daylight at a sufficiently high oxygen concentration (kinetic regime) in the presence of different  $\beta$ -carotene concentrations: 0 - 0%; 1 - 0.001% ( $1.86 \times 10^{-5} M$ ); 2 - 0.005% ( $9.30 \times 10^{-5} M$ ); 3 - 0.01% ( $1.86 \times 10^{-4} M$ ); 4 - 0.02% ( $3.72 \times 10^{-4} M$ )




Figure 5

Kinetic curves of peroxide accumulation during oxidation of SO at room temperature under daylight at oxygen insufficient conditions (diffusion regime) in the presence of different  $\beta$ -carotene concentrations: 0 - 0%; 1 - 0.001% ( $1.86 \times 10^{-5} M$ ); 2 - 0.005% ( $9.30 \times 10^{-5} M$ ); 3 - 0.01% ( $1.86 \times 10^{-4} M$ ); 4 - 0.02% ( $3.72 \times 10^{-4} M$ )

The general parameter activity  $A$  unifies the effectiveness on the one hand, and the strength of the additive in the oxidation process, on the other:

$$A = F / ORR$$

After processing the kinetic curves in Figures 4 and 5 the kinetic parameters  $F$ ,  $ORR$  and  $A$  characterizing the oxidation of SO at room temperature under daylight in the presence of  $\beta$ -carotene were obtained (Table I). The data in Table I illustrate that  $\beta$ -carotene retards the oxidation of SO under light in a much higher degree when the process proceeds at a sufficiently high oxygen concentration (kinetic regime).

The experiments at room temperature in the dark showed that  $\beta$ -carotene did not influence the oxidation stability of sunflower oil in a kinetic ( $IP_0 = 48$  days), as well as in a diffusion regime ( $IP_0 = 140$  days). The same was true when SO was oxidized at  $100^\circ C$  in the dark ( $IP_0 = 7.3$  h). The results obtained with the thermal oxidation of safflower seed oil at  $75^\circ C$  in the presence of different concentrations of  $\beta$ -carotene (Henry *et al.*, 1998) were similar.

The presented data demonstrated that  $\beta$ -carotene increased the oxidation stability of tocopherol containing sunflower oil during its oxidation at room temperature and under daylight. It should be pointed out that the natural sensitizers in sunflower oil, e.g. the chlorophylls, may generate singlet oxygen during

Table I  
Kinetic parameters characterising the oxidation of sunflower oil at room temperature under daylight in the presence of  $\beta$ -carotene ( $\beta$ -Car),  $PV_o = 13 \text{ meq kg}^{-1}$

Conditions	$\beta$ -Carotene concentration		F	ORR	A
	[ $\beta$ -Car] (wt%)	[ $\beta$ -Car] $\times 10^5$ (M)			
Sufficiently high oxygen concentration (Kinetic regime) $IP_o = 2.2 \text{ days}$ $W_o = 1.93 \times 10^{-7} \text{ Ms}^{-1}$	0.001	1.86	2.0	0.75	2.7
	0.005	9.30	3.0	0.60	5.0
	0.010	18.60	4.1	0.43	9.5
	0.020	37.20	6.3	0.30	21.0
Oxygen insufficient conditions (Diffusion regime) $IP_o = 21.0 \text{ days}$ $W_o = 3.62 \times 10^{-8} \text{ Ms}^{-1}$	0.001	1.86	1.3	0.87	1.5
	0.005	9.30	1.4	0.70	2.0
	0.010	18.60	1.4	0.67	2.1
	0.020	37.20	1.5	0.53	2.8

oil oxidation under light (Foote and Denny, 1968; Korycka-Dahl and Richardson, 1978). The stabilizing effect of  $\beta$ -carotene observed may be due to its ability to quench the singlet oxygen (Raws and Van Santen, 1970; Foote, 1979; Kiritsakis and Dugan, 1985; Lee and Min, 1990), which explains its cooperative (synergistic) action with tocopherols.

The presented data are in accordance with the findings of other authors concerning the synergistic action of  $\beta$ -carotene with  $\alpha$ -tocopherol (Palozza and Krinsky, 1992b; Haila and Heinonen, 1994; Li *et al.*, 1995; Perrin *et al.*, 1996; Yin and Cheng, 1997),  $\gamma$ -tocopherol (Heinonen *et al.*, 1996), phospholipids (Bratkowska and Zwierzykowski, 1988), quercitrin (Maoka and Ito, 1994), tert-butylated hydroquinone (Tsai *et al.*, 1989), and 3,4-dihydroxyphenyl ethanol (Servili *et al.*, 1996).

#### ACKNOWLEDGEMENTS

One of the authors (NVY) is very grateful to the Alexander von Humboldt foundation for the financial support.

#### REFERENCES

- Appel, M. J. and Woutersen, R. A. (1996). Effect of dietary  $\beta$ -carotene and selenium on initiation and promotion of pancreatic carcinogenesis in azaserine-treated rats. *Carcinogenesis*, **17**, 1411-1416.
- Bhagavan, H. N., Nair, P. P. (1992). Antioxidants in dietary fats. *Food Sci. Technol.*, **53**, 329-336.
- Bratkowska I. and Zwierzykowski, W. (1988). Effect of carotenoids on the autoxidation of rapeseed oil with phospholipid additives. *Tluszczce Jadalne*, **26**, 3-12.
- Burri, B. J. (1997).  $\beta$ -Carotene and human health: A review of current research. *Nutr. Research*, **17**, 547-580.
- Burton, G. W. (1989). Antioxidant action of carotenoids. *J. Nutr.*, **119**, 109-111.
- Burton, G. W. and Ingold, K. U. (1984).  $\beta$ -Carotene: an unusual type of lipid antioxidant. *Science*, **224**, 569-573.
- Fakourelis, N., Lee, E.C. and Min, B. D. (1987). Effect of chlorophyll and  $\beta$ -carotene on the oxidation stability of olive oil. *J. Food Sci.*, **52**, 234-235.
- Faria, J. A. F. and Mukai, M. K. (1983). Use of a gas chromatographic reactor to study lipid photooxidation. *J. Amer. Oil Chem. Soc.*, **60**, 77-81.
- Foote, C. S. (1979). Quenching of Singlet Oxygen in Singlet Oxygen, p. 139-171, H. H. Wasserman and R. W. Murray (Eds.), Academic Press, New York.
- Foote, C. S. and Denny, W. R. (1968). Chemistry of singlet oxygen. VII. Quenching by  $\beta$ -carotene. *J. Amer. Chem. Soc.*, **90**, 6233-6235.
- Haila, K. and Heinonen, M. (1994). Action of  $\beta$ -carotene on purified rapeseed oil during light storage. *Food Sci. Technol.*, **27**, 573-577.
- Haila, K. W., Lievonen, S. M. and Heinonen, M. I. (1996). Effects of lutein, lycopene, annatto and  $\gamma$ -tocopherol on autoxidation of triglycerides. *J. Agric. Food Chem.*, **44**, 2096-2100.
- Handelman, G. J. (1996). Carotenoids as Scavengers of Active Oxygen Species in *Handbook of Antioxidants*, p. 259-314, E. Cadenas and L. Packer, (Eds.), Marcel Dekker, New York.
- Heinonen, M., Lampi, A., Piironen, V. and Haila, K. (1996). Interaction of Carotenoids and Tocopherols in Food Emulsions in *Oils-Fats-Lipids 1995*, Proceedings of the 21st World Congress of the International Society for Fat Research, The Hague, October 1995, p. 339, P. J. Barnes & Associates (Eds.), Bridgewater, United Kingdom.
- Henk, H. J. (1941). Die Stabilisierung von Fetten, Olen und Seifen. *Fette und Seifen*, **48**, 90-91.
- Henry, L. K., Catignani, G. L. and Schwartz, S. J. (1998). The influence of carotenoids and tocopherols on the

- stability of safflower seed oil during heat-catalyzed oxidation. *J. Amer. Oil Chem. Soc.*, **75**, 1399-1402.
- Herriset, A. (1948). Further study of the antioxygen properties of certain carotenoids. *Bull. Soc. Chim. Biol.*, **30**, 187-195.
- Holman, R. T. (1949). Spectrophotometric studies of the oxidation of fats. VIII. Coupled oxidation of carotene. *Arch. Biochem.*, **21**, 51-57.
- Hughes, D. A. (1998). Beta-carotene and carotenoids: antioxidants and their role in human health. *Pharm. News*, **5**, N<sup>o</sup>4, 7-10.
- Ivanov, St. A. and Aitzetmüller, K. (1995). Untersuchungen über die Tocopherol- und Tocotrienolzusammensetzung der Samenöle einiger Vertreter der Familie Apiaceae. *Fat Sci. Technol.*, **97**, 24-29.
- Jorgensen, K. and Skibsted, L. H. (1993). Carotenoid scavenging of radicals. Effect of carotenoid structure and oxygen pressure on antioxidative activity. *Z. Lebensm. Unters. Forsch.*, **196**, 423-429.
- Jung, M. Y. and Min, D. B. (1991). Effects of quenching mechanisms of carotenoids on the photosensitized oxidation of soybean oil. *J. Amer. Oil Chem. Soc.*, **68**, 653-658.
- Kasaikina, O. T. (1995). Oxidation of polyene hydrocarbons. *Khim. Fiz.*, **14**, 72-85.
- Kennedy, T. A. and Liebler, D. C. (1992). Peroxyl radical scavenging by  $\beta$ -carotene in lipid bilayers. Effect of oxygen partial pressure. *J. Biol. Chem.*, **267**, 4658-4663.
- Kiritsakis, A. and Dugan, L. R. (1985). Studies in photooxidation of olive oil. *J. Amer. Oil Chem. Soc.*, **62**, 892-896.
- Kohlmeier, L. and Hastings, S. B. (1995). Epidemiologic evidence of a role of carotenoids in cardiovascular disease prevention. *Am. J. Clin. Nutr.*, **62**, S1370-S1376.
- Korycka-Dahl, M. B. and Richardson, F. (1978). Activated oxygen species and oxidation of food constituents. *CRC Crit. Rev. Food Sci. Nutr.*, **10**, 209-241.
- Kotareddy, B. and Devi, K. (1997). Development of  $\beta$ -carotene-rich extruded maize products. *J. Food Sci. Technol.*, **34**, 63-66.
- Krinsky, N. I. (1968). Protective function of carotenoid pigments. *Photophysiology*, **3**, 123-195.
- Krinsky, N. I. (1989). Antioxidant function of carotenoids. *Free Radical Biol. Med.*, **7**, 617-635.
- Krinsky, N. I. (1993). Action of carotenoids in biological systems. *Ann. Rev. Nutr.*, **13**, 561-587.
- Lee, S.-W. and Min, D. B. (1990). Effects, quenching mechanisms, and kinetics of carotenoids in chlorophyll-sensitized photooxidation of soybean oil. *J. Agric. Food Chem.*, **38**, 1630-1634.
- Le Tutour, B. and Guedon, D. (1992). Antioxidative activities of *Olea europaea* leaves and related phenolic compounds. *Phytochemistry*, **31**, 1173-1178.
- Li, Z.-L., Wu, L.-M., Ma, L.-P., Liu, Y.-C. and Liu, Z.-L. (1995). Antioxidant synergism and mutual protection of  $\alpha$ -tocopherol and  $\beta$ -carotene in the inhibition of radical-initiated peroxidation of linoleic acid in solution. *J. Phys. Org. Chem.*, **8**, 774-780.
- Maoka, T. and Ito, Y. (1994). Synergistic antioxidant effect of  $\beta$ -carotene and quercetin. *Igakuto Seibutsugaku*, **129**, 63-73.
- Marinova, E. M. and Yanishlieva, N. V. (1992). Effect of temperature on the antioxidative action of inhibitors in lipid autoxidation. *J. Sci. Food Agric.*, **60**, 313-318.
- Olcott, H. J. (1934). Antioxidants and autoxidation of fats. II. *J. Amer. Chem. Soc.*, **56**, 2492-2493.
- Palozza, P., Calviello, G. and Bartoli, G. M. (1995). Prooxidative activity of  $\beta$ -carotene under 100% oxygen in liver microsomes. *Free Radical Biol. Med.*, **19**, 887-892.
- Palozza, P. and Krinsky, N. I. (1992a). Antioxidant effects of carotenoids in vivo and in vitro: an overview. *Methods Enzymol.*, **213**, 403-420.
- Palozza, P. and Krinsky, N. I. (1992b).  $\beta$ -Carotene and  $\alpha$ -tocopherol are synergistic antioxidants. *Arch. Biochem. Biophys.*, **297**, 184-187.
- Palozza, P. and Krinsky, N. I. (1993). Antioxidant effects of carotenoids on biological membranes. *Acta Med. Rom.*, **31**, 131-140.
- Perrin, D., Eloualja, H. and Martin, R. (1996). Antioxidant properties of all-trans- $\beta$ -carotene/ $\alpha$ -tocopherol mixtures in the induced oxidation of ethyl linoleate. *J. Chim. Phys. Phys. - Chim. Biol.*, **93**, 1462-1471.
- Peto, R., Doll, R., Buckley, J. D. and Sporn, M. B. (1981). Can dietary  $\beta$ -carotene materially reduce human cancer rates? *Nature*, **290**, 201-208.
- Popov, A., Yanishlieva, N. and Slavceva, J. (1968). Methode zum Nachweis von Antioxidantien in Methyloleate für kinetische Untersuchungen. *Compt. Rend. Acad. Bulg. Sci.*, **21**, 443-446.
- Pryor, W. A., Cornicelli, J. A., Devall, L. J., Tait, B., Trivedi, B. K., Witiak, D. T. and Wu, M. D. (1993). A rapid screening test to determine the antioxidant potencies of natural and synthetic antioxidants. *J. Org. Chem.*, **58**, 3521-3532.
- Pryor, W. A., Strickland, T. and Church, D. F. (1988). Comparison of the efficiencies of several natural and synthetic antioxidants in aqueous sodium dodecyl sulphate micelle solutions. *J. Amer. Chem. Soc.*, **110**, 2224-2229.
- Rajalakshi, D. and Narasimhan, S. (1996). Food Antioxidants: Sources and Methods of Evaluation in *Food Antioxidants*, p. 65-157. D. L. Madhavi, S. S. Deshpande and D. K. Salunke (Eds.), Marcel Dekker, New York.
- Raws, H. R. and Van Santen, P. J. (1970). A possible role for singlet oxygen in the initiation of fatty acid autoxidation. *J. Amer. Oil Chem. Soc.*, **47**, 121-125.
- Seelert, K. (1991).  $\beta$ -Carotin. Hochwirksam bei koronaren Herzerkrankungen. *Deutsche Apoth. Zeitung*, **131**, 72-76.
- Servili, M., Baldioli, M., Miniati, E. and Montedoro, G. F. (1996). Antioxidant activity of new phenolic compounds extracted from virgin olive oil and their interaction with  $\alpha$ -tocopherol and  $\beta$ -carotene. *Riv. Ital. Sost. Grasse*, **73**, 55-59.
- Terao, J. A., Nagao, A., Park D.-Ki and Lim, B. P. (1992). Lipid hydroperoxide assay for antioxidant activity of carotenoids. *Methods Enzymol.*, **213**, 454-460.
- Terao, J., Yamauchi, R., Murakami, H. and Matsushita, S. (1980). Inhibitory effects of tocopherols and  $\beta$ -carotene on singlet oxygen initiated photooxidation of methyl linoleate and soybean oil. *J. Food Process. Preserv.*, **4**, 79-93.
- Thompson, C. R. and Steenbock, H. (1944). The effect of  $\alpha$ -tocopherol and  $\beta$ -carotene in the oxidation of plant and animal fats. *Arch. Biochem.*, **4**, 15-23.
- Tsai, R. S., Huang, J. J. and Chang, R. L. (1989). The inhibitory effect of  $\beta$ -carotene on the photooxidation of peanut oil. *J. Chin. Agric. Chem. Soc.*, **27**, 233-242.
- Tsuchihashi, H., Kigoshi, M., Iwatsuki, M. and Niki, E. (1995). Action of  $\beta$ -carotene as antioxidant against lipid peroxidation. *Arch. Biochem. Biophys.*, **323**, 137-147.

- Warner, K. and Frankel, E. N. (1987). Effects of  $\beta$ -carotene on light stability of soybean oil. *J. Amer. Oil Chem. Soc.*, **64**, 213-218.
- Werner, H. (1942). Das Verderben von tierischen und pflanzlichen Fetten, insbesondere Ranzigkeit und dessen Vermeidung. *Z. Fleisch- und Milchhyg.*, **52**, 207-211.
- Wodsak, W. (1951). Antioxidantien für fettlösliche Vitamine. *Fette und Seifen*, **53**, 73-76.
- Yanishlieva, N. V., Aitzetmüller, K. and Raneva, V. G. (1998).  $\beta$ -Carotene and lipid oxidation. *Fett/Lipid*, **100**, 444-462.
- Yanishlieva, N. V., and Marinova, E. M. (1992). Inhibited oxidation of lipids I. Complex estimation and comparison of the antioxidative properties of some natural and synthetic antioxidants. *Fat Sci. Technol.*, **94**, 374-379.
- Yanishlieva, N. and Popov, A. (1971). Über einige Eigentümlichkeiten in der Kinetik zu Beginn der Autoxydation von Estern ungesättigter Fettsäuren 1.Mitt. Abhängigkeit des Mechanismus der Kettenbildung und der Initiierungsgeschwindigkeit im Methyloleat und Methyllinoleat von der Temperatur und vom Hydroperoxidgehalt, *Die Nahrung*, **15**, 395-402.
- Yanishlieva, N, Popov, A. and Marinova, E. (1978). Eine modifizierte Methode zur Bestimmung der Peroxidzahl in kleinen Lipidproben, *Compt. Rend. Acad. Bulg. Sci.*, **31**, 869-871.
- Yin, M.-Ch. and Cheng, W-Sh. (1997). Oxymyoglobin and lipid oxidation in phosphatidylcholine liposomes retarded by  $\alpha$ -tocopherol and  $\beta$ -carotene. *J. Food Sci.*, **62**, 1095-1097.
- Ziegler, R. G., Colavito, E. A., Hairge, P., Mcadams, M. J., Schoenberg, J. B., Mason, T. J. and Fraumeni, J. F. (1996). Importance of  $\alpha$ -carotene,  $\beta$ -carotene, and other phytochemicals in the etiology of lung cancer. *J. Natl. Cancer Inst.*, **88**, 612-615.

Recibido: Mayo 1999  
Aceptado: Febrero 2000