

Enzymatic aqueous technology for simultaneous coconut protein and oil extraction

By **Beatriz P.M. Sant'Anna**, **Suely P. Freitas**¹ and **Maria A. Z. Coelho**²

¹ E-mail: freitasp@unisys.com.br

² Escola de Química, Universidade Federal do Rio de Janeiro, CT, sl. E-113, 21949-900, Rio de Janeiro - RJ, Brazil

RESUMEN

Tecnología enzimática para la extracción simultánea de aceite y proteína de coco.

En este trabajo diferentes enzimas hidrolíticas fueron usados para la extracción de aceite y proteína de coco: *Celluclast*, *Termamyl*, *Viscozyme*, *Neutrast* y *Protease* producidos por Novo Nordisk. Experimentos preliminares fueron encaminados a la selección de los extractos enzimáticos, de las concentraciones de las enzimas y de los tiempos de incubación. La técnica de análisis de varianza, mediante un delineamiento factorial en tres niveles, fue usada para la optimización de los procesos con relación a los parámetros pH y razón sustrato/agua. Comparando con la extracción acuosa todos los enzimas hidrolíticos incrementan la extracción de aceite y proteína vegetal de coco. Los coeficientes de correlación muestran que el pH fué el parámetro más significativo ($p < 0,10$). Las condiciones óptimas para la extracción de aceite y proteína fueron: concentración de las enzimas *Viscozyme* y *Neutrast* 0.6% (w/w) y 0.3% (w/w), respectivamente, tiempo de incubación 30 minutos, razón sustrato/agua 1:6 y pH 7.

PALABRAS-CLAVE: Aceite de coco – Enzima – Extracción acuosa – Proteína de coco.

SUMMARY

Enzymatic aqueous technology for simultaneous coconut protein and oil extraction.

In this study, the following commercial enzymes were evaluated in the enzymatic extraction of oil and protein from coconut: *Celluclast*, *Termamyl*, *Viscozyme*, *Neutrast* y *Protease*. Preliminary experiments were carried out for selection of enzymes, enzyme concentration and incubation time. In order to verify the effects of pH and water/substrate ratio, a response surface methodology was applied based in a 3² full factorial experimental design. The pH is the most meaningful parameter on oil and protein extraction yields, with a significance level higher than 90%. The maximal extraction yields of oil and protein emulsion (83%) was reached using *Viscozyme* L and subsequently *Neutrast* 1.5 MG at concentrations of 0.6% (w/w) and 0.3% (w/w), respectively, total incubation time of 60 minutes, substrate/water ratio 1:6 and pH around 7 (no adjustment).

KEY-WORDS: Aqueous extraction – Coconut oil – Coconut protein – Enzyme.

1. INTRODUCTION

Coconut oil is one of the most stable oils what is attributed to its high proportion (90%) of saturated

fats (Inform, 1994). About half of fatty acids in coconut oil are short chains (lauric, myristic and palmitic acids). These medium chain length acids and their derivatives are hydrophilic, which promotes interfacial properties that are valuable in detergents, lubricants, cosmetics and other emulsions applications (Seguro & Motoki, 1994).

Conventional processes for the extraction of coconut oil involve mechanical treatment, by pressing coconut meat, and a solvent step, that submit the cake for further extraction with n-hexane (Christensen, 1991). Although these technologies are economically justifiable, they have certain well-known drawbacks: damage to the environment and quality loss of finished products (e.g. high free fatty acids and lower resistance to rancidity).

Several studies have been carried out on aqueous enzymatic oil extraction (Dominguez *et al.*, 1994; Rosenthal *et al.*, 1996). The enzymatic extraction of vegetable oils, developed at Embrapa - Food Agroindustry reach the high yield extraction of tropical fruits' oil (avocado, pupunha, pequi, tucumã) when the incubation process is based on a combination of pectinolytic enzymes (Freitas *et al.* 1993). Furthermore, the high yield extraction of seeds oils is obtained on a combination of cellulase and protease enzymes (Freitas *et al.*, 1996).

The enzyme-assisted coconut oil extraction significantly raises the extracted oil yield in aqueous systems (Tano-Debrah, 1997). The main limitation of this process is the low efficiency of the oil separation due to stability of the lipoprotein emulsions. On the other hand, the most important advantage is associated with the simultaneous production of oil and hydrolyzed protein, both largely used in cosmetics formulations. Furthermore, this process allows the removal of the impurities in the aqueous phase.

The present work investigates the effect of relevant operational parameters including enzyme concentration, incubation time, pH and substrate/water ratio on coconut protein and oil extraction yield.

2. MATERIALS AND METHODS

Coconut meat used for all experiments was obtained from a local supplier at Rio de Janeiro, Brazil. The proximate composition of the coconut meat and obtained protein-oil emulsions were determined by AOAC methods (1995).

Commercial enzymatic complexes, donated by Novo Nordisk A/S, were employed:

- *Viscozyme L* (100 FBG/g), a multienzyme complex with cellulase, hemicellulase, arabanase, xylanase and -glucanase activities;
- *Termamyl 120 L* (120 KNU/g), an endoamylase which will hydrolyze 1,4- α - glucosidic linkages in amylose and amylopectin;
- *Pectinex 3XL* (3000 FDU/g) from *Aspergillus niger*, with pectinolytic effect;
- *Celluclast 1.5L* (1500 NCU/g), a cellulase preparation;
- *Alcalase* (2.4 AU/g) and *Neutrase 1.5MG* (1.5 AU/g), respectively an alkaline and a neutral protease produced from *Bacillus amyloliquefaciens*.

2.1. Enzyme-assisted protein-oil extraction

For each experiment 100 g of coconut meat was triturated and blended with warm water (45°C). The mixture was squeezed and strained through a porous membrane. The milk obtained was heated to 100°C for 30 min and left to cool. The solid residue was blended with water for 5 min and the mixture was submitted to enzymatic treatment. The enzymatic reactions were carried out with a pectinolytic or cellulolytic enzyme at 50°C prior the incubation with a protease at 60°C, in a shaker under temperature and agitation control. After this step, the mixture was again filtered through a porous membrane, originating a second coconut milk. This milk was mixed with the previously extracted coconut milk and allowed to stand, undisturbed, for 1 hour in order to separate the oily layer. The aqueous layer was drained off and the oil-protein-water mixture was centrifuged at 5000 rpm and 12°C during 45 min. This procedure resulted in frozen protein-oil emulsion, released by the enzymatic reaction, which could be then separated from the remaining aqueous layers. The yield was determined by its dry weight in 100g of coconut meat (c.m.). Figure 1 illustrates this procedure.

2.2. Statistical analysis

Preliminary experiments were developed for selection of enzymes, enzyme concentration, time reaction and incubation parameter levels. Response surface methodology was applied to investigate the effects of pH and water/substrate ratio, based in a 3² full factorial experimental design.




Figure 1
Experimental procedure for coconut protein-oil emulsion extraction in aqueous enzymatic media.

3. RESULTS AND DISCUSSION

Results from proximate composition of coconut meat and emulsion obtained indicate that a protein fraction, transferred to aqueous phase during enzymatic process, remains bind to the oil as a lipoprotein emulsions (Table I).

a) Enzyme Selection: Initially, an enzyme selection was carried out based on combinations of protease (*Neutrase 1.5MG*) and above-mentioned pectinolytic and cellulolytic enzymes. Experiments were carried out in duplicate, with substrate-water ratio of 1:6, during 30 minutes for each enzymatic extract employed. The first enzymatic step, at 50°C, employed 0.3% of hydrolytic enzyme and the second one, at 60°C, 0.1% of protease. The selected enzyme combination, as well as the results obtained, are shown in Figure 2. In all cases the enzymatic treatment increased the protein-oil yield as compared with control. The combination between *Viscozyme* and *Neutrase* was selected as the most suitable enzymatic treatment reaching an increase of about 60% with reference to the control.

New experiments were performed in the same conditions, employing *Viscozyme L* (0.3%) and replacing *Neutrase* by *Alcalase* (0.1%). No significant difference in protein-oil yield was obtained (Figure 2). The *Neutrase* present the higher activity in the natural pH of the coconut emulsion and was applied for further incubation parameter selection.

Table I
Proximate composition of coconut meat and protein-oil emulsion (dry base, g/100g)

	Coconut	Protein-Oil emulsion
Oil	46.41	80.88
Protein	8.20	13.38
Ash	1.68	0.38
Fibers	33.78	1.04
Carbohydrates*	9.88	4.33

*By difference

b) Enzyme Concentration and Incubation

Time: Figure 3 shows the simultaneous effects of the concentrations and incubation times in the yield process for the *Viscozyme* and *Neutralse* enzymes. The maximum protein-oil emulsion yield, 50.4 g by 100 g of the substrate (92%) was observed by enzymatic treatment with 0.6% *Viscozyme L* and 0.3% *Neutralse*, during incubation time of 30 minutes each in the natural pH (Figure 3). High levels of cellulase and hemicellulase present in *Viscozyme L*, enabled to release the oil and the protein to the aqueous media, due to the rupture of the cell walls that involve the oil globule. Treatment with 0.6% *Viscozyme L* and 0.1% *Neutralse* was too effective, although, in this case, the protein-oil emulsion obtained presented higher stability due to lower concentration of protease employed. Protease contributes to enhancing oil extraction, since it breaks the lipoprotein emulsions. Thirty minutes incubation time for each treatment was more efficient in the majority of the combinations tested.

c) pH and Substrate/Water Ratio: The pH and substrate/water ratio effects was based in a full factorial experimental design 3², in the previous




Figure 2

Selection for hydrolytic enzymes: 0.3% of Pectinase or Cellulase enzyme (50°C) and 0.1% of Neutralse (60°C), in all experiments. P - Pectinex 3XL; C - Celluclast; T - Termamyl 120L; V - *Viscozyme,L*




Figure 3

Effects of the incubation time and enzyme concentration in coconut protein-oil emulsion extraction. V - *Viscozyme*; N - *Neutralse*

selected conditions: 0.6% of *Viscozyme L*, 0.3% of *Neutralse 1.5MG* and 30 minutes of incubation time for each enzymatic treatment. Three different pH levels were tested: 6, 7 and 9, where 7 correspond to the emulsion pH without any adjustment. To obtain the extreme values, NaOH 1N and HCl 1N solutions were added.

It was possible to conclude that an increase in pH has a negative effect on the enzymatic extraction of coconut protein-oil yield. This is due to higher activities of selected enzymes at natural pH. As can be observed in response surface plots (Figure 4), the pH profile presents an accentuated curve if compared to the substrate:water one, showing strong dependence between the process yield and pH (significance level superior to 94%, according to variance analysis). The best condition corresponding to the maximum oil yield was reached in the natural pH of the mixture and

Fitted Surface; Variable: Protein-oil YIELD (%)
 2 3-level factors, 9 Runs;




Figure 4

Response Surface for protein-oil emulsion yield as a function of the water/substrate ratio and pH. 0.6% of *Viscozyme L*, 0.3% of *Neutralse 1.5MG* and 30 minutes of incubation time for each enzymatic treatment.

substrate:water ratio 1:9. However, the oil yield at substrate:water ratio 1:9 was slightly superior if compared with substrate:water ratio 1:6.

4. CONCLUSIONS

This work allowed selecting enzymes and determining extraction parameters that provide best coconut protein-oil yield in the aqueous process. It was possible to conclude that pH is the most meaningful parameter of the enzymatic process, with a significance level higher than 90%.

The results pointed to the selection of *Viscozyme* L and *Neutralse 1.5 MG* at concentrations of 0.6% (w/w) and 0.3% (w/w), respectively, incubation time of 30 minutes, substrate/water ratio 1:6 and pH around 7 (no adjustment).

The aqueous enzyme-assisted extraction method has shown to be an effective technique to obtain a higher oil yield. This process generates a practically nonfat fiber, thus proving its efficacy.

ACKNOWLEDGMENTS

The authors would like to thank the FAPERJ - Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro and CNPq - Conselho Nacional de Desenvolvimento Científico e Tecnológico for providing the scholarship. The authors are also grateful to Novo Nordisk which gently provided commercial enzymes.

REFERENCES

- AOAC. Official methods of analysis of the Association of Official Analytical Chemists. 40.ed. Washington, DC, 1995. cap.33, p. 10.
- Christensen, F.M. Extraction by aqueous enzymatic process. Inform: International News on Fat, Oils and related Materials. New York, **2**, n.11, 984 - 987, 1991.
- Dominguez, H.M.; Núñez, J. and Lema, J.M. Enzymatic pretreatment to enhance oil extraction from fruits and oil seeds: a review. Food Chemistry, **49**, 271-286, 1994.
- Freitas, S., Lago, R.C.A., Jablonka, F.H. and Hartman, L. Extraction aqueuse enzymatique de l'huile d'avocat a partir de la pulpe fraîche. Revue Française des Corps Gras, França, **41**, 365-371, 1993.
- Freitas, S.; Hartman, L. and Couri, S. The combined application of extrusion and enzymatic technology for extraction of soya bean oil. Alemanha, Fett and Lipid, Alemanha, **99**, 333-337, 1996.
- Rosenthal, A., Pyle L.D. and Niranjana, K. Aqueous and enzymatic processes for edible oil extraction. Enzyme Microbiology Technology, **19**, Inglaterra, novembro, 402-420, 1996.
- Seguro, K. and Motoki, M. Trends in Japanese soy protein research. Inform, International News on Fat, Oils and related Materials, 5, n. 3, 308-313, 1994.
- Tano-Debrah, K. and Obta, Y. Aqueous extraction of coconut oil by an enzyme assisted- process. Journal Science Food Agricultural, **74**, 497-502, 1997.

Recibido: Marzo 2002
Aceptado: Septiembre 2002