

HISTO-ANATOMICAL AND PRELIMINARY TLC INVESTIGATIONS ON HYPERICUM HIRSUTUM L. (HYPERICACEAE) SPECIES

GEORGE DAN MOGO ANU¹, CORNELIA BEJENARU², ANDREI BI^{1*},
LUDOVIC EVERARD BEJENARU¹

¹Department of Pharmacognosy & Phytotherapy, Faculty of Pharmacy, University of Medicine and Pharmacy of Craiova, e-mail: andreibita@gmail.com

²Department of Vegetal & Animal Biology, Faculty of Pharmacy, University of Medicine and Pharmacy of Craiova

Keywords: *Hypericum hirsutum* L., histo-anatomy, polyphenols, thin layer chromatography

ABSTRACT

Starting from *Hypericum hirsutum* L. plants in blossom, harvested in May 2016, from Râ ca Forest, Olt County (Romania), the cross-sections of root, aboveground stem and leaf were acquired and examined through microphotography technique. Also, the content of polyphenols was determined by thin layer chromatography as part of a comprehensive pharmacognostic analysis of aerial parts (*Hyperici hirsuti herba*). From the 12 separated chromatographic bands, one chlorogenic acid derivative was identified.

INTRODUCTION

Hypericum hirsutum L., Hairy St. John's wort, *Hypericaceae* family, is a perennial species spontaneous in Europe and Asia, growing mainly in forest edges and scrubs. It has downy stems and much longer leaves, compared to the most commonly known and used *H. perforatum* species [6].

Hypericum species contain some active principles (naphthodianthrones, flavonoids, essential oil, prenylated phloroglucinols, xanthonenes, proanthocyanidins, catechic tannin) [3, 7, 8, 13, 18, 21] with peculiar pharmacological activities: antidepressant [16], biostimulating, cicatrizing, antimicrobial [3, 5, 13, 14], antiviral [19], anti-inflammatory [17], antioxidant, anti-acetylcholinesterase [12], antieczematous, antihemorrhagic [3, 19], leishmanicidal [9].

In the specialty papers, there are scarce and incomplete data concerning *H. hirsutum* histo-anatomy [10, 15] and chemical composition [7, 8, 21].

The aim of our paper was the histo-anatomical investigation of the root, aboveground stem and leaf of the *H. hirsutum* species and the preliminary analysis of the polyphenols content from the aerial parts (*Hyperici hirsuti herba*).

MATERIAL AND METHOD

Histo-anatomical investigation

The vegetal material was collected from *H. hirsutum* plants in blossom, in May 2016, from Râ ca Forest, Olt County, Romania.

The fixation and preservation of roots, aboveground stems and leaves were achieved in 70% ethanol. The cross-sections and longitudinal-radial sections were obtained using botanical razor. After washing with distilled water, the sections were clarified using 10% sodium hypochlorite solution (Javel water). Then, the clarifying agent was removed by washing with distilled water. Congo red–chrysoidine mixture (Genevese reagent) was used for the staining of sections. Depending on the chemical composition of cell membranes, the reactive induced various stains: pink to red for cellulose and mucilage, pale red for cytoplasm, yellow for suberin and brown for lignin [2].

Stained and mounted sections were analyzed on a Krüss binocular photon microscope (objectives x4, x10, x40) and then photographed using a Sony DSLR-A380 digital system adapted to the microscope. The description of microscopic sections was accomplished according to some classical authors [20].

Thin-layer chromatography (TLC) analysis

Preliminary analysis of polyphenols was performed on the aerial parts of *H. hirsutum* species (*Hyperici hirsuti herba*), by using a CAMAG (Muttentz, Switzerland) system in the following experimental conditions [1, 4, 11]: stationary phase TLC silica gel 60 F₂₅₄ 20×10 cm precoated glass plates (Merck, Darmstadt, Germany) pre-washed with chloroform–methanol (1:1, v/v); mobile phase chloroform–ethyl acetate–toluene–formic acid–methanol (15:20:10:10:1, in volumes) in a vapor-equilibrated chromatographic tank (20×10 cm twin trough chamber, CAMAG); sample – 20% methanolic extract of *Hyperici hirsuti herba*; standards (Merck) – 0.05% methanolic solutions of caffeic acid, chlorogenic acid, quercetin and rutin; migration distance 80 mm; sample (1–10 µL) and standards (2 µL) application – CAMAG Linomat 5 semiautomatic system (spray gas nitrogen, dosage speed 150 nL/s, band length 8 mm); detection – CAMAG TLC Scanner 3 photodensitometer, UV 254 nm, without derivatization, deuterium–wolfram lamp, scanning speed 20 mm/s, data resolution 100 µm/step; measurement mode – absorption; spectra acquisition, processing, and quantification analysis – winCATS ver. 1.4.3 software package.

RESULTS AND DISCUSSIONS

Histo-anatomical investigation

Root

In cross-section, the root has circular shape and secondary structure due to the two secondary meristematic zones: phellogen and libero-ligneous cambium. From the outside towards the inside of the root, the following histological sequence was highlighted in cross-section. Peridermis is made of suber, phellogen and phelloderm. Suber is composed of 4–5 layers of suberin-impregnated flattened cells. From point to point, suber is exfoliated. One layer of subero-phellodermic cambium has anterior-posterior flattened cells, thin-walled, the radial walls being slightly undulated. Phelloderm is made of 2–3 cellular layers with slightly collenchymatized walls. Conducting tissues are arranged in two concentric rings. Phloem tissue forms a thin external ring, made of sieve tubes, phloem parenchyma and annex cells. Libero-ligneous cambium is disposed between xylem and phloem tissues. Xylem tissue forms the internal ring made of metaxylem vessels with different caliber, disorderly dispersed into the libriform tissue. Thus, the protoxylem vessels with small diameter, accompanied by few xylem parenchyma cells, are pushed toward the center of the root. Metaxylem vessels show reticulate thickenings, on longitudinal-radial sections. Medullary rays are multi-cellular, uniseriate, cellulose, at the level of phloem tissue ring but lignified to the xylem tissue ring. Medullary parenchyma is missing (Figures 1 and 2).



Figure 1. Cross-section through *H. hirsutum* root: (a) metaxylem; (b) libriform tissue; (c) medullary ray (Congo red–chrysoidine staining, ×400).



Figure 2. Longitudinal-radial section through *H. hirsutum* root: (a) metaxylem with reticular thickenings (Congo red–chrysoidine staining, $\times 400$).

Aboveground stem

In cross-section, the aboveground stem has circular shape and secondary structure generated by the libero-ligneous cambium. Epidermis is made of big heterodiametric cells with thickened external wall covered by a tick cuticle. The epidermal cells are tangentially elongated with thin radial walls and thickened tangential external and internal walls. From point to point, we found stomata and unicellular long tector hairs with sharp peak. The cortex is made up of two parts: external area of 2–3 layers of chlorenchyma and internal area, well represented, from parenchymatic type. The conducting tissues are arranged in two concentric rings. Phloem tissue forms a thin, external ring consisting of sieve tubes, phloem parenchyma and annex cells. Libero-ligneous cambium is situated between xylem and phloem tissues. Xylem tissue forms the internal ring composed of metaxylem vessels with various caliber, disorderly scattered into the libriform mass, pushing toward the center protoxylem vessels with small diameter, accompanied by few xylem parenchyma cells. In longitudinal-radial sections, xylem vessels exhibit reticulate and spiral thickenings. The medullary rays are multi-cellular, uniseriate, cellulosic at the level of phloem tissue ring but lignified into the xylem tissue ring. Medullary parenchyma is well developed, of meatus type (Figures 3–5).




Figure 3. Cross-section through *H. hirsutum* aboveground stem: (a) metaxylem; (b) libriform tissue; (c) medullary ray (Congo red–chrysoidine staining, $\times 400$).




Figure 4. Cross-section through *H. hirsutum* aboveground stem: (a) metaxylem; (b) libriform tissue; (c) protoxylem; (d) xylem parenchyma; (e) medullary parenchyma (Congo red–chrysoidine staining, $\times 400$).



Figure 5. Longitudinal-radial section through *H. hirsutum* aboveground stem: (a) metaxylem with spiral thickening; (b) metaxylem with reticular thickening (Congo red–chrysoidine staining, $\times 400$).

Leaf's limb

At the level of median rib, leaf's lamina is prominent on abaxial side. In cross-section, from the outside towards the inside of leaf's limb, the following histological sequence is observed. Upper epidermis is composed of a single layer of large cells, flattened, with thickened tangential external and internal walls and thin radial walls. External walls are bulged and covered by a thick cuticle. At this level, from point to point, are found long unicellular tector hairs with sharp peak. Mesophyll is made of a single layer of palisade parenchyma consisting of large, cylindrical cells, rich in chloroplasts, with small intercellular spaces, as well as of 3–4 layers of lacunose parenchyma composed of small cells, disorderly arranged, with large aeriferous spaces. Into the mesophyll are found many small libero-ligneous conducting fascicles, surrounded by assimilatory fascicular sheaths. At this level, cystoliths of calcium oxalate are also observed. Mesophyll has bifacial dorsiventral structure. Lower epidermis is made of a single layer of small cells, anterior-posterior slightly flattened, with thin radial walls and thickened tangential external and internal walls. At this level, there are many anisocytic stomata and unicellular long tector hairs, with sharp peak. In cross-section, the median rib has gutter aspect prominent on abaxial face. On the outside,

the epidermis consists of small cells, anterior-posterior slightly flattened, with the external wall covered with a thick cuticle. To the adaxial pole, exhibits rare unicellular sharp tector hairs. Under the epidermis, at both adaxial and abaxial poles, less angular collenchyma is found. At the central area, a single libero-ligneous conducting fascicle surrounded by a assimilatory fascicular sheath is located into a leaf's parenchyma mass. Xylem vessels are seriate disposed into the libero-ligneous fascicle. At this level, the medullary rays are uniseriate, cellulose. The leaf's lamina has bifacial dorsiventral, hypostomatic structure (Figures 6 and 7).




Figure 6. Cross-section through *H. hirsutum* leaf's limb: (a) upper epidermis; (b) cuticle; (c) angular collenchyma; (d) leaf's parenchyma; (e) xylem tissue; (f) phloem tissue; (g) medullary ray (Congo red–chrysoidine staining, $\times 400$).




Figure 7. Cross-section through *H. hirsutum* leaf's limb: (a) upper epidermis; (b) palisade parenchyma; (c) lacunose parenchyma; (d) libero-ligneous conducting fascicle; (e) lower epidermis; (f) tector hair (Congo red–chrysoidine staining, $\times 400$).

TLC analysis

The experimental data on the preliminary TLC analysis of polyphenols from *Hyperici hirsuti herba* are highlighted in Figures 8–10. One chlorogenic acid derivative (R_f 0.13) was identified from the 12 separated chromatographic bands, in an amount of 281.8 mg per 100 g of dried vegetal product.




Figure 8. TLC chromatogram of polyphenols from *Hyperici hirsuti* herba methanolic extract (UV 254 nm, without derivatization). From left to right: first five applications – sample (1–5 µL); subsequent four applications – standards (2 µL); last five applications – sample (6–10 µL).




Figure 9. Densitogram of polyphenols (UV 254 nm) separated from *Hyperici hirsuti* herba methanolic extract.




Figure 10. Chlorogenic acid derivative in situ UV spectra of standard and compound separated from the analyzed sample.

CONCLUSIONS

Histo-anatomical investigation of the root, aboveground stem and leaf of *Hypericum hirsutum* species and preliminary TLC analysis of polyphenols from *Hyperici hirsuti herba* were realized. The root has circular shape and secondary structure due to the presence of two meristematic zones (phellogen and libero-ligneous cambium). The aboveground stem has circular shape and secondary structure generated by the libero-ligneous cambium. Leaf's limb has bifacial dorsiventral, hypostomatic structure. Using thin-layer chromatographic technique, a single chlorogenic acid derivative was isolated and quantified from the 12 separated chromatographic bands.

BIBLIOGRAPHY

1. **Altemimi, A., Watson, D.G., Kinsel, M., Lightfoot, D.A.**, 2015 – *Simultaneous extraction, optimization, and analysis of flavonoids and polyphenols from peach and pumpkin extracts using a TLC-densitometric method*, Chem. Cent. J. 9:39.
2. **Andrei, M., Paraschivoiu, R.M.**, 2003 – *Microtehnica botanică*, Ed. Niculescu, București, 2003, 222 pag.
3. **Barnes, J., Anderson, L.A., Phillipson, J.D.**, 2001 – *St John's wort (Hypericum perforatum L.): a review of its chemistry, pharmacology and clinical properties*, J. Pharm. Pharmacol. 53(5):583–600.
4. **Boji, M., Simon Haas, V., Sari, D., Maleš, Z.**, 2013 – *Determination of flavonoids, phenolic acids, and xanthines in mate tea (Ilex paraguariensis St.-Hil.)*, J. Anal. Methods Chem. 2013:658596.
5. **Cecchini, C., Cresci, A., Coman, M.M., Ricciutelli, M., Sagratini, G., Vittori, S., Lucarini, D., Maggi, F.**, 2007 – *Antimicrobial activity of seven Hypericum entities from Central Italy*, Planta Med. 73(6):564–566.
6. **Ciocârlan V.**, 2000 – *Flora ilustrată a României. Pteridophyta et Spermatophyta*, ediția a 2-a revizuită și adăugită, Ed. Ceres, București, 1138 pag.
7. **Cirak, C., Radusiene, J., Jakstas, V., Ivanauskas, L., Seyis, F., Yayla, F.**, 2016 – *Secondary metabolites of seven Hypericum species growing in Turkey*, Pharm. Biol. 54(10):2244–2253.
8. **Cirak, C., Radusiene, J., Jakstas, V., Ivanauskas, L., Yayla, F., Seyis, F., Camas, N.**, 2016 – *Secondary metabolites of Hypericum species from the Drosanthe and Olympia sections*, S. Afr. J. Bot. 104:82–90.
9. **Dagnino, A.P., Barros, F.M., Ccana-Ccapatinta, G.V., Prophiro, J.S., Poser, G.L., Romão, P.R.**, 2015 – *Leishmanicidal activity of lipophilic extracts of some Hypericum species*, Phytomedicine, 22(1):71–76.
10. **Gîtea, D., Ișco, M., Tîmba, M., Pașca B.**, 2011 – *Secretory structures at species of Hypericum genera from Bihor County, Romania. Note I. Vegetative organs*, Farmacia 59(3):424–431.
11. **Gîrd, C.E., Nencu, I., Costea, T., Dușu, L.E., Popescu, M.L., Ciupitu, N.**, 2014 – *Quantitative analysis of phenolic compounds from Salvia officinalis L. leaves*, Farmacia 62(4):649–657.
12. **Hernandez, M.F., Falé, P.L.V., Araújo, M.E.M., Serralheiro, M.L.M.**, 2010 – *Acetylcholinesterase inhibition and antioxidant activity of the water extracts of several Hypericum species*, Food Chem. 120(4):1076–1082.
13. **Maggi, F., Cecchini, C., Cresci, A., Coman, M.M., Tirillini, B., Sagratini G, Papa F, Vittori S.**, 2010 – *Chemical composition and antimicrobial activity of the essential oils from several Hypericum taxa (Guttiferae) growing in Central Italy (Appennino Umbro-Marchigiano)*, Chem. Biodivers. 7(2):447–466.
14. **Niculescu, Mariana** - *Morfologia și anatomia plantelor*, VOL. I, Ed. Sitech, Craiova, 2009, 201 p., ISBN 978-606-8021-87-4

15. **Niculescu, M., F g ra M.**, 2015 - *The grassland natura 2000 habitats found in Lespezi quarry and the surroundings, Dambovita county*, 15th International Multidisciplinary Scientific GeoConference SGEM 2015, www.sgem.org, SGEM2015 Conference Proceedings, ISBN 978-619-7105-39-1 / ISSN 1314-2704, June 18-24, 2015, Book5 Vol. 1, 877-882 pp, DOI: 10.5593/SGEM2015/B51/S20.116, <http://www.sgem.org/sgemlib/spip.php?article6369&lang=en>
16. **Niculescu, L., Niculescu, M., Soare, R., Bonea, D.**, 2015 - *Plant communities characteristic to the montane and subalpine springs and streams in the Parang Mountains (Southern Carpathians), Romania* - JEPE, Vol. 16, No. 4 (2015), <https://docs.google.com/a/jepe> journal.info/viewer?a=v&pid=sites&srcid=amVwZS1qb3VybmFsLmluZm98amVwZS1qb3VybmFsGd4OjgwMzlwZGUwMDQ2N2E2NQ
17. **Nogueira, T., Medeiros, M.A., Marcelo-Curto, M.J., García-Pérez, B.E., Luna-Herrera, J., Costa, M.C.**, 2013 – *Profile of antimicrobial potential of fifteen Hypericum species from Portugal*, Ind. Crops Prod. 47:126–131.
18. **Perrone, R., De Rosa, P., De Castro, O., Colombo, P.**, 2013 – *Leaf and stem anatomy in eight Hypericum species (Clusiaceae)*, Acta Bot. Croat. 72(2):269–286.
19. **Sarris, J., Fava, M., Schweitzer, I., Mischoulon, D.**, 2012 – *St John's wort (Hypericum perforatum) versus sertraline and placebo in major depressive disorder: continuation data from a 26-week RCT*, Pharmacopsychiatry 45(7):275–278.
20. **Savikin, K., Dobri , S., Tadi , V., Zduni , G.**, 2007 – *Anti-inflammatory activity of ethanol extracts of Hypericum perforatum L., H. barbatum Jacq., H. hirsutum L., H. richeri Vill. and H. androsaemum L. in rats*, Phytother. Res. 21(2):176–180.
21. **Smelcerovic, A., Spiteller, M.**, 2006 – *Phytochemical analysis of nine Hypericum L. species from Serbia and the F.Y.R. Macedonia*, Pharmazie 61(3):251–252.
22. **Stojanovi , G., Đor evi , A., Šmelcerovi , A.**, 2013 – *Do other Hypericum species have medical potential as St. John's wort (Hypericum perforatum)?* Curr. Med. Chem. 20(18):2273–2295.
23. **Toma, C., Rugin , R.**, 1998 – *Anatomia plantelor medicinale. Atlas*, Ed. Academiei Române, Bucure ti, 1998, 320 pag.
24. **Umek, A., Kreft, S., Kartnig, T., Heydel, B.**, 1999 – *Quantitative phytochemical analyses of six Hypericum species growing in Slovenia*, Planta Med. 65(4):388–390.