

THE INFLUENCE OF DIFFERENT TREATMENTS OF SOIL POLLUTED WITH CRUDE OIL ON PLANT BIOMASS

MARIANA MARINESCU, ANCA LACATUSU, EUGENIA GAMENT, M. MARINESCU,

National Research Institute for Soil Science, Agrochemistry and Environmental Protection, Bucharest, Romania, e-mail: maramarinescu2003@yahoo.com

Keywords: plant biomass, treatments, soil polluted, crude oil.

ABSTRACT

The aim of this research was to improve the bioremediation technology for soils that are polluted with crude oil through biostimulation, bioaugmentation and phytoremediation. The soil artificial polluted with crude oil have been treated with a natural biodegradable product and bacterial inoculum. The plant used in the greenhouse experiment is maize. In this paper is presented the influence of different treatments on plant biomass. It will be shown the influence of crude oil on plant growth (plant number/pot, plant length, leaves number and plant biomass) in a soil treated with a natural biodegradable product and inoculated with bacterial inoculum. According with the results obtained, this method could be used in field for rehabilitation and reuse in agriculture of polluted soils with crude oil.

INTRODUCTION

The development of human civilization throughout history has led to growing disruption of the natural balance and the occurrence of different types of pollution. The world depends on oil, and the use of oil as fuel has led to intensive economic development worldwide. The great need for this energy source has led to the gradual exhaustion of natural oil reserves. However, mankind will witness the results of oil consumption for centuries after its cessation. Environmental pollution with petroleum and petrochemical products has been recognized as a significant and serious problem (Alexander, 1995, 2000). Most components of oil are toxic to humans and wildlife in general, as it is easy to incorporate into the food chain. This fact has increased scientific interest in examining the distribution, fate and behaviour of oil and its derivatives in the environment (Alexander, 1995, 2000; Semple et al., 2001, 2003; Stroud et al., 2009). Oil spills in the environment cause long-term damage to aquatic and soil ecosystems, human health and natural resources.

One of the most important classes of organic pollutants in the environment is crude oil constituents of petrochemicals. Accidental oil pollution has become nowadays a common phenomenon that can cause environmental and social disasters (Burger, 1993; Burns, 1993; Pepper, 1996). Potential sources of direct pollution of soil and subsoil can be covered by tanks, separators old from wastewater treatment plants, settling basins, slurries and waste pits of tar, ramp CF loading and unloading, underground pipelines, sewerage networks, etc.

Plants are designed to increase the activity of microorganisms in rhizosphere by optimizing the parameters of the environment, such as moisture, soil reaction. Further, roots growth involves the penetration of needed oxygen for contaminant/pollutant oxidation process (Joner and Leyval, 2003; Lin and Mendelsohn, 1998). Although hydrocarbons biodegradation around rhizosphere are known, still have not been clarified the mechanisms that influence the microorganisms growth and activity (Hutchinson et al., 2001; Wiltse et al., 1998).

MATERIAL AND METHOD

The main objective of this research is testing the natural hydrocarbon absorbent named ECOSOL. To achieve data concerning the bioremediation of polluted soil with petroleum hydrocarbons was achieved a greenhouse experiment. The soil used for this experiment was a cambic chernozems. The plant used in the experiment was maize.

The experimental variants are:

- ✓ V₁, control (unpolluted soil);
- ✓ V₂, polluted soil with 5% crude oil;
- ✓ V₃, polluted soil with 10% crude oil;
- ✓ V₄, polluted soil with 5% crude oil + 1 kg ECOSOL/m² polluted soil;
- ✓ V₅, polluted soil with 5% crude oil + 1 kg ECOSOL/m² polluted soil + bacterial inoculum;
- ✓ V₆, polluted soil with 5% crude oil + 2 kg ECOSOL/m² polluted soil;
- ✓ V₇, polluted soil with 5% crude oil + 2 kg ECOSOL/m² polluted soil + bacterial inoculum;
- ✓ V₈, polluted soil with 10% crude oil + 2 kg ECOSOL/m² polluted soil;
- ✓ V₉, polluted soil with 10% crude oil + 2 kg ECOSOL/m² polluted soil + bacterial inoculum;
- ✓ V₁₀, polluted soil with 10% crude oil + 4 kg ECOSOL/m² polluted soil;
- ✓ V₁₁, polluted soil with 10% crude oil + 4 kg ECOSOL/m² polluted soil + bacterial inoculum.

RESULTS AND DISCUSSIONS

Soil artificial polluted with 5 and 10% crude oil by volume, was treated with Ecosol in variant uninoculated and inoculated with selected bacteria. At the end of the first experimental year of testing, the polluted soil material was allowed to stand over the winter, when were not applied water or mixing and aerating procedures. Storage temperature was close to the exterior environment of Green House.

The seeding was carried out in April of the second experimental year, at a depth of 8-10 cm of 5 grains per pot. The first plants have emerged Control and in late in the experimental pots polluted with 5% crude oil. In experimental pots with soil polluted 10% crude oil, plants do not emerged at all excessive concentration of pollutant exerting a very severe phytotoxic effect, preventing total germination.

The soil was kept in an optimal umidity state (approximately corresponding to the water capacity in the field). The beneficial effects of the applied treatments were emphasized in maize plants growth, which grew only in experimental variants polluted with 5% crude oil, best developed in experimental variant V7 with 100 g Ecosol and inoculated with selected bacteria.

In the third experimental year, the soil was homogenized before experiment reorganization. Sowing was performed in April, at a depth of 8-10 cm with a total of 5 beans in each pot. As expected, the first seedlings have emerged in control. Unlike in the previous experimental year, although the maize plants have emerged with delay in experimental pots polluted with crude oil at both concentrations.

The benefit effects of treatments applied were found in maize plants growth in these second experimental year, which could grow only in the variants polluted with 5% crude oil, the better option and inoculated with 0.5% (100g/pot) Ecosol with selected bacteria.

Biometric determinations carried out on the maize plants are presented in tables. The evolution of plant number/pot, plant length, leaves number and plant biomass in soil polluted with 5% and 10% crude oil in different experimental variants – analysis of variance are presented in table 1-4.

Table 1.

The evolution of plant number/pot in soil polluted with crude oil in different experimental variants – analysis of variance

Experimental variant		Plants number/pot	
		2nd year	3rd year
unpolluted soil	V1	5	5
polluted soil with 5% crude oil	V2	4	5
polluted soil with 5% crude oil + 1 kg ECOSOL/m ² polluted soil	V4	5	5
polluted soil with 5% crude oil + 1 kg ECOSOL/m ² polluted soil + bacterial inoculum	V5	4	5
polluted soil with 5% crude oil + 2 kg ECOSOL/m ² polluted soil	V6	5	5
polluted soil with 5% crude oil + 2 kg ECOSOL/m ² polluted soil + bacterial inoculum	V7	5	5
polluted soil with 10% crude oil	V3	-	5
polluted soil with 10% crude oil + 2 kg ECOSOL/m ² polluted soil	V8	-	5
polluted soil with 10% crude oil + 2 kg ECOSOL/m ² polluted soil + bacterial inoculum	V9	-	5
polluted soil with 10% crude oil + 4 kg ECOSOL/m ² polluted soil	V10	-	5
polluted soil with 10% crude oil + 4 kg ECOSOL/m ² polluted soil + bacterial inoculum	V11	-	5

Table 2.

The evolution of plant length in soil polluted with crude oil in different experimental variants – analysis of variance

Experimental variant		Length (cm)	
		2nd year	3rd year
unpolluted soil	V1	139	122
polluted soil with 5% crude oil	V2	38	66
polluted soil with 5% crude oil + 1 kg ECOSOL/m ² polluted soil	V4	43	80
polluted soil with 5% crude oil + 1 kg ECOSOL/m ² polluted soil + bacterial inoculum	V5	55	83
polluted soil with 5% crude oil + 2 kg ECOSOL/m ² polluted soil	V6	59	89
polluted soil with 5% crude oil + 2 kg ECOSOL/m ² polluted soil + bacterial inoculum	V7	62	114
polluted soil with 10% crude oil	V3	-	56
polluted soil with 10% crude oil + 2 kg ECOSOL/m ² polluted soil	V8	-	83
polluted soil with 10% crude oil + 2 kg ECOSOL/m ² polluted soil + bacterial inoculum	V9	-	87

polluted soil with 10% crude oil + 4 kg ECOSOL/m ² polluted soil	V10	-	88
polluted soil with 10% crude oil + 4 kg ECOSOL/m ² polluted soil + bacterial inoculum	V11	-	112

Table 3.

The evolution of plant leaves number in soil polluted with crude oil in different experimental variants – analysis of variance

Experimental variant		Leaves number	
		2nd year	3rd year
unpolluted soil	V1	12	14
polluted soil with 5% crude oil	V2	7	8
polluted soil with 5% crude oil + 1 kg ECOSOL/m ² polluted soil	V4	8	8
polluted soil with 5% crude oil + 1 kg ECOSOL/m ² polluted soil + bacterial inoculum	V5	8	8
polluted soil with 5% crude oil + 2 kg ECOSOL/m ² polluted soil	V6	8	8
polluted soil with 5% crude oil + 2 kg ECOSOL/m ² polluted soil + bacterial inoculum	V7	8	9
polluted soil with 10% crude oil	V3	-	8
polluted soil with 10% crude oil + 2 kg ECOSOL/m ² polluted soil	V8	-	8
polluted soil with 10% crude oil + 2 kg ECOSOL/m ² polluted soil + bacterial inoculum	V9	-	8
polluted soil with 10% crude oil + 4 kg ECOSOL/m ² polluted soil	V10	-	9
polluted soil with 10% crude oil + 4 kg ECOSOL/m ² polluted soil + bacterial inoculum	V11	-	9

Table 4.

The evolution of plant biomass in soil polluted with crude oil in different experimental variants – analysis of variance

Experimental variant		Biomass (g)	
		2nd year	3rd year
unpolluted soil	V1	106	177
polluted soil with 5% crude oil	V2	5	46
polluted soil with 5% crude oil + 1 kg ECOSOL/m ² polluted soil	V4	6	62
polluted soil with 5% crude oil + 1 kg ECOSOL/m ² polluted soil + bacterial inoculum	V5	7	64
polluted soil with 5% crude oil + 2 kg ECOSOL/m ² polluted soil	V6	8	69
polluted soil with 5% crude oil + 2 kg ECOSOL/m ² polluted soil + bacterial inoculum	V7	9	109
polluted soil with 10% crude oil	V3	-	38

polluted soil with 10% crude oil + 2 kg ECOSOL/m ² polluted soil	V8	-	68
polluted soil with 10% crude oil + 2 kg ECOSOL/m ² polluted soil + bacterial inoculum	V9	-	72
polluted soil with 10% crude oil + 4 kg ECOSOL/m ² polluted soil	V10	-	77
polluted soil with 10% crude oil + 4 kg ECOSOL/m ² polluted soil + bacterial inoculum	V11	-	109

Comparison between experimental variants showed a better development of maize plants in variants with soil polluted with 5% crude oil and inoculated, the same observing also in variants of soil polluted with 10% oil and inoculated. Inoculation of soil polluted with bacteria appears to confer enhanced plant vigor of maize in combination with soil conditioning with 0.5 % and 1 % Ecosol.

Doubling the dose of Ecosol (0.5% and 1%) added to the existing maize plants could flourish in conditions of soil pollution with 10% crude oil. Obviously, the more developed plants and flowers of maize variants in which the two technological, soil conditioning Ecosol and inoculation with selected bacteria, acted synergistically.

In the experimental variants of soil polluted with 5% crude oil, biomass differences are significantly lower compared to plant biomass determined in unpolluted soil. Although uninsured statistically, biomass values gradually increase in experimental variants, from the natural attenuation to variant with 0.5% Ecosol inoculated with selected bacteria. In the third experimental year and the second year of experiment with plant, differences between experimental variants polluted end significance, so that all variants undergo bioremediation process developed higher biomass values compared to that obtained with natural attenuation in the embodiment and the variant in which the soil was conditioned with 0.5 % Ecosol (100 g/pot) and inoculated with selected bacteria generated an increase significant biomass bioremediation variants compared to the rest. The experimental variants polluted with 10% oil in the second experimental year, the first of experimentation with plant biomass was phytotoxic effect of excessive oil pollution completely preventing the plants to germinate. The following year, biomass plants in variants bioremediation was significantly higher than that of plants from variant natural attenuation and the optimal variant in which the soil was conditioned with 1% Ecosol and inoculated with selected bacteria, plant biomass was significantly higher compared to other experimental variants bioremediation. The data were correlated clear dynamics of soil pollutant disappearance.

The beneficial effects of the treatments were found in development of maize plants grown in the second experimental year, who have developed version only contaminated with 5% of oil, the more the solution and inoculated with 100 g Ecosol with selected bacteria.

Biomass plants in bioremediation variants was significantly higher than that of plants in natural attenuation variants, and the experimental variants considered optimal in the soil was conditioned with higher Ecosol doses and inoculated with selected bacteria, plant biomass was significantly higher compared with other experimental variants bioremediation. The data were correlated clearly with the dynamics of soil pollutant disappearance.

CONCLUSIONS

The cleaning up of petroleum hydrocarbons in the soil environment is a real world problem. Better understanding of the mechanisms and factors which affect biodegradation is of great ecological significance, since the choice of bioremediation strategy depends on it. Microbial degradation processes aid the elimination of spilled oil from the environment,

together with various physical and chemical methods. This is possible because microorganisms have enzyme systems to degrade and utilize different hydrocarbons as a source of carbon and energy. Even if the optimal conditions for microbial degradation are provided, the extent of hydrocarbon removal is strongly affected by its bioavailability and stages of weathering.

The beneficial effects of the treatments were found in development of maize plants grown in the second experimental year, who have developed in experimental variants only contaminated with 5% of oil, the more the solution and inoculated with 100 g Ecosol with selected bacteria. In the first year of experimenting with plants in pots contaminated with excessive concentration of 10% oil pollutant exercised severely phytotoxic effect, preventing total germination. In the second year experimental plant, maize plants have sprung up in crude oil contaminated pots at both concentrations of force and the enormous gap size between plants affected by oil pollution and developed a clean soil very visible. Inoculation of soil polluted with bacteria selected in combination with soil conditioning with 0.5 % and 1 % Ecosol had a beneficial effect on plant vigor of maize. Obviously, the more developed plants and flowers of maize variants in which the two technological measures act synergistically.

BIBLIOGRAPHY

1. Alexander, M., 1995 -*How toxic are toxic chemicals in soil? Environmental Science and Technology*, Vol. 29, No. 11, pp. 2713–2717, ISSN 0013-936X.
2. Alexander, M., 2000 -*Aging, bioavailability, and overestimation of risk from environmental pollutants*, *Environmental Science and Technology*, Vol. 34, No. 20, pp. 4259–4265, ISSN 0013-936X.
3. Burger, A.E., 1993 - *Estimating the mortality of seabirds following oil spills – effects of spill volume*, *Marine Pollution Bulletin* 26, pag. 239–248.
4. Burns, K.A., Garrity, S.D., Levings, S.C., 1996 - *How many years until mangrove ecosystems recover from catastrophic oil-spills*, *Marine Pollution Bulletin* 26, pag. 239–248, 1993.1. Cunningham, S.D., Anderson, T.A., Schwab, A.P., Hsu, F.C., *Phytoremediation of soils contaminated with organic pollutants*, *Advances in Agronomy* 56, pag. 55–113.
5. Hutchinson, S.L., Schwab, A.P., Banks, M.K., 2001 - *Bioremediation and biodegradation: phytoremediation of aged petroleum sludge: effect of irrigation techniques and scheduling*, *Journal of Environmental Quality* 30, pag. 1516–1522.
6. Joner, E.J., Leyval, C., 2003 - *Phytoremediation of organic pollutants using mycorrhizal plants: a new aspect of rhizosphere interactions*, *Agronomie* 23, pag. 495–502.
7. Lin, Q., Mendelssohn, I.A., 1998 - *The combined effects of phytoremediation and biostimulation in enhancing habitat restoration and oil degradation of petroleum contaminated wetlands*, *Ecological Engineering* 10, pag. 263–274,.
8. Pepper, Ian L., Gerba, Charles P., Brusseau, Mark L, 1996 - *Pollution Science*, Academic Press.
9. Semple, K.T.; Morris, A.W.J. & Paton, G.I., 2003 -*Bioavailability of hydrophobic organic contaminants in soils: fundamental concepts and techniques for analysis*. *European Journal of Soil Science*, Vol. 54, pp. 809-818, ISSN 0022-4588.
10. Semple, K.T.; Reid, B.J. & Fermor, T.R., 2001 -*Impact of composting strategies on the treatment of soils contaminated with organic pollutants*. *Environmental Pollution*, Vol. 112, No. 2, pp. 269-283, ISSN 0269-7491.
11. Stroud, J.L.; Paton, G.I. & Semple, K.T., 2009 -*Predicting the biodegradation of target hydrocarbons in the presence of mixed contaminants in soil*. *Chemosphere*, Vol. 74, No. 4, pp. 563–567, ISSN 0045-6535
12. Wiltse, C.C., Rooney, W.L., Chen, Z., Schwab, A.P., Banks, M.K., 1998 - *Greenhouse evaluation of agronomic and crude oil-phytoremediation potential among alfalfa genotypes*, *J. Environ. Qual.* 27, pag. 169–173.