


SURFICIAL GEOLOGY OF THE PORTLAND 1° X 2° QUADRANGLE, MAINE


Compiled By:
Woodrow B. Thompson, Maine Geological Survey

Area Compilers and Contributors:
Geoffrey W. Smith, University of Ohio, Athens
Woodrow B. Thompson, Maine Geological Survey


Maine Geological Survey
DEPARTMENT OF CONSERVATION
Augusta, Maine 04333
Walter A. Anderson, State Geologist
1987
OPEN-FILE NO. 87-6



COMPILED RESPONSIBILITY



Funding for the preparation and publication of this map was provided by a grant from the U.S. Department of Energy (grant no. DE-FG05-81NE4664) and the Maine Geological Survey.



EXPLANATION

Geologic Unit	Materials	Topography	Origin
a Stream alluvium (includes Holocene flood plain, stream terrace, and alluvial fan deposits)	Sand, gravel, and silt.	Flat to gently sloping on flood plains and stream terraces; gently to moderately sloping on alluvial fans.	Deposited on flood plains and stream beds by postglacial streams.
s Swamp, marsh, and bog deposits (includes both fresh-water and salt-water marshes)	Peat, muck, clay, silt, and sand.	Flat.	Formed by accumulation of sediments and organic material in depressions and other poorly drained areas.
b Beach deposits	Sand and gravel.	Gently to moderately sloping, with low ridges and mounds.	Includes beach sediments formed by wave and current action, and sand dunes derived from these deposits.
eb Emerged beach deposits	Sand and gravel.	Low ridges or sloping surfaces. May be associated with wave-cut benches on hillsides.	Formed by wave erosion of till or other materials during the late-glacial marine submergence of parts of southern Maine.
e Eolian deposits	Sand.	Dune ridges and mounds, or blanket deposit that conforms to surface of underlying unit.	Windblown sand. Derived from wind erosion of glacial sediments and deposited in late-glacial to postglacial time.
L Lake-bottom deposits	Silt, clay, and sand. Commonly well stratified, and may be rhythmically bedded.	Flat to gently sloping except where dissected by modern streams.	Composed of sediments that washed out of late Wisconsinan glacial ice and accumulated on the floors of glacial lakes. Map unit may also include a few non-glacial lake deposits.
m Glaciomarine deposits (fine-grained facies)	Silt, clay, sand, and minor amounts of gravel. Commonly a clayey silt (the Presumpscot Formation). Sand is dominant in some places, but may be underlain by finer grained sediments. Locally fossiliferous. May unit includes small areas of till and other units that are not completely covered by marine sediments.	Flat to gently sloping except where dissected by modern streams. Commonly has a branching network of steep-walled stream gullies.	Composed of glacial sediments that accumulated on the ocean floor. Formed during the late-glacial marine submergence of lowland areas in southern Maine.
ms Glaciomarine deposits (coarse-grained facies)	Sand, gravel, and minor amounts of silt.	Flat to moderately sloping. Steeper on ice-contact slopes and delta fronts. May be kettled where deposited over stagnant ice blocks.	Deposited where glacial meltwater streams and currents entered the sea. Includes glaciomarine deltas, subaqueous fans and bars (subaqueous outwash), and outwash that prograded into shallow marine waters and locally covered earlier glaciomarine silt and clay deposits.
go Glacial outwash deposits	Sand and gravel.	Flat to gently sloping. Steeper on ice-contact slopes and delta fronts. May be kettled where deposited over stagnant ice blocks.	Deposited by meltwater streams in front of the retreating late Wisconsinan ice margin. Includes non-marine outwash plains, deltas, and fans.
g Ice-contact glaciofluvial deposits (exclusive of eskers)	Sand, gravel, and silt.	Flat-topped kame terraces and deltas which are locally kettled and bounded by steep sides, or hummocky terrain with numerous kames and kettles.	Deposited by meltwater streams adjacent to stagnant glacial ice.

Geologic Unit	Materials	Topography	Origin
gr Eskers	Gravel and sand. May include minor amounts of till. Portions of many eskers below the marine limit are partly or entirely buried by glaciomarine deposits.	Individual or multiple ridges. Complex eskers may have anastomosing patterns and be gradational with other types of ice-contact deposits.	Chiefly deposited by meltwater streams flowing in tunnels within or beneath the late Wisconsinan ice sheet. May unit also includes small undifferentiated areas of units "g" and "go".
sm Stagnation moraine	Mostly till, but also includes variable percentages of undifferentiated sand and gravel.	Undulating topography with local hummocks and ridges.	Deposited during the dissipation of stagnant glacial ice.
om End moraines	Till or sand and gravel. May be very boundary. Commonly interbedded with or overlain by glaciomarine sediments in areas that experienced late-glacial marine submergence. Only the largest and most dense clusters of smaller moraines are shown here as a separate unit (em). Elsewhere, short lines mark the crests of moraine ridges, which are locally so numerous that only selected individuals are represented.	Ridges. Commonly arcuate, discontinuous, and in groups. May be multi-crested and hummocky. Side ranges: L-40 ft high, 5-200 ft wide, and 30 ft to over 10 mi long.	Deposited in the marginal zone of the late Wisconsinan ice sheet, by glacial ice and/or meltwater flowing out of the ice.
rm Ribbed moraine	Till is the principal constituent, but stratified sediments are present in some of the deposits.	Numerous hummocks and short sub-parallel ridges which typically occur in lake basins and other lowland areas.	Origin uncertain. Deposited either at the margin of or beneath the late Wisconsinan ice sheet.
t Till	Heterogeneous mixture of sand, silt, clay and stones. May include many boulders. Generally massive, but in many places contains beds and lenses of variably washed and stratified sediments.	Generally a blanket deposit that conforms to the underlying bedrock topography. Also forms drumlins and other glacially streamlined hills.	Deposited directly by glacial ice.
Thin drift	Area of many bedrock outcrops and/or thin surficial deposits (generally less than 3 m thick). The type of surficial material is known or inferred.	Topography of these areas reflects the configuration of the bedrock surface, and ranges from smooth undulating hills to knobby terrain and high mountains.	Commonly the result of non-deposition of glacial sediments, but the surficial materials in some coastal areas have been locally removed by marine erosion in late-glacial time.
tdu Thin drift, undifferentiated	Area of many bedrock outcrops and/or near-surface bedrock where the surficial materials have not been mapped.	Same as other thin-drift areas.	Same as other thin-drift areas.
rk Bedrock	Area of extensive bedrock outcrop, or thin bedrock lens. May be a thin cover of soil and vegetation. Surficial geology is essentially absent. Particularly common on the ridge crests and steeper slopes of mountainous areas.	Hilly to mountainous terrain.	Same as the thin-drift areas.


GEOLOGIC SYMBOLS

	Contact	Boundary between adjacent map units.
	Moraine ridge	Lines mark the crests of individual moraines. Symbol also is used in conjunction with unit ms to show orientation of drift ridges of uncertain origin.
	Glacial striation locality	Includes striations, grooves, drag-and-tails, and other types of ice-flow indicators on bedrock outcrops. Dot indicates point of observation. Arrowhead is omitted where ice-flow direction is uncertain. Flag indicates older trends.
	Glacially streamlined landform	Symbol shows long-axis orientation of drumlins, fluted till ridges, roches moutonnées, and other hills that have been elongated parallel to the flow of glacial ice.
	Cirque	Steep-walled, semicircular bedrock basin formed by glacial erosion in high mountainous areas.
	Meltwater channel	Channel eroded by glacial meltwater streams. Arrow indicates known or inferred direction of stream flow.
	▲ 288 Glaciomarine delta	Number indicates surveyed altitude (in feet) of contact between topset and foreset beds, or of meltwater channel on delta surface, which approximately marks position of sea level in late-glacial time.
	▲ 345 Glaciolacustrine delta	Number indicates approximate altitude (in feet) of former glacial-lake surface.
	▲ 699 Delta of uncertain origin	Delta formed near limit of late-glacial marine submergence. Number indicates approximate altitude (in feet) of contact between topset and foreset beds.

SOURCES OF GEOLOGIC INFORMATION

- Blum, A. L., 1960, Late Pleistocene changes of sea level in southwestern Maine. *Maine Geol. Surv.*, 43 p.
- Borne, H. W., Jr., 1973, Late Wisconsinan fluctuations of the Laurentide ice sheet in southern and eastern New England, in Black, R. F., Goldthwait, R. P., and Wilman, H. B., (eds.), *The Wisconsinan Stage*. *Geol. Soc. Amer. Memoir* 150, p. 37-45.
- Calderell, D. W., 1965, Eolian features in Freeport and Wayne, Maine. In *History, A. M., II, (ed.), New England Intercollegiate Geological Conference Guidebook for Field Trips in Southern Maine*, p. 59-102.
- Calderell, D. W., Hanson, L. S., and Holland, W. R., 1979, Drumlins in southwestern Maine containing abundant stratified drift. *Geol. Soc. Amer. Bull.*, 90, p. 250-253.
- Husey, A. M., II, 1971, *Geologic map of the Portland Quadrangle, Maine*. *Maine Geol. Surv.*, Map 59-1, 19 p.
- Byland, F., Thompson, W. B., and Stuckert, R., Jr., 1978, Late Wisconsinan wood and other tree remains in the Presumpscot Formation, Portland, Maine. *Maritime Sediments*, v. 14, p. 103-120.
- Keller, S. B., unpublished reconnaissance data.
- Freestott, G. C., Jr., 1981, Unpublished availability and surficial geology of the Royal, upper Presumpscot, and upper Saco River basins, Maine. *U.S. Geol. Surv. Water Resources Invest.*, 79-1287.
- Freestott, G. C., Jr., Smith, G. W., and Thompson, W. B., 1976, Reconnaissance surficial geology of the Cumberland Center (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-10.
- Freestott, G. C., Jr., Smith, G. W., and Thompson, W. B., 1977, Reconnaissance surficial geology of the Old Orchard Beach (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-25.
- Freestott, G. C., Jr., Smith, G. W., and Thompson, W. B., 1977, Reconnaissance surficial geology of the South Portland (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-40.
- Freestott, G. C., Jr., and Thompson, W. B., 1976, Reconnaissance surficial geology of the North Windham (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 76-31.
- Freestott, G. C., Jr., and Thompson, W. B., 1976, Reconnaissance surficial geology of the Old Orchard Beach (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 76-25.
- Freestott, G. C., Jr., Smith, G. W., and Thompson, W. B., 1977, Reconnaissance surficial geology of the South Portland (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-40.
- Freestott, G. C., Jr., and Thompson, W. B., 1976, Reconnaissance surficial geology of the Farmington (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-7.
- Smith, G. W., 1977, Reconnaissance surficial geology of the Barwick (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-9.
- Smith, G. W., 1977, Reconnaissance surficial geology of the Hallowell (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-10.
- Smith, G. W., 1977, Reconnaissance surficial geology of the Dover (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-12.
- Smith, G. W., 1977, Reconnaissance surficial geology of the Kennebunk (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-13.
- Smith, G. W., 1977, Reconnaissance surficial geology of the Kears Falls (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-14.
- Smith, G. W., 1977, Reconnaissance surficial geology of the Newfield (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-15.
- Smith, G. W., 1977, Reconnaissance surficial geology of the Portland (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-16.
- Smith, G. W., 1977, Reconnaissance surficial geology of the York (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-18.
- Smith, G. W., 1977, Reconnaissance surficial geology of the Kennebunk glacial advance: a reappraisal! *Geol. Soc. Amer. Bull.*, v. 88, p. 99-104.
- Smith, G. W., 1980, Chronology of late Wisconsinan deglaciation of coastal Maine. In Borne, H. W., Jr., Labille, P., and Thompson, W. B., (eds.), *Late Pleistocene History of Northeastern New England and Adjacent Quebec*. *Geol. Soc. Amer., Special Paper* 197, p. 29-44.
- Smith, G. W., and Thompson, W. B., 1976, Reconnaissance surficial geology of the Gorham (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 76-42.
- Smith, G. W., and Thompson, W. B., 1977, Reconnaissance surficial geology of the Buxton (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-19.
- Smith, G. W., and Thompson, W. B., 1980, Reconnaissance surficial geology of the Freeport (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 80-12.
- Stuiver, M., and Borne, H. W., Jr., 1975, Late Quaternary marine invasion in Maine: its chronology and associated crustal movements. *Geol. Soc. Amer. Bull.*, v. 86, p. 99-104.
- Thompson, W. B., 1976, Reconnaissance surficial geology of the Cape Elizabeth (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 76-43.
- Thompson, W. B., 1976, Reconnaissance surficial geology of the Cornish (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 76-44.
- Thompson, W. B., 1976, Reconnaissance surficial geology of the Portland West (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 76-45.
- Thompson, W. B., 1976, Reconnaissance surficial geology of the Freeport Beck (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 76-46.
- Thompson, W. B., and Freestott, G. C., Jr., 1977, Reconnaissance surficial geology of the Portland East (7.5') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-40.
- Thompson, W. B., and Smith, G. W., 1976, Reconnaissance surficial geology of the Gray (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 76-47.
- Thompson, W. B., and Smith, G. W., 1977, Reconnaissance surficial geology of the Sebago Lake (15') quadrangle, Maine. *Maine Geol. Surv.*, Open-File Map 77-42.

INDEX TO SOURCES OF GEOLOGIC INFORMATION



SITES OF SPECIAL INTEREST

● Location of special site

This list includes locations of important stratigraphic sections of Pleistocene deposits in Maine, and places where good examples of certain glacial features can be seen. The sites were selected partly on the basis of accessibility, ease of observation, and relative permanence. Sites where good examples of certain glacial features can be seen, but only a few of the best examples are included here.

Site	Town	Name/Description	Principal References
1	Eliot	Great Hill; pit exposure showing interior of drumlin.	Calderell, Hanson, and Holland (4)
2	Kennebunk	Great Hill; coastal exposure showing two tills separated by radio-carbonated marine sediments.	Smith (24)
3	Kennebunk	Kennebunk landfill; type locality for Kennebunk Resurgence (Bloom); pit exposure showing DeGeer moraine and other glaciomarine deposits indicative of oscillatory ice-margin retreat (Smith).	Bloom (1); Borne (2); Smith (23)
4	Buxton	Group of prominent DeGeer moraines.	Bloom (1)
5	Portland	Portland fossil-wood locality; pit exposure showing spruce logs and other plant remains buried in the Presumpscot Formation.	Byland, Thompson, and Stuckert (6)
6	Freeport	Dune of Maine; modern eolian sand deposit with active dunes.	Calderell (3)

RADIOCARBON-DATED SITES

Explanation of symbols used to designate sites on the map:

- Material in place between late Wisconsinan tills.
- Material that predates or is contemporaneous with the advance of the late Wisconsinan ice sheet.
- ▣ Material that postdates or is contemporaneous with the recession of the late Wisconsinan ice sheet.
- Material that approximately dates the onlay of the sea during the late-glacial marine transgression.
- Material that approximately dates the offlay of the sea.

Site	Name/Town	Date (yr B.P.)	Laboratory No.	Material	Reference
A	Great Hill (Kennebunk)	13,800 ± 100	GL-192	marine shells between tills	Smith (24)
B	Kennebunk	13,200 ± 120	Y-2208	shells in deformed glacio-marine sediments	Stuiver and Borne (28)
C	Little Falls (Gorham)	11,400 ± 90	SI-4747	shells in marine sediments	Smith (24)