### Baptist Health South Florida Scholarly Commons @ Baptist Health South Florida

### All Publications

2015

### Development of an RCU Vertical Care Expedites Low Acuity Patients Through the ED

Griselle Pastor West Kendall Baptist Hospital, griselled@baptisthealth.net

Julie Lamoureux *West Kendall Baptist Hospital*, julieal@baptisthealth.net

Andrew Friedman Baptist Health South Florida, andrewf@baptisthealth.net

Matthew Kadrie Baptist Health South Florida, matthewk@baptisthealth.net

Sebastian Abdelnur *West Kendall Baptist Hospital*, sebastianab@baptisthealth.net

See next page for additional authors

Follow this and additional works at: https://scholarlycommons.baptisthealth.net/se-all-publications Part of the <u>Emergency Medicine Commons</u>, and the <u>Nursing Commons</u>

### Citation

Pastor, Griselle; Lamoureux, Julie; Friedman, Andrew; Kadrie, Matthew; Abdelnur, Sebastian; and Scott, Joseph, "Development of an RCU Vertical Care Expedites Low Acuity Patients Through the ED" (2015). *All Publications*. 3090. https://scholarlycommons.baptisthealth.net/se-all-publications/3090

This Conference Poster -- Open Access is brought to you for free and open access by Scholarly Commons @ Baptist Health South Florida. It has been accepted for inclusion in All Publications by an authorized administrator of Scholarly Commons @ Baptist Health South Florida. For more information, please contact Carrief@baptisthealth.net.

### Authors

Griselle Pastor, Julie Lamoureux, Andrew Friedman, Matthew Kadrie, Sebastian Abdelnur, and Joseph Scott



# West Kendall Baptist Hospital

An academic affiliate of the **FIU** Herbert Wertheim College of Medicine

**BAPTIST HEALTH SOUTH FLORIDA** 



## Background

The number of Emergency Department (ED) visits increased in the United States by 34% between 1995 and 2010. During the same period, the number of EDs decreased by 11%. As a result, EDs experience overcrowding which hampers the delivery of safe and quality care. Factors that influence the degree of overcrowding include patient volume, flow within the ED, boarding, and physical space limitations. Overcrowding in the ED can lead to decreased patient satisfaction, rushed and unpleasant treatment environments, suffering for those who wait, and poor patient outcomes.

Urgent care centers effectively treat lower acuity patients thereby diverting them from EDs. In turn, nearby EDs may experience a higher percentage of patients who require extensive and lengthy evaluations, delaying the evaluation and treatment of lower acuity patients who do present to the ED.

To encourage EDs to address these issues, the Centers for Medicare & Medicaid Services (CMS) implemented throughput benchmarks that must be met and are nationally reported. These include:

- 1. OP18-Median time from ED arrival to ED departure for outpatients (includes observation patients). National benchmarks are top decile: 91 minutes, median: 131 minutes
- 2. OP20-Median time from ED arrival to diagnostic evaluation by a qualified medical personnel for outpatients (includes observation patients). National benchmarks are top decile: 11 minutes, median: 23 minutes

Since opening its doors in 2011, the ED at West Kendall Baptist Hospital (WKBH) has experienced yearly volume increases of 10-15%. With insufficient space to develop a separate "fast-track", we developed a Rapid Care Unit (RCU) in our triage area to address the issues listed above.

### **Project Goals**

The main goals of this project were to:

- 1. Reduce the median length of stay in the ED for low acuity patients.
- 2. Reduce the median time from arrival to diagnostic evaluation by qualified medical personnel for low acuity patients
- 3. Maintain patient satisfaction scores at or above the 90th percentile.
- 4. Utilize existing staffing and space within the ED

# Development of an RCU Vertical Care Expedites Low Acuity Patients Through the ED Griselle Diaz, MBA, MSN, RN, NE-BC, Julie Lamoureux, DMD, MSc, Andrew Friedman, BSSA, MBA, Matthew Kadrie, MBA, Sebastian Abdelnur, MD, Joseph Scott, MD, FACEP

### Methods of Implementation

A triage room is designated for private evaluation, testing, and treatment of Emergency Severity Index (ESI) Level 4 & 5 patients.



Instead on lying in beds, patients sit in recliners in a previously underutilized discharge lounge adjacent to the triage area, pending evaluation and results. Staying "vertical" facilitates the patients' movement through the ED.



This Rapid Care Unit (RCU) is staffed by an Allied Health Professional (ARNP/PA), a registered nurse, and an ED tech. An ED physician located in the main treatment area is immediately available for consultation.

By modeling patient flow, IT Process Reengineering has created a virtual ED that provides updated recommendations for optimal staffing hours. This model, using FlexSim simulation software, is able to assess the impact the RCU and related changes will have across the ED.


The location in triage facilitates movement of patients from the RCU to the main treatment area if the ESI level was underestimated. It also allows the assigned Allied Health Professional to assist in the main treatment area if RCU volume is low.

Opening date: December 1, 2014

The median time from arrival to diagnostic evaluation (OP20) for FY2014 for adult patients of acuity 4 and 5 was 39 minutes. After implementation, the median time decreased significantly to 36 minutes (z = 4.405, p < 0.001), still higher than the national median of 23 minutes.

The median LOS (OP18) for FY 2014 for adult patients of acuity 4 and 5 was 182 minutes. After implementation, the median time decreased significantly to 133 minutes (z = 19.997, p <0.001), comparable to the national median time of 131 minutes.

The Press-Ganey patient satisfaction scores remained above the 95th percentile throughout the process.



Our RCU model demonstrates that dedicating a portion of the triage area to this population can result in improved flow, and efficient care without sacrificing patient satisfaction. The relative decrease in OP20 (7.6%), although statistically significant, was less important clinically than the decrease in OP18 (26.9%). The decrease in LOS put our median very close to the national median.

1. American Medical Association (2014). NQF-Endorsed Voluntary Consensus Standards for Hospital Care. Retrieved on September 20, 2014 from, https:// www.qualitynet.org 2. Hospital Outpatient Quality Reporting Benchmarks and Trends: Fourth Quarter 2012-Fourth Quarter 2013. (July, 2014). Retrieved on March 1, 2015 from, https:// www.qualitynet.org/dcs/ContentServer?c=Page&pagename=QnetPublic% 2FPage%2FQnetTier2&cid=1228768205213 3. White, B. A., Biddinger, P. D., Chang, Y., Grabowski, B., Carignan, S., & Brown, D. (2013). Boarding Inpatients in the Emergency Department Increases Discharged Patient Length of Stay. The Journal of Emergency Medicine, 44 (1), 230-235 4. National Center for Health Statistics. Health, United States 2012: With Special

Feature on Emergency Care. Hyattsville, MD. 2013





### Outcomes

### Discussion

### References