

## Review

# Plant-driven selection of microbes in the rhizosphere and plant-microbe feedbacks

A. ESSARIOUI<sup>1</sup>, A. AMAMOU<sup>2</sup>, R. MEZIANI<sup>1</sup>, M. AFECHTAL<sup>3</sup>, F. MOKRINI<sup>4</sup>, N. HAMAMOUCHE<sup>5</sup>

(Reçu le 06/03/2017; Accepté le 03/04/2017)

## Abstract

Plant impacts on soil microbial communities and plant-microbe feedbacks have become the focus of much research. Recent advances in plant-microbe interactions investigations show that plants are able to shape their rhizosphere microbiome through diverse mechanisms. In this review, we gather findings from across multiple studies on the role of plants in altering the structure and functions of microbes in the rhizosphere. In addition, we discuss the roles of diverse phytochemicals in mediating these effects. Finally, we highlight that selective enrichment of specific microorganisms in the rhizosphere has either negative feedbacks, with pathogen accumulation in the rhizosphere; or, perhaps most importantly, positive feedbacks as a result of the recruitment of a beneficial microflora. Insights into the mechanisms that underpin plant selection of microbial communities with positive feedbacks will provide new opportunities to increase crop production.

**Keywords:** Plants, microbes, rhizosphere, feedbacks.

## Résumé

L'effet des plantes sur les microorganismes du sol et les rétroactions entre les microbes et les plantes sont devenus le focus de beaucoup de recherche. Les dernières avancées dans le domaine des interactions plantes-microbes montrent que les plantes sont capables de façonner le microbiome de leur rhizosphère via des mécanismes divers. Dans cette revue, nous rassemblons des résultats de multiples études sur le rôle des plantes dans l'altération de la structure et des fonctions des microorganismes au niveau de la rhizosphère. En outre, nous discutons les rôles de différentes substances phytochimiques dans la médiation de ces effets. Finalement, nous soulignons que l'enrichissement sélectif de microorganismes spécifiques a aussi bien des rétroactions négatives, avec l'accumulation de pathogènes dans la rhizosphère, que positives, suite au recrutement de microflore bénéfique. Une meilleure compréhension des mécanismes qui régissent la sélection de microorganismes ayant des rétroactions positives ouvrira de nouvelles possibilités pour l'amélioration de la production agricole.

**Mots-clés:** Plantes, microorganismes, rhizosphère, rétroactions.

## INTRODUCTION

A broad range of biotic and abiotic forces shape soil microbial communities. Within a given soil type and set of climatic conditions, there is compelling evidence that plants set the stage for growth conditions experienced by soil microbial communities through a variety of mechanisms (Dini-Andreote and van Elsas, 2013). In particular, plants are capable of modulating microbial population density, identity, and relative abundance; and influencing the functional activities that are carried out by soil microbial communities. These characteristics are inter-related, although specific relationships between microbial community structure and function are often unclear (Fuhrman, 2009). Understanding the factors that promote and maintain microbial diversity is a critical task for enhancing ecosystem processes and functioning. Significant role may be played by plants in this regard, through plant direct impacts on microbial activity and fitness, or through indirect alterations of soil properties or microbial interactions.

Interactions between soil microorganisms and plants occur primarily in the rhizosphere (Barea et al., 2005; Prithiviraj et al., 2007). The root system, which was long considered only as a means of anchorage and uptake of nutrients and water from the soil, is now consensually recognized as a key element in the interplay between plants and their milieu (Bais et al., 2006), making the root environment the most likely place to observe effects of plants on associated soil microbial communities (Kowalchuk et al., 2002; Bais et al., 2006). Indeed, the rhizosphere is a biologically and chemically highly diverse environment where complex and dynamic interactions occur among plant roots, microbes, and the soil (Hartmann et al., 2009). Compelling evidence was reported on the ability of plants to alter associated soil microbial communities through the well-known rhizosphere effect, wherein microbial populations and activity are markedly higher in soil adhering to plant roots compared to bulk soil (Starkey, 1958). Within this habitat, plants and microbes have evolved intimate relationships that enable them to coexist (Hartman et al., 2009). Specifically, plants provide nutrient resources to the rhizosphere that support

<sup>1</sup> National Institute of Agronomic Research, Morocco; Regional Center of Errachidia

<sup>2</sup> National Institute of Agronomic Research, Morocco; Regional Center of Settlat

<sup>3</sup> National Institute of Agronomic Research, Morocco; Regional Center of Kénitra

<sup>4</sup> National Institute of Agronomic Research, Morocco; Regional Center of Agadir

<sup>5</sup> University Sultan Moulay Slimane, Béni Mellal, Morocco

microbial activities. In turn, microbes provide a wide range of services that impact directly plant health and productivity (Nihorimbere et al., 2011).

A coupling of the concepts that plants are capable of exerting selection on associated microbes and that associated microbes may impact plant fitness, suggests the existence of interactions in which plants achieve fitness benefits through targeted manipulation of associated microbes. Such interactions are likely to be dependent upon the provision of resources to soil microbes, suggesting a selective force that may explain the seemingly inefficient loss of nutrients through plant roots. Expanding our knowledge on the specific mechanisms by which plants influence microbes and the resulting consequences for plant fitness will provide valuable insight into novel strategies for enhancing plant productivity in agricultural systems. Here; we (i) provide an overview on plant-driven impacts on soil microbial community characteristics (density, composition and structure, and function), (ii) explain mechanisms by which plants exert substantial effects on associated microbes, and (iii) examine plant response to rhizosphere microbiome, known as plant-microbe feedbacks.

## PLANT-DERIVED IMPACTS ON SOIL MICROBIAL COMMUNITIES

### Methods for assessing plant effects on soil microbes

A wide variety of techniques, ranging from traditional approaches to modern molecular methods, have been used to evidence differential host plant selective effects. These techniques have been reviewed extensively in previous work (Mazolla, 2004; Rastoji and Sani, 2011). Traditional culture-based approaches have been effective in demonstrating effects of particular plant species on soil bacterial populations (Loranger-Merciris et al., 2006). However, the major drawback of culture-based techniques is that >99% of the microorganisms in any environment observed through a microscope are not cultivable by standard culturing techniques (Hugenholtz, 2002). This limits comprehensive understanding of community structure and functions. Further insight into differential influence of plant host on microbial community can be reached through community profiling based on the characterization of biomolecules such as lipids without relying on culturing (Banowetz et al., 2006). Indeed, fatty acids are present in a relatively constant proportion of the cell biomass, and fatty acid signatures that can differentiate major taxonomic groups within a community exist in microbial cells. Specifically, soil microbial phospholipid fatty acid (PLFA) profiles and fatty acid methyl ester (FAME) profiles have been used to detect distinct microbial footprints and show specific microbial responses to various plant hosts (Germida et al., 1998; Carney and Matson, 2006). With recent advances in the development and application of molecular tools, the fields of microbial ecology and plant-microbe interactions have made an unprecedented bound. The use of techniques such as denaturing/temperature gradient gel electrophoresis (DGGE/TGGE), ribosomal intergenic spacer analysis (RISA), terminal restriction fragment length polymorphism (TRFLP), real-time PCR, amplicon sequencing, and

metatranscriptomics of environmental DNA showed that specific microbial communities are associated with distinct plant species, and that plant species have significant effects on the structure and function of resident soil microbial community (Miethling et al., 2000; Kuske et al., 2002; Broeckling et al., 2008; Garbeva et al., 2008; Micallef et al., 2009; LeBlanc et al., 2015).

### Effect of plants on soil microbial density

Plant host identity exerts significant impact on the density of associated microbes. Not only does plant community above-ground biomass correlates positively with soil microbial biomass (Wang et al., 2011), plant hosts also differ in the density of soil microbes that are supported in their rhizosphere. This has been demonstrated for particular taxa as well as for overall microbial population densities. For example, alfalfa supported larger populations of inoculated *Sinorhizobium meliloti* compared to rye (Miethling et al., 2000), and the population density of antibiotic producing *Pseudomonas* in soil was found to differ among plant hosts (Bergsma-Vlami et al., 2005a). In addition, some prairie perennial plant species were shown to differ in the density of antagonistic *Streptomyces* associated with their rhizospheres (Bakker et al., 2014). Furthermore, past work in recently deglaciated soils showed that plant species have differential effects on soil microbial biomass (Bardgett and Walker, 2004). Consistent with this, higher bacterial cell counts were found to be associated with wild oat roots than bulk soil (DeAngelis et al., 2006). Taken together, these findings from across multiple systems prove that some plant species support denser soil microbial communities than others. The difference in impact on soil microbial biomass among plant species suggests that the composition, structure, and function of soil microbial community will depend on the identity of plant host.

### Effect of plants on soil microbial community composition, structure, and function


For several years, great research efforts have been devoted to depict the impact of plant host on soil microbial community composition, structure, and function. Regardless of the investigation tools and approaches, findings from diverse studies seem to support the validity and ubiquity of plant-derived effects on soil microbial communities (Figure 1). For instance, plant hosts have been shown to alter the identity of ammonia-oxidizing bacteria (Briones et al., 2002) and denitrifying bacteria present in soil (Bremer et al., 2009). Clear impact of plant species effects on the composition, structure, diversity and taxonomic identity of pathogen-antagonistic microbial groups was also found (Berg et al., 2002; Berg et al., 2006). Interestingly, the strength of selective effect has been shown to differ among host plants. Specifically, compared to microbial communities associated with tomato, flax caused a greater shift away from the baseline conditions of uncultivated soil (Lemanceau et al., 1995). The ability of host plants to differentially select among soil microbes is also suggested by host specificity in mycorrhizal associations, although partners in such interactions span a continuum from specialist to generalist. That is, some mycorrhizal fungal species associate with

specific plants, while others have a broad host range (Johnson et al., 2005). Furthermore, invasive plants have been shown to significantly alter soil microbial communities in invaded soils (Batten et al., 2006) and occasionally reduce significantly soil microbial diversity (Broz et al., 2007). Monitoring soil microbial communities over the course of a change in plant cover can also reveal the selective effect of host plant. In some cases, it appears that the soil microbial community stabilizes after a period of adaptation to a host plant, with subsequent host switching leading to dramatic microbial community shifts (Badri et al., 2008). The study of the temporal dynamic of plant host effects on associated microbes, supported meta-transcriptomics data on potato associated microbes, suggests development-dependent changes over time in the rhizobiomes (Rasche et al., 2006; Chaparro et al., 2014).

Not only do different plant species have disparate effects on soil microbes, but also variants within the same species show distinct selective effects on soil communities. For instance, significantly distinct rhizosphere communities were found to be associated with specific genotypes of *Arabidopsis*, barley, canola, maize, potato, and sugar cane (Siciliano et al., 1998; Rasche et al., 2006; Micallef et al. 2009; Bulgarelli et al., 2012; Peiffer et al., 2013; Bulgarelli et al., 2015; Lebeis et al., 2015; Yeoh et al., 2015). Additionally, studies of genetically modified plants have demonstrated that small changes in plant genotype within the same species can result in significant impacts on associated microbial communities (Giovanni et al., 1999). In one fascinating example, the culture of opine-producing transgenic lotus plants induces an increase in the rhizosphere of bacterial communities that are able to utilize these molecules as sole carbon source. The density of members of these was about 10,000 times higher in the rhizosphere of the opine-producing plants than in that of wild-type (control) (Oger et al., 2004). This is suggestive of strong and rapid plant-imposed selection on soil microbial communities.

The effects of host plants extend to many functional measures of soil microbial activity. Plant induced effects can span from very general functions, as was seen for total

microbial respiration (Innes et al., 2004), to specific microbial functional traits. For example, plant identity and presence were found to exert substantial impact on soil denitrifying microbial communities; in bare soil, the denitrifying enzyme activity was 80% lower than in planted soil (Bremer et al., 2009). The plants influenced enzyme activity indirectly through their effect on the composition of the denitrifier community. Consistent with this, Alfalfa and rye supported microbial communities with significantly different community level physiological profiles (Miethling et al., 2000). Additionally, the proportion of auxin-producing *Pseudomonads* was higher for heterozygous maize plants compared to either of the parent lines (Picard and Bosco, 2005). Further, rice cultivars supported different amounts of activity by associated ammonia-oxidizing bacteria (Briones et al., 2002). Pathogen antagonism and related functional traits have been specifically studied as variables that respond to plant host selection. Recently, compelling evidence of plant host and plant community richness impact on soil *Streptomyces* nutrient use profiles and growth dynamics has been reported, suggesting that *Streptomyces* respond to selection imposed by plants (Essarioui et al., 2014, 2016). The proportion, composition, richness and diversity of pathogen-antagonistic microbes have been shown to be plant species and plant diversity dependent (Garbeva et al., 2008). An enhanced proportion, but reduced diversity of *Verticillium* antagonists, was found in rhizosphere compared to bulk soil (Berg et al., 2006). This lowered antagonist diversity in the rhizosphere is an evidence of plant-driven selection favoring some microbes over others (Berg et al., 2005). Cultivars of wheat differed in their ability to support antibiotic producing *Pseudomonads* (Mazzola et al., 2004). In other work, *Streptomyces griseoviridis* was shown to colonize the root surface of turnip rape (*Brassica rapa* subspecies *oleifera*) more readily than of carrot (Kortemaa et al., 1994) and the ability of various antibiotic producing *Pseudomonad* genotypes to colonize the rhizosphere of sugar beets was shown to be variable (Bergsma-Vlami et al., 2005a). Furthermore, the amount of antibiotic produced on a per cell basis by *Pseudomonads* in the rhizosphere differed among plant species (Bergsma-Vlami et al., 2005b). These find-



**Figure 1: Example of plant host effect on associated microbes in the rhizosphere. Comparison of the average bacterial community composition and relative abundances, at the phylum level, in root samples from three different plant species; (a) *Pilosella aurantiaca*, (b) *Leucanthemum vulgare*, and (c) *Trifolium hybridum*. Data from Aleklett et al., (2015).**



ings strengthen the view that specific microbial functional phenotypes are under significant selection by plant hosts in the rhizosphere.

### Mechanisms underlying plant-derived impacts on soil microbial communities

The number of studies through the literature that have gone beyond measuring plant impacts on soil microbial community composition, structure and function, to explore the actual mechanisms underpinning microbe selection by plants remain limited. Results from across multiple plant-microbe interaction systems point to the chemical nature of resources provided by plants as the major driver of plant selection of microbes. However, few studies explicitly tested this assumption leaving a great gap of knowledge on the relative importance of plant originated resource inputs in shaping soil microbial communities.

Nutritional resources available to soil communities are largely of plant origin; sloughed-off roots, mucilage, leachates, senescent tissue, litter, and root exudates. A large body of work has emphasized root exudates as having a particular importance in determining rhizosphere microbial community characteristics (Walker *et al.*, 2003), although their significance relative to other rhizo-deposits has not always been clearly demonstrated (Dennis *et al.*, 2010). There is a general belief that root exudation may be the key determinant of microbial community structure in the rhizosphere, whereas inputs such as plant litter and senescent tissue may play a larger role in influencing bulk soil microbial communities.

Plants exude a wide range of compounds into the soil by diffusion, ion channels and vesicular transport (Bertin *et al.* 2003). Root exudates are often divided into two classes of compounds. Low-molecular weight compounds such as amino acids, organic acids, sugars, phenolics, and other secondary metabolites account for much of the diversity of root exudates, whereas high molecular weight exudates, such as mucilage (polysaccharides) and proteins, are less diverse but often compose a larger proportion of the root exudates by mass (Bais *et al.*, 2006). These compounds modify soil chemical properties and define microbial nutritional niches in the rhizosphere (Lynch and Whipps, 1990; Bardgett *et al.*, 1998; Vandenkoornhuys *et al.* 2007; Bever *et al.*, 2012; Miransari, 2013).

The quantity and diversity of plant exudates vary among plant species and genotype (Haichar *et al.*, 2008; Micallef *et al.*, 2009; Compant *et al.*, 2010; Bever *et al.*, 2012; Philippot *et al.*, 2013; Pérez-Jaramillo *et al.*, 2015). Total nutrient inputs from plants will constrain microbial densities in soil, and the chemical variety of resources available to microbes will define in important ways microbial community composition, structure and function in the rhizosphere. The discovery that different plant species can have distinct microbial communities associated with their roots suggests that plant shape and dimension nutritional environment in the rhizosphere, and consequently create fitness penalties and rewards that are unequally distributed among taxa (Grayston *et al.*, 1998; Bertin *et al.*, 2003; Kumar *et al.*, 2007; Marshner *et al.*, 2011; Berendsen *et al.*, 2012). The provision of specific carbon substrates may offer a selec-

tive advantage to microbes with enzymatic capabilities to utilize those compounds. Methods for collecting plant root exudates *in vitro* have been previously described (Meharg and Killham, 1991; Nagahashi and Douds, 2000). This has opened new paths for studying the effects of plant exudates on soil microbial communities. Importantly, such collected exudates can have effects similar to whole plants when applied to soil (Badri *et al.*, 2008; Broeckling *et al.*, 2008). Quite recently, direct amendment of natural blends of phytochemicals, predominantly phenolic-related compounds, obtained from *Arabidopsis* to soil altered the bacterial community by stimulating or inhibiting different community members (Badri *et al.*, 2013).

In some cases, observed functional differences emanating from plant species and plant richness manipulation suggest plausible connections to root exudation. For example, carbon source utilization by soil microbes varied among host plant species, and the carbon substrates responsible for the observed patterns matched known major components of root exudates (Grayston *et al.*, 1998). Similarly, differences in soil bacterial community level physiological profiles among pioneering plant species were driven by the utilization of particular carbon compounds (Yan *et al.*, 2008), although it was not demonstrated that these compounds were constituents of the root exudates of the plants in question. Additionally, assumed differences in the chemical nature and diversity of rhizo-deposits among plant species and between monocultures and polycultures were linked to differential selections imposed by plant host and plant community richness on *Streptomyces* and *Fusarium* nutrient use profiles and antagonistic capacities (Essarioui *et al.*, 2014, 2016, 2017). Furthermore, plant community richness has also the potential to influence soil microbial interactions and function by defining their resource competitive environment through the quantity and diversity of resources they contribute to the soil. In recent a study, plant richness was found to impact *Fusarium* and *Streptomyces* resource niche overlap and inhibitory capacity in different ways (Essarioui *et al.*, 2017). *Fusarium* species had greater niche (nutrient) overlap with *Streptomyces* in monoculture than in polyculture, which was suggested to be a significant factor in generating highly antagonistic *Streptomyces* communities in monoculture. Conversely, *Streptomyces* had greater niche (nutrient) overlap with *Fusarium* in polyculture than in monoculture, which was hypothesized to result in selection for more inhibitory *Fusarium* populations in polyculture. This suggests that beyond their direct impact on soil microbes, plants are able to shape their microbiome indirectly via the mediation of cross-kingdom microbial interactions.

The emerging picture from multiple studies is that bioactive molecules with inhibitory properties from plants can also play a role in structuring rhizosphere microbiome by impacting negatively microbe fitness (Broeckling *et al.*, 2008; De-la-Pena *et al.*, 2008; Badri *et al.*, 2009; De-la-Pena *et al.*, 2010). For example, root glucosinolate content in *Brassica napus* was negatively correlated with root infection by *Azorhizobium caulinodans* (O'Callaghan *et al.*, 2000). Selective effects imposed by plants may also be indirect, resulting from changes to the physical or chemical environment, such as modifications to water content, soil pH (Starkey 1958) or other factors. For example, rice cul-

tivars that supported differing activity levels by ammonia-oxidizing bacteria were also found to exhibit differences in oxygen availability in the root zone (Briones *et al.*, 2002), suggesting that the observed host plant effects were modulated through atmospheric chemistry.

Plants may also impact soil microbial communities via phytochemicals that act as signals and alter gene expression, change microbial phenotypes and modulate outcomes of microbial competitive interactions. Probably the best-deciphered plant-microbe signaling systems are those underlying the well-established symbioses involving nitrogen fixing bacteria in legumes and mycorrhizal fungi. The role of plant signals in the formation of root nodules have been deeply explored in the past decades, and the outcomes of these studies have been systematically reviewed and updated (Downie, 2010; Oldroyd, 2013; Venturi and keel, 2016). Legumes release flavonoids (2-phenyl-1,4-benzopyrone derivatives) that alter patterns of gene expression in rhizobia, initiating a series of complex and specific interactions that ultimately lead to the fixation of atmospheric nitrogen inside of nodules (Rossen *et al.*, 1985; Redmond *et al.*, 1986; Ding and Oldroyd, 2009). Specific plant derived flavonoid compounds elicit also variable responses among distinct mycorrhizal partners during pre-symbiotic growth (Scervino *et al.*, 2005) resulting in specific plant-mycorrhizae associations driven by efficient chemical cross-talk. Other molecules secreted by plant roots act as signals for arbuscular mycorrhizal fungi (AMF) as well. For example, the strigolactone plant hormones have been recently shown to play a central role in initiating sym-

biotic relationships between plants and AMF (Ruyter-Spira *et al.*, 2013; Schmitz and Harrison, 2014). Additionally, cutin monomers have been implicated as a specific class of plant signaling factors which play a primary role in the stimulation of AMF (Gobatto *et al.*, 2012; Wang *et al.*, 2012).

In addition to the well-studied signaling pathways in the classical rhizobial and mycorrhizal symbiotic relationships with plants, patterns of signal-driven plant-microbe cross-communication in the rhizosphere, albeit not in such close association, have been elucidated. Many plant-associated bacteria require density-dependent signaling mechanisms, known as quorum sensing (QS) (Fuqua *et al.*, 1994, 2001), for the regulation of gene expression involved in various traits that facilitate the colonization of the plant-associated environments. Examples of such phenotypes are virulence, conjugation, secretion of hydrolytic enzymes, and the production of secondary metabolites (Van Bomdon, 2003; Newton and Fray, 2004). Gene expression coordination at the bacterial community level occurs through the production and response to quorum levels of signals, mainly N-acyl-homoserine lactones (AHLs) (Fuqua *et al.*, 2001). Importantly, plant-produced compounds have been reported to mimic and interfere with QS acting as agonists or antagonists of bacterial AHLs QS systems (Teplitski *et al.*, 2000; Keshavan *et al.*, 2005; Subramoni *et al.*, 2011; Gonzales and venture, 2013), though only a few of these compounds have been identified and the significance of their interference is still an open question.




Figure 2: Mechanisms by which plants impact soil microbial community in the rhizosphere


Biological evolutionary processes endowed plants to also produce and secrete a wide range of volatile compounds from their leaves, flowers, fruits, and roots (Dudariva et al., 2006). These chemicals are mainly represented by terpenoids, phenylpropanoids, and fatty acid and amino acid derivatives. Volatile compounds provide the producer with fitness benefits by acting against herbivores and pathogens, and conferring reproductive advantages. Specifically, volatiles can serve as antimicrobial molecules or as attractants for enemies of root-feeding herbivores (Rasmann et al., 2005). This highlights the role that plant produced volatiles are likely to play in structuring soil microbial communities.

In summary, root exudates, bioactive molecules, signaling molecules, and volatile compounds have been recognized as major phytochemicals by which plant tailor their rhizomicrobiome (Figure 2). This complex and intimate communication suggest feedback loops between plants and microbes, which ultimately will alter soil processes and impact both partners fitness.

## PLANT-MICROBE FEEDBACKS

The modulation of soil microbiome propounds the view that some plants invest more than others in broadening the spectrum of supported microbes for potentially greater beneficial feedbacks. This can lead to selective enrichment

of specific microbial species and functional phenotypes in the rhizosphere with the potential to improve overall plant health and development. Recent work suggests that plants maintain high microbial densities to sustain a level of competition that will reduce pathogen viability (Kinkel et al., 2011). For example, increased populations of *Bacillus subtilis* was shown to correlate with the suppression of the pathogen *Ralstonia solanacearum* in pepper, *Capcicum anuum* (Lee et al., 2012). As a corollary of microbial enrichment in the rhizosphere, strong resource competition may also impact microbial coevolutionary trajectories and end up selecting for highly antagonistic phenotypes, critical to pathogen suppression at the root level (Kinkel et al., 2014; Essarioui et al., 2017). Additionally, Recruitments of mutualistic microbes improve nutrient supply to plants and enhance their growth. Beside symbiotic associations with AMF and rhizobia that facilitate phosphate and nitrogen uptake, plant select through their root exudates for siderophores-producing microbes that enhance soluble iron availability in the soil (Hartmann et al., 2009; Carvalhais et al., 2013). These findings and many others from multiple studies (Bertin et al., 2003; Akiyama et al., 2005; Hassan and Mathesius, 2011) suggest the possibility of plant-microbe feedbacks having positive outcomes for plant fitness. However, negative feedbacks can exist as well between plants and soil microbial communities (McCarthy-Neumann and Kobe, 2010b), with pathogens



**Figure 3: Schematic representation of plant-microbes interactions and feedbacks in the rhizosphere. Plants communicate with microbes through the secretion of a variety of phytochemicals. Microbes respond with positive or negative feedbacks, impacting thereby plant fitness**



accumulating over time in the presence of a given host plant. Plant-microbe communication strategies are prone to hijacking by microbial plant parasites to increase their fitness. For example, isoflavones, that facilitate the recruitment of endosymbiotic nitrogen-fixing bacteria, were found to attract also the oomycete pathogen *Phytophthora sojae* and guide its growth towards the host (Morris et al., 1998; Subramanian et al., 2007; Cameroun et al., 2013). Despite the potential exploitation by opportunistic and parasitic microbes, creating an advantageous rhizosphere milieu may allow plant to select for microbial functional phenotypes that are more likely to provide a variety of fitness benefits and enhance their ecological competitiveness.

Plant-microbe feedbacks are also believed to be important to plant community dynamics (Reynolds et al., 2003), where the relative impact among plant species is the critical factor (McCarthy-Neumann and Kobe, 2010a); feedbacks may be species-specific, or may be more general, affecting all plants similarly (Casper et al., 2008). Invasive plant-induced change in microbial community may contribute to a positive feedback loop that selectively improve invasive plant performance and lead to the exclusion of native plant species that experience relatively poorer performance under the new microbial environment (Batten et al., 2007). On the other hand, specific negative feedbacks may slow competitive exclusion and work to sustain plant diversity (Bever et al., 2010).

## CONCLUSIONS AND PROSPECTS

Accumulating evidence demonstrates the importance of plant-microbe interplays in structuring soil microbial communities in the rhizosphere and impacting plant fitness (Figure 3). Considerable progress has been made with regard to bringing into focus the significance of these relationships to plant development and health. This highlights the agricultural potential of plant-microbe interactions management in inducing plant-beneficial microbial communities in soil and, thus, enhancing sustainable plant productivity. However, much research has focused only on plant impact on individual microbial strains and their feedbacks to that specific plant. Yet, there is increasing consensus that plant-microbe interactions are highly complex and involve consortia of plant and microbial species in a given assembly. Additionally, a wide range of factors such as edaphic conditions, climatic conditions, and microbe-microbe interactions, can influence plants, microbes, and the outcomes of their interactions. Therefore, further work is still needed to unravel the intricacies of the multipartite communication between microbes and hosts all along with the factors that govern their interactions. A greater understanding of the dynamics of root microbial communities has the potential to allow for the development of agricultural practices that support the recruitment of beneficial microbial communities. Overall, soil microbiome can be an-as-yet untapped resource to ameliorate plant health and productivity and promote global food security.

## REFERENCES

- Akiyama K., Matsuzaki K., Hayashi H. (2005). Plant sesquiterpenes induce hyphal branching in arbuscular mycorrhizal fungi. *Nature* 435: 824–827.
- Aleklett K., Leff J.W., Fierer N., Hart M. (2015). Wild plant species growing closely connected in a subalpine meadow host distinct root-associated bacterial communities. *Peer J*. DOI 10.7717/peerj.804
- Badri D.V., Chaparro J.M., Zhang R., Shen Q., Vivanco J.M. (2013). Application of natural blends of phytochemicals derived from the root exudates of *Arabidopsis* to the soil reveal that phenolic-related compounds predominantly modulate the soil microbiome. *J. Biol. Chem.* 288: 4502–4512.
- Badri D.V., Quintana N., El Kassis E.G., Kim H.K., Choi Y.H., Sugiyama A., Verpoorte R., Martinoia E., Manter D.K., Vivanco J.M. (2009). An ABC transporter mutation alters root exudation of phytochemicals that provoke an overhaul of natural soil microbiota. *Plant Phys.* 151: 2006–2017.
- Bais H.P., Weir T.L., Perry L.G., Gilroy S., Vivanco J.M. (2006). The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu Rev Plant Biol.* 57:233–266.
- Bakker M., Otto-Hanson L., Lange A., Bradeen J., Kinkel L. (2013). Plant monocultures produce more antagonistic soil *Streptomyces* communities than high-diversity plant communities. *Soil Biol. Biochem.* 65:304–312.
- Banowetz G.M., Whittaker G.W., Dierksen K.P., Azevedo M.D., Kennedy A.C., Griffith S.M., Steiner J.J. (2006). Fatty acid methyl ester analysis to identify sources of soil in surface water. *J. Environ. Qual.* 3:133–140.
- Bardgett R.D., Walker L.R. (2004). Impact of coloniser plant species on the development of decomposer microbial communities following deglaciation. *Soil Biol. Biochem.* 36: 555–559.
- Batten K.M., Scow K.M., Espeland E.K. (2007). Soil microbial community associated with an invasive grass differentially impacts native plant performance. *Microb. Ecol.* 55: 220–228.
- Berendsen R.L., Pieterse C.M.J., Bakker P.A.H.M. (2012). The rhizosphere microbiome and plant health. *Trends Plant Sci.* 17:478–486.
- Bertin C., Yang X., Weston L.A. (2003). The role of root exudates and allelochemicals in the rhizosphere. *Plant Soil.* 256:67–83.
- Bever J.D., Platt T.G., Morton E.R. (2012). Microbial population and community dynamics on plant roots and their feedbacks on plant communities. *Ann. Rev. Microbiol.* 66:265–283.
- Bever J.D., Dickie I.A., Facelli E., Facelli J.M., Klironomos J., Moora M., Rillig M.C., Stock W.D., Tibbett M., Zobel M. (2010). Rooting theories of plant community ecology in microbial interactions. *Trend Ecol. Evol.* 25: 468–478.
- Berg G., Opelt K., Zachow C., Lottmann J., Gotz M., Costa R., Smalla K., (2006). The rhizosphere effect on bacteria antagonistic towards the pathogenic fungus *Verticillium* differs depending on plant species and site. *FEMS Microbiol. Ecol.* 56: 250–261.

- Berg G., Zachow C., Lottmann J., Gotz M., Costa R., Smalla K. (2005). Impact of plant species and site on rhizosphere-associated fungi antagonistic to *Verticillium dahliae* Kleb. *Appl. Environ. Microbiol.* 71: 4203-4213.
- Berg G., Roskot N., Steidle A., Eberl L., Zock A., Smalla K. (2002). Plant-dependent genotypic and phenotypic diversity of antagonistic rhizobacteria isolated from different *Verticillium* host plants. *Appl. Environ. Microbiol.* 68: 3328-3338.
- Bergsma-Vlami M., Prins M.E., Raaijmakers J.M., (2005a). Influence of plant species on population dynamics, genotypic diversity and antibiotic production in the rhizosphere by indigenous *Pseudomonas* spp. *FEMS Microbiol. Ecol.* 52: 59-69.
- Bergsma-Vlami M., Prins M.E., Staats M., Raaijmakers J.M. (2005b). Assessment of genotypic diversity of antibiotic-producing *Pseudomonas* species in the rhizosphere by denaturing gradient gel electrophoresis. *Appl. Environ. Microbiol.* 1: 993-1003.
- Bremer C., Braker G., Matthies D., Beierkuhnlein C., Conrad R. (2009). Plant presence and species combination, but not diversity, influence denitrifier activity and the composition of nirK-type denitrifier communities in grassland soil. *FEMS Microbiol. Ecol.* 70: 377-387.
- Briones A.M., Okabe S., Umehiya Y., Ramsing N.B., Reichardt W., Okuyama H. (2002). Influence of different cultivars on populations of ammonia-oxidizing bacteria in the root environment of rice. *Appl. Environ. Microbiol.* 68: 3067-3075.
- Broeckling C.D., Broz A.K., Bergelson J., Manter D.K., Vivanco J.M. (2008). Root exudates regulate soil fungal community composition and diversity. *Appl. Environ. Microbiol.* 74: 738-744.
- Cameron D.D., Neal A.L., van Wees S.C.M., Ton J. (2013). Mycorrhizal induced resistance: more than the sum of its parts? *Trends Plant Sci.* 18:539-545.
- Carney K.M., Matson P.A. (2006). The influence of tropical plant diversity and composition on soil microbial communities. *Microb. Ecol.* 52: 226-238.
- Carvalho L.C., Dennis P.G., Fan B., Fedoseyenko D., Kierul K., Becker A., von Wiren N., Borriss R., (2013). Linking Plant Nutritional Status to Plant-Microbe Interactions. *PLoS One* 8:e68555
- Casper B.B., Bentivenga S.P., Ji B., Doherty J.H., Edensor H.M., Gustafson D.J. (2008). Plant-soil feedback: Testing the generality with the same grasses in serpentine and prairie soils. *Ecol.* 89: 2154-2164.
- Changting W., Genxu W., Wei L., Pengfei W. (2011). The Effects of Plant-Soil-Enzyme Interactions on Plant Composition, Biomass and Diversity of alpine Meadows in the Qinghai-Tibetan Plateau. *Inter. J. Eco.* Article ID 180926, 10 pages.
- Chaparro J.M., Badri D.V., Vivanco J.M. (2014). Rhizosphere microbiome assemblage is affected by plant development. *ISME J.* 8: 790-803.
- Compant S., Clément C., Sessitsch A. (2010). Plant growth-promoting bacteria in the rhizo- and endosphere of plants: their role, colonisation, mechanisms involved and prospects for utilisation. *Soil Biol. Biochem.* 42: 669-678.
- DeAngelis K.M., Brodie E.L., DeSantis T.Z., Andersen G.L., Lindow S.E., Firestone M.K. (2009). Selective progressive response of soil microbial community to wild oat roots. *ISME J.* 3: 168-178.
- De-la-Pena C., Lei Z., Watson B.S., Sumner L.W., Vivanco J.M. (2008). Root-microbe communication through protein secretion. *J. Biol. Chem.* 283: 25247-25255.
- Dennis P.G., Miller A.J., Hirsch P.R. (2010). Are root exudates more important than other sources of rhizodeposits in structuring rhizosphere bacterial communities? *FEMS Microbiol. Ecol.* 72: 313-327.
- Ding Y., Oldroyd G.E. (2009). Positioning the nodule, the hormone dictum. *Plant Signal. Behav.* 4: 89-93.
- Downie J.A. (2010). The roles of extracellular proteins, polysaccharides and signals in the interactions of rhizobia with legume roots. *FEMS Microbiol. Rev.* 34: 150-170.
- Dudareva N., Negre F., Nagegowda D.A., Orlova I. (2006). Plant volatiles: recent advances and future perspectives. *Crit. Rev. Plant Sci.* 25: 417-440.
- Ehrenfeld J.G., Ravit B., Elgersma K. (2005). Feedback in the plant-soil system. *Annu. Rev. Environ. Res.* 30: 75-115.
- Essarioui A., Kistler H.C., Kinkel L.L. (2017). Plant community richness mediates inhibitory interactions and resource competition between *Streptomyces* and *Fusarium* populations in the rhizosphere. *Microb. Ecol.* DOI: 10.1007/s00248-016-0907-5
- Essarioui A., Kistler H.C., Kinkel L.L. (2016). Nutrient use preferences among soil *Streptomyces* suggest greater resource competition in monoculture than polyculture plant communities. *Plant Soil.* Doi 10.1007/s11104-016-2968-0
- Essarioui A., LeBlanc N., Kistler H.C., Kinkel L.L. (2014). Plant host and plant community richness impact the dynamics of resource use by soil *Streptomyces* (Abstract). *Phytopathology.* 104, supplement 3: 67-67.
- Fuqua C., Parsek M.R., Greenberg E.P. (2001). Regulation of gene expression by cell-to-cell communication: acyl-homoserine lactone quorum sensing. *Annu. Rev. Genet.* 35: 439-468.
- Fuqua W.C., Winans S.C., Greenberg E.P. (1994). Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. *J. Bacteriol.* 176:269-275.
- Garbeva P., van Elsas J.D., van Veen J.A. (2008). Rhizosphere microbial community and its response to plant species and soil history. *Plant Soil.* 302: 19-32.
- Giovanni G.D.D., Watrud L.S., Seidler R.J., Widmer F., (1999). Comparison of parental and transgenic alfalfa rhizosphere bacterial communities using Biolog GN metabolic fingerprinting and enterobacterial repetitive intergenic consensus sequence-PCR (ERIC-PCR). *Microb. Ecol.* 37: 129-139.
- Gobbato E., Marsh J.F., Vernié T., Wang E., Maillet F., Kim J., Miller J.B., Sun J., Bano S.A., Ratet P., Mysore K.S., Dénarié J., Schultze M., Oldroyd G.E. (2012). A GRAS-type transcription factor with a specific function in mycorrhizal signaling. *Curr. Biol.* 22: 2236-2241.
- Gonzalez J.F., Venturi V. (2013). A novel widespread inter-kingdom signaling circuit. *Trends Plant Sci.* 18: 167-174.
- Grayston S.J., Wang S., Campbell C.D., Edwards A.C. (1998). Selective influence of plant species on microbial diversity in the rhizosphere. *Soil Biol. Biochem.* 30: 369-378.



- Haichar F.E., Marol C., Berge O., Rangel-Castro J.I., Prosser J.I., Balesdent J., Heulin T., Achouak W. (2008). Plant host habitat and root exudates shape soil bacterial community structure. *ISME J.* 2: 1221–1230.
- Hassan S., Mathesius U., (2011). The role of flavonoids in root rhizosphere signalling: opportunities and challenges for improving plant-microbe interactions. *J. Exp. Bot.* 63: 3429–3444.
- Hugenholtz P. Exploring prokaryotic diversity in the genomic era. *Genome Biol.* 3: Reviews 0003.
- Innes L., Hobbs P.J., Bardgett R.D. (2004). The impacts of individual plant species on rhizosphere microbial communities in soils of different fertility. *Biol. Fert. Soil.* 40: 7-13.
- Johnson D., Ijdo M., Genney D.R., Anderson I.C., Alexander I.J. (2005). How do plants regulate the function, community structure, and diversity of mycorrhizal fungi? *J. Exp. Bot.* 56: 1751-1760.
- Keshavan N.D., Chowdhary P.K., Haines D.C., González J.E. (2005). L-Canavanine made by *Medicago sativa* interferes with quorum sensing in *Sinorhizobium meliloti*. *J. Bacteriol.* 187: 8427–8436.
- Kinkel L., Schlatter D., Xiao K., Baines A. (2014). Sympatric inhibition and niche differentiation suggest alternative coevolutionary trajectories among *Streptomyces*. *ISME J.* 8: 492-492.
- Kinkel L., Bakker M., Schlatter D. (2011). A coevolutionary framework for managing disease-suppressive soils. *Annu. Rev. Phytopathol.* 49: 47-67.
- Kortema H., Rita H., Haahtela K., Smolander A. (1994). Root-colonization ability of antagonistic *Streptomyces griseoviridis*. *Plant Soil.* 163: 77-83.
- Kumar R., Bhatia R., Kukreja K., Behl R.K., Dudeja S.S., Narula N. (2007). Establishment of *Azotobacter* on plant roots: chemotactic response, development and analysis of root exudates of cotton (*Gossypium hirsutum* L.) and wheat (*Triticum aestivum* L.). *J. Basic Microbiol.* 47: 436–439.
- Kuske C.R., Ticknor L.O., Miller M.E., Dunbar J.M., Davis J.A., Barns S.M., Belnap J. (2002). Comparison of soil bacterial communities in rhizospheres of three plant species and the interspaces in an arid grassland. *Appl. Environ. Microbiol.* 68: 1854-1863.
- LeBlanc N., Kinkel L.L., Kistler H. C. (2015). Soil fungal communities respond to grassland plant community richness and soil edaphics. *Microb. Ecol.* 70:188-95. doi: 10.1007/s00248-014-0531-1.
- Lee B., Lee S., Ryu C.M. (2012). Foliar aphid feeding recruits rhizosphere bacteria and primes plant immunity against pathogenic and nonpathogenic bacteria in pepper. *Ann. Bot.* 110: 281–290.
- Lemanceau P., Corberand T., Gardan L., Latour X., Laguerre G., Boeufgras J.M., Alabouvette C. (1995). Effect of two plant species, flax (*Linum usitatissimum* L) and tomato (*Lycopersicon esculentum* Mill), on the diversity of soilborne populations of fluorescent *Pseudomonads*. *Appl. Environ. Microbiol.* 61: 1004-1012.
- Loranger-Merciris G., Barthes L., Gastine A., Leadley P. (2006). Rapid effects of plant species diversity and identity on soil microbial communities in experimental grassland ecosystems. *Soil Biol. Biochem.* 38: 2336-2343.
- Lynch J.M., Whipps J.M. (1990). Substrate flow in the rhizosphere. *Plant Soil.* 129:1–10.
- Marschner P., Crowley D., Rengel Z. (2011). Rhizosphere interactions between microorganisms and plants govern iron and phosphorus acquisition along the root axis—model and research methods. *Soil Biol. Biochem.* 43: 883–894.
- Mazzola M., Funnell D.L., Raaijmakers J.M., (2004). Wheat cultivar-specific selection of 2,4-diacetylphloroglucinol-producing fluorescent *Pseudomonas* species from resident soil populations. *Microb. Ecol.* 48: 338-348.
- McCarthy-Neumann S., Kobe R.K. (2010a). Conspecific and heterospecific plant-soil feedbacks influence survivorship and growth of temperate tree seedlings. *J. Ecol.* 98: 408-418.
- McCarthy-Neumann S., Kobe R.K. (2010b). Conspecific plant-soil feedbacks reduce survivorship and growth of tropical tree seedlings. *J. Ecol.* 98: 396-407.
- Meharg A.A., Killham K. (1991). A novel method of quantifying root exudation in the presence of soil microflora. *Plant Soil.* 133: 111-116.
- Mendes R., Kruijt M., De Bruijn I., Dekkers E., Van der Voort M., Schneider J. H.M., Piceno Y.M., DeSantis T.Z., Andersen G.L., Bakker P.A.H.M., Raaijmakers J.M. (2011). Deciphering the rhizosphere microbiome for disease-suppressive bacteria. *Science.* 332: 1097-1100.
- Newton J.A. Fray R.G. (2004). Integration of environmental and host-derived signals with quorum sensing during plant–microbe interactions. *Cell Microbiol.* 6: 213–224.
- Micallef S.A., Shiaris M.P., Colon-Carmona A. (2009). Influence of *Arabidopsis thaliana* accessions on rhizobacterial communities and natural variation in root exudates. *J. Exp. Bot.* 60: 1729–1742.
- Miethling R., Wieland G., Backhaus H., Tebbe C.C. (2000). Variation of microbial rhizosphere communities in response to crop species, soil origin, and inoculation with *Sinorhizobium meliloti* L33. *Microb. Ecol.* 40: 43-56.
- Miransari M. (2013). Soil microbes and the availability of soil nutrients. *Acta Physiol. Plant.* 35:3075–3084.
- Morris P.F., Bone E., Tyler B.M. (1998). Chemotropic and contact responses of *Phytophthora sojae* hyphae to soybean isoflavonoids and artificial substrates. *Plant Physiol.* 117: 1171–1178
- Nagahashi, G., Douds D.D. (2000). Partial separation of root exudate components and their effects upon the growth of germinated spores of AM fungi. *Mycol. Res.* 104:1453-1464.
- O’Callaghan K.J., Stone P.J., Hu X.J., Griffiths D.W., Davey M.R., Cocking E.C. (2000). Effects of glucosinolates and flavonoids on colonization of the roots of *Brassica napus* by *Azorhizobium caulinodans* ORS571. *Appl. Environ. Microbiol.* 66: 2185-2191.
- Oger P., Mansouri H., Nesme X., Dessaux Y. (2004). Engineering root exudation of lotus toward the production of two novel carbon compounds leads to the selection of distinct microbial populations in the rhizosphere. *Microb. Ecol.* 47: 96-103.

- Oldroyd G.E. (2013). Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. *Nat. Rev. Microbiol.* 11: 252–263.
- Pérez-Jaramillo J.E., Mendes R., Raaijmakers J.M. (2015). Impact of plant domestication on rhizosphere microbiome assembly and functions. *Plant. Mol. Biol.* 1–10. doi:10.1007/s11103-015-0337-7
- Philippot L., Raaijmakers J.M., Lemanceau P., van der Putten W.H. (2013). Going back to the roots: the microbial ecology of the rhizosphere. *Nat. Rev. Microbiol.* 11: 789–799.
- Picard C., Bosco M. (2005). Maize heterosis affects the structure and dynamics of indigenous rhizospheric auxins-producing *Pseudomonas* populations. *FEMS Microbiol. Ecol.* 53: 349–357.
- Rasmann S., Köllner T.G., Degenhardt J., Hiltbold I., Toepfer S., Kuhlmann U., Gershenzon J., Turlings Ted C.J. (2005). Recruitment of entomopathogenic nematodes by insect-damaged maize roots. *Nature.* 434: 732–737.
- Rastogi G., Sani R. (2011). Molecular techniques to assess microbial community structure function and dynamics in the environment. I. Ahmad, F. Ahmad, J. Pichtel (Eds.), *Microbes and Microbial Technology: Agricultural and Environmental Applications*, Springer, New York, pp. 29–57.
- Redmond J.W., Batley M., Djordjevic M.A., Innes R.W., Kuempel P.L., Rolfe B.G. (1986). Flavones induce expression of nodulation genes in *Rhizobium*. *Nature.* 323: 632–635.
- Reynolds H.L., Packer A., Bever J.D., Clay K. (2003). Grassroots ecology: Plant-microbe-soil interactions as drivers of plant community structure and dynamics. *Ecology.* 84: 2281–2291.
- Rossen L., Shearman C.A., Johnston A.W., Downie J.A. (1985). The nodD gene of *Rhizobium leguminosarum* is autoregulatory and in the presence of plant exudate induces the nodA,B,C genes. *EMBO J.* 4: 3369–3373
- Ruyter-Spira C., Al-Babili S., van der Krol S., Bouwmeester H. (2013). The biology of strigolactones. *Trends Plant Sci.* 18: 72–83
- Scervino J.M., Ponce M.A., Erra-Bassells R., Vierheilig H., Ocampo J.A., Godeas A. (2005). Flavonoids exhibit fungal species and genus specific effects on the presymbiotic growth of *Gigaspora* and *Glomus*. *Mycol. Res.* 109: 789–794.
- Schmitz A.M., Harrison M.J. (2014). Signaling events during initiation of arbuscular mycorrhizal symbiosis. *J. Integr. Plant Biol.* 56: 250–261.
- Siciliano S.D., Theoret C.M., de Freitas J.R., Hucl P.J., Germida J.J. (1998). Differences in the microbial communities associated with the roots of different cultivars of canola and wheat. *Can. J. Microbiol.* 44: 844–851.
- Starkey R.L. (1958). Interrelationships between microorganisms and plant roots in the rhizosphere. *Bacteriol. Rev.* 22: 154–172.
- Subramanian S., Stacey G., Yu O., (2007). Distinct, crucial roles of flavonoids during legume nodulation. *Trends Plant Sci.* 12:282–285.
- Subramoni S., Gonzalez J.F., Johnson A., Péchy-Tarr M., Rochat L., Paulsen I., Loper J.E., Keel C., Venturi V. (2011). Bacterial subfamily of LuxR regulators that respond to plant compounds. *Appl. Environ. Microbiol.* 77: 4579–4588.
- Teplitski M., Robinson J.B., Bauer W.D. (2000). Plants secrete substances that mimic bacterial N-acyl homoserine lactone signal activities and affect population density-dependent behaviors in associated bacteria. *Mol. Plant Microbe Interact.* 13:637–48.
- Von Bodman S.B., Bauer W.D., Coplin D.L., (2003) Quorum sensing in plant-pathogenic bacteria. *Annu. Rev. Phytopathol.* 41: 455–482
- Vandenkoornhuyse P., Mahe S., Ineson P., Staddon P., Ostle N., Cliquet J.B., Francez A.J., Fitter A.H., Young J.P., (2007). Active root-inhabiting microbes identified by rapid incorporation of plant derived carbon into RNA. *Proc. Nat. Acad. Sci. U S A* 104: 16970–16975
- Walker T.S., Bais H.P., Grotewold E., Vivanco J.M., (2003). Root exudation and rhizosphere biology. *Plant Phys.* 132: 44–51.
- Wang C., Wang G., Liu W., Pengfei W. (2011). The Effects of Plant-Soil-Enzyme Interactions on Plant Composition, Biomass and Diversity of alpine Meadows in the Qinghai-Tibetan Plateau. *Int. J. Ecol.* Article ID 180926, 10 pages.
- Wang E., Schornack S., Marsh J.F., Gobbato E., Schwesinger B., Eastmond P., Schultze M., Kamoun S., Oldroyd G.E. (2012) A common signaling process that promotes mycorrhizal and oomycete colonization of plants. *Curr. Biol.* 22: 2242–2246.
- Yan W., Artz R.R.E., Johnson D. (2008). Species-specific effects of plants colonizing cutover peatlands on patterns of carbon source utilisation by soil microorganisms. *Soil Biol. Biochem.* 40: 544–549.