

Original Paper

Open Access

Prey consumption and development of *Chrysoperla externa* (Neuroptera: Chrysopidae) on *Spodoptera frugiperda* (Lepidoptera: Noctuidae) eggs and larvae and *Anagasta kuehniella* (Lepidoptera: Pyralidae) eggs

Wagner S Tavares¹, Ivan Cruz², Rafael B Silva³, José E Serrão⁴, José C Zanuncio^{5*}

¹Departamento de Fitotecnia, Universidade Federal de Viçosa, 36570-000, Viçosa, Brazil

²Embrapa Milho e Sorgo, Rodovia MG 424, Km 65, Caixa Postal 151, 35701-970, Sete Lagoas, Brazil

³Programa de Pós-graduação em Ecologia e Recursos Naturais, Universidade Federal de São Carlos, 13565-905, São Carlos, Brazil

⁴Departamento de Biologia Geral, Universidade Federal de Viçosa, 36570-000, Viçosa, Brazil

⁵Departamento de Biologia Animal, Universidade Federal de Viçosa, 36571-000, Viçosa, Brazil

*Corresponding author: E-mail: zanuncio@ufv.br

Abstract

The Chrysopidae family comprises natural enemies of agricultural and forest pests. This work evaluated the prey consumption and development of one species, *Chrysoperla externa* (Neuroptera: Chrysopidae), fed with *Spodoptera frugiperda* (Lepidoptera: Noctuidae) or *Anagasta kuehniella* (Lepidoptera: Pyralidae) in laboratory conditions. *Chrysoperla externa* was reared with: newly laid or one-day-old *S. frugiperda* eggs; newly hatched, one- or two-day-old *S. frugiperda* larvae; or one-day-old *A. kuehniella* eggs. The number of prey offered varied with the development stage of *C. externa*. Larvae of *C. externa* and prey were transferred every 24 hours to fresh vials. Duration of the larval stage of *C. externa* was similar when fed with newly laid or one-day-old *S. frugiperda* eggs, newly hatched *S. frugiperda* larvae or *A. kuehniella* eggs. Larval survival of *C. externa* was $90.0 \pm 2.5\%$ when fed with *A. kuehniella* eggs and $73.3 \pm 18.32\%$ with newly hatched *S. frugiperda* larvae. *Chrysoperla externa* consumed high numbers of eggs of *A. kuehniella* and high weights of one-day-old eggs or newly hatched larvae of *S. frugiperda* or eggs of *A. kuehniella*. *Chrysoperla externa* could not be successfully reared in the laboratory on one- or two-day-old *S. frugiperda* larvae, but could on eggs of both preys and newly hatched *S. frugiperda* larvae.

Keywords: biological control, fall armyworm, flour moth, green lacewing, rearing

Introduction

Fall armyworm, *Spodoptera frugiperda* (Smith, 1797) (Lepidoptera: Noctuidae), is a major pest of corn and sorghum crops, and their larvae can feed on plants at all growth stages (Castillejos et al, 2001; Hoballah et al, 2004; Matos Neto et al, 2004). Other than synthetic pesticides, alternative methods to reduce populations of *S. frugiperda* include spraying botanical extracts on young plants (Tavares et al, 2009, 2010a, 2010b) and release of natural enemies including predatory insects, for example of the Chrysopidae, Carabidae, Coccinellidae, Pentatomidae, Anthicoridae, Reduviidae and Pentatomidae families (Figueiredo et al, 2006; Zanuncio et al, 2008; Silva et al, 2009).

Chrysoperla externa (Hagen, 1861) (Neuroptera: Chrysopidae) and other lacewings can control arthropod pests (Barbosa et al, 2008), such as *Alabama argillacea* (Hübner, 1818) (Lepidoptera: Noctuidae), *Aphis gossypii* (Glover, 1877), *Schizaphis graminum* (Rondani, 1852), *Rhodobium porosum* (Sanderson, 1901) (Hemiptera: Aphididae), *Coccus* spp. (Hemiptera: Coccidae), *Orthezia* spp. (Hemiptera: Ortheziidae), *Pinnaspis* spp. (Hemiptera: Diaspididae), *Sele-*

naspis spp. and *Leptopharsa heveae* (Drake and Poor, 1935) (Heteroptera: Tingidae) in various field crops (Gao et al, 2007; Pappas et al, 2007; Barbosa et al, 2008; Souza et al, 2008). *Chrysoperla externa* is a natural enemy that feeds on eggs and young larvae, and so can be utilized in biological control programs (Hoballah et al, 2004; Hagerty et al, 2005). Benefits of using *C. externa* as a mass-released biological control agent are that its larvae are tolerant to handling and adults produce large numbers of offspring (Auaud et al, 2003). *Chrysoperla externa* has been maintained in the laboratory on an artificial diet; however, *S. frugiperda* may be an alternative food source.

Prey consumption of lacewings has been studied in the laboratory (Nakahira et al, 2005; Pappas et al, 2008a, 2008b; Souza et al, 2008), greenhouse (Cole et al, 2006; Barbosa et al, 2008) and field (Hagerty et al, 2005; Kovanci et al, 2007). The predator has been trialled with different prey types (Silva et al, 2004; Kunkel and Cottrell, 2007) and diets (Sattar et al, 2007; Sattar and Abro, 2009), including algae (Zaki and Gesraha, 2001), honeydew (Hogervorst et al, 2008) and pollen grains (Berkvens et al, 2008). *Anagasta kuehniella* (Zeller, 1879) (Lepidoptera: Py-

ralidae) is a potential food source for lacewings but its rearing demands intensive labor and space (Pappas et al, 2007, 2008a, 2008b). Another potential food source for *C. externa*, *S. frugiperda*, was offered ad libitum to larvae (Aquad et al, 2003). The development of *C. externa* reared on different prey types has been studied (Silva et al, 2004; Gao et al, 2007; Souza et al, 2008), but data on the effect of different ages of *S. frugiperda* on the development of *C. externa* are scarce.

The objective of this study was to evaluate the consumption rates and development of *C. externa* reared on different age *S. frugiperda* eggs and larvae or *A. kuehniella* eggs.

Materials and Methods

Experiments were carried out at EMBRAPA Maize and Sorghum in Sete Lagoas, Minas Gerais State, Brazil in the laboratory (25 ± 1°C, 70 ± 10% R.H. and 12:12 L:D).

Anagasta kuehniella, *C. externa*, and *S. frugiperda* were obtained from laboratory-reared colonies at EMBRAPA Maize and Sorghum. *Anagasta kuehniella* larvae were reared in plastic trays (30 x 20 x 10 cm) covered with organza and fed with a diet of 600 g of corn meal, 600 g of wheat bran and 3% yeast (Wade et al, 2008; Tavares et al, 2009). *Chrysoperla externa* adults were kept in PVC tubes (700 mm wide x 30 cm height) and their larvae were fed an artificial diet of 45 g of honey, 45 g of yeast, and 10 ml of water (Lawo and Romeis, 2008). *S. frugiperda* adults were kept in cages, and their larvae were kept in 50 ml disposable plastic cups sealed with a transparent acrylic cover. Adults were fed with a solution of 25 g of sugar, 0.5 g of ascorbic acid, and 500 ml of water and their larvae were fed with 8 g of artificial diet consisting of 2 kg of beans, 950.4 g of wheat germ, 608.8 g of yeast, 61.2 g of ascorbic acid, 37.8 g of methyl parahydroxybenzoate (nipagim), 240 g of agar, 49.8 ml of formaldehyde, 16 l of water, and 49.8 ml of inhibitor solution (418 g of propionic acid, 42 g of phosphoric acid, and 540 ml of water).

Newly hatched F2 generation *C. externa* larvae were removed from the colony and isolated in glass tubes with flat bottoms (2 cm wide x 10 cm height). These larvae were reared on one of six diets: (T1) newly laid or (T2) one-day-old *S. frugiperda* eggs; (T3) newly-hatched, (T4) one- or (T5) two-day-old *S. frugiperda* larvae; or (T6) one-day-old *A. kuehniella* eggs. The hypothesis was that older eggs and larvae of *S. frugiperda* may have lower nutritional value and/or raised physical barriers against predation by *C. externa* larvae (Aquad et al, 2003). The number of prey offered depended on the life stage of *C. externa*: 40 units (number of eggs/larvae) daily for first instars; 80 units daily for second instars; and 120 units daily for third instars. The presence of scales in the eggs of *S. frugiperda* was not observed (i.e. mixed eggs with or without scales) (Beserra and Parra, 2004). *Chrysoperla externa* larvae and their prey were transferred with a slip-away brush every 24 h to new glass tubes.

We recorded observations every 24 h to determine: survival of *C. externa* larvae; duration of each *C. externa* life stage; and prey consumption (number and weight of eggs or larvae consumed). The design was entirely randomized, with four replications, each with five *C. externa* larvae per treatment.

Data were subjected to analysis of variance (ANOVA) and to Tukey's post-hoc tests ($P < 0.05$) with the computer program MSTAT-C, version 2.1 (Supplier: EMBRAPA Maize and Sorghum) (Russel, 1989).

Results

Chrysoperla externa had three instars. The duration of each life stage was similar when larvae were fed with newly laid or one-day-old *S. frugiperda* eggs, newly-hatched *S. frugiperda* larvae or *A. kuehniella* eggs (Table 1). Survival of first instar *C. externa* was higher with newly hatched *S. frugiperda* larvae or *A. kuehniella* eggs. However, the survival of *C. externa* varied between instars, with higher rates for second instars fed one-day-old *S. frugiperda* eggs or *A. kuehniella* eggs, and for third instar larvae reared on eggs of both prey (Table 2).

Table 1 - Duration (mean ± standard error of mean) of the developmental stages (DS) (first instar – FI, second instar – SI, third instar – TI, larval – L, pupal – P, and from larval to adult – LA) of *Chrysoperla externa* (Neuroptera: Chrysopidae) fed with newly laid (T1) or one-day-old *Spodoptera frugiperda* (Lepidoptera: Noctuidae) eggs (T2); newly hatched (T3), one- (T4) or two-day-old (T5) *S. frugiperda* larvae; or one-day-old *Anagasta kuehniella* (Lepidoptera: Pyralidae) eggs (T6).

DS	T1	T2	T3	T4	T5	T6
	Duration (days)					
FI	3.0 ± 0.20 ^a	2.9 ± 0.09 ^a	2.90 ^a	3.40 ^a	2.60 ^a	3.20 ^a
SI	3.1 ± 0.21 ^a	3.0 ± 0.10 ^a	3.10 ^a	-	-	2.80 ^a
TI	3.8 ± 0.28 ^a	3.4 ± 0.14 ^a	3.7 ± 0.07 ^a	-	-	3.30 ^a
L	9.9 ± 0.85 ^a	9.3 ± 0.70 ^a	9.7 ± 0.67 ^a	-	-	9.30 ^a
P	10.2 ± 0.92 ^a	10.0 ± 0.80 ^a	8.3 ± 0.53 ^b	-	-	10.6 ± 0.46 ^a
LA	19.7 ± 1.87 ^a	19.0 ± 1.70 ^a	18.7 ± 1.57 ^a	-	-	19.9 ± 1.39 ^a
CV	8.45%					

means followed by the same letter per line do not differ by the test of Tukey ($P < 0.05$); CV= Coefficient of variation

Table 2 - Survival (mean \pm standard error of mean) of the developmental stages (DS) (first instar – FI, second instar – SI, third instar – TI, larval – L, pupal – P, and from larval to adult – LA) of *Chrysoperla externa* (Neuroptera: Chrysopidae) fed with newly laid (T1) or one-day-old *Spodoptera frugiperda* (Lepidoptera: Noctuidae) eggs (T2); newly hatched (T3), one- (T4) or two-day-old (T5) *S. frugiperda* larvae; or one-day-old *Anagasta kuehniella* (Lepidoptera: Pyralidae) eggs (T6).

DS	T1	T2	T3	T4	T5	T6
	Survival (%)					
FI	55.0 \pm 2.50 ^b	55.0 \pm 2.50 ^b	95.0 \pm 2.96 ^a	0.00 ^c	0.00 ^c	95.0 \pm 2.50 ^a
SI	79.2 \pm 4.40 ^b	85.4 \pm 4.74 ^a	71.3 \pm 2.97 ^b	0.00 ^c	0.00 ^c	100.0 ^a
TI	100.00 ^a	100.0 ^a	21.7 \pm 5.42 ^b	0.00 ^c	0.00 ^c	93.8 \pm 2.46 ^a
L	45.0 \pm 2.50 ^c	45.0 \pm 2.50 ^c	73.3 \pm 18.32 ^b	0.00 ^d	0.00 ^d	90.0 \pm 2.50 ^a
P	45.0 \pm 2.50 ^b	45.0 \pm 2.50 ^b	20.0 \pm 5.00 ^c	0.00 ^d	0.00 ^d	90.0 \pm 2.50 ^a
LA	45.0 \pm 2.50 ^b	50.0 \pm 2.77 ^b	20.0 \pm 5.00 ^c	0.00 ^d	0.00 ^d	90.0 \pm 2.50 ^a
CV	6.47%					

means followed by the same letter per line do not differ by the test of Tukey ($P < 0.05$); CV = Coefficient of variation

Chrysoperla externa did not reach the pupal stage on a diet of one- or two-day-old *S. frugiperda* larvae. *Chrysoperla externa* did pupate on a diet of eggs or newly hatched larvae of *S. frugiperda* or *A. kuehniella* eggs, although the duration of the pupal stage was shorter when they were fed young larvae than with eggs of either prey (Table 1). Pupae survival of *C. externa* fed with *S. frugiperda* eggs was higher than with newly hatched *S. frugiperda* larvae. However, pupae survival was highest for *C. externa* fed with *A. kuehniella* eggs (Table 2).

Chrysoperla externa reached maturity on a diet of newly laid or one-day-old *S. frugiperda* eggs, newly hatched *S. frugiperda* larvae or *A. kuehniella* eggs, and the interval between larvae and adult did not differ between these treatments (Table 1). The survival from larvae to adult of *C. externa* was higher with *A. kuehniella* eggs than with other prey types (Table 2). Adults of *C. externa* showed normal morphology when reared on a diet of newly laid or one-day-old *S. frugiperda* eggs, newly hatched *S. frugiperda* larvae or *A. kuehniella* eggs.

Prey consumption (number of eggs or larvae) of *C. externa* increased with its development from 44.7 to 330.8 *A. kuehniella* eggs. First instar *C. externa* showed higher consumption of one-day-old eggs or newly hatched *S. frugiperda* larvae than with other prey types. *Chrysoperla externa* consumed higher numbers of *A. kuehniella* eggs during its larval stage than with *S. frugiperda* eggs or newly hatched *S. frugiperda* larvae (Table 3). Some *C. externa* larvae preyed on newly hatched *S. frugiperda* larvae but did not consume them.

First instar *C. externa* larvae consumed a greater weight of one- or two-day-old *S. frugiperda* larvae. Second instar *C. externa* larvae consumed a greater weight of newly hatched *S. frugiperda* eggs or larvae or *A. kuehniella* eggs. Third instar *C. externa* larvae consumed more one-day-old *S. frugiperda* eggs or *A. kuehniella* eggs. Overall, *C. externa* larvae consumed more one-day-old eggs or newly hatched *S. frugiperda* larvae and *A. kuehniella* eggs (Table 4).

Discussion

The three instars detected in *C. externa* were similar to those reported for this predator fed with *Bemisia tabaci* biotype B (Gennadius, 1889) (Hemiptera: Aleyrodidae) nymphs (Silva et al, 2004) and *Myzus persicae* (Sulzer, 1776) (Hemiptera: Aphididae) (Barbosa et al, 2008). Other lacewings have also exhibited three instar stages, including *Ceraeochrysa cubana* (Hagen, 1861) (Neuroptera: Chrysopidae) fed with *A. gossypii* (Alcantra et al, 2008) and *Dichochrysa prasina* (Burmeister, 1839) (Neuroptera: Chrysopidae) fed with *A. kuehniella*, *Ephestia kuehniella* (Burmeister, 1879) (Lepidoptera: Noctuidae) or *M. persicae* (Papas et al, 2007, 2008a). This suggests that newly hatched larvae or eggs of *S. frugiperda* or *A. kuehniella* are adequate for *C. externa* because unsuitable prey or unfavorable environmental conditions may increase or reduce the number of instars (Michaud, 2005; Vandekerckhove et al, 2006). In spite of this, additional nutrients provided in artificial diets, such as essential amino acids and mineral salts offered along with prey can be useful when mass-rearing the predators (Isikber and Copland, 2002; Ragkou et al, 2004; Berkvens et al, 2008).

The equal duration of larval stages of *C. externa* – except those fed with one- or two-day-old *S. frugiperda* larvae – differed from the relatively shorter first and third instars *C. externa* exhibited when fed with *B. tabaci* biotype B nymphs (Silva et al, 2004). The larval stage of *Chrysoperla rufilabris* (Burmeister, 1839) (Neuroptera: Chrysopidae) was shorter with *Monella caryella* (Fitch) or *Melanocallis caryaefoliae* (Davis, 1910) (Hemiptera: Aphididae) alone than with both prey types together, which was attributed to its generalist feeding behavior (Kunkel and Cottrell, 2007). The equal duration of the larval stages of *C. externa* fed with newly hatched *S. frugiperda* larvae or *A. kuehniella* eggs agreed with results for neuropteran predators with this and other prey (Aquad et al, 2003; Pappas et al, 2007, 2008a; Souza et al, 2008).

All *C. externa* fed with one- or two-day-old *S. frugiperda* larvae died, suggesting that this predator

Table 3 - Consumption (mean \pm standard error of mean) of the developmental stages (DS) (first instar – FI, second instar – SI, third instar – TI, and larval – L) of *Chrysoperla externa* (Neuroptera: Chrysopidae) fed with newly laid (T1) or one-day-old *Spodoptera frugiperda* (Lepidoptera: Noctuidae) eggs (T2); newly hatched (T3), one- (T4) or two-day-old (T5) *S. frugiperda* larvae; or one-day-old *Anagasta kuehniella* (Lepidoptera: Pyralidae) eggs (T6).

DS	T1	T2	T3	T4	T5	T6
	Consumption (number of eggs or larvae)					
FI	38.1 \pm 1.9 ^b	31.9 \pm 2.6 ^b	71.5 \pm 4.7 ^a	85.7 \pm 5.8 ^a	43.2 \pm 5.2 ^b	44.7 \pm 0.8 ^b
SI	43.4 \pm 4.8 ^b	48.5 \pm 13.8 ^b	119.9 \pm 28.7 ^a	-	-	129.9 \pm 2.5 ^a
TI	160.9 \pm 21.5 ^b	169.8 \pm 9.4 ^b	176.3 \pm 30.9 ^b	-	-	330.8 \pm 2.8 ^a
L	242.4 \pm 28.2 ^b	250.2 \pm 15.8 ^b	367.7 \pm 64.3 ^b	-	-	505.4 \pm 6.1 ^a
CV	9.16%					

means followed by the same letter per line do not differ by the test of Tukey ($P < 0.05$); CV = Coefficient of variation

failed to successfully attack the larger larvae. Larvae of one- or two-day-old *S. frugiperda* exhibited considerable movement away from attacking *C. externa* larvae and their exterior became increasingly tough with age. A lower survival rate of first instar *C. externa* larvae fed with *S. frugiperda* eggs may be attributed to the architecture of the egg masses, which are often covered with scales that hinder the access of natural enemies (Beserra and Parra, 2004; Souza et al, 2008). The chorion hardness of Noctuidae eggs may further thwart first instar *C. externa* larvae, as they have weak mouthparts (López-Arroyo et al, 2000).

A lower survival rate has been reported for *C. externa* fed with *Toxoptera citricida* (Kirkaldy, 1907) (Hemiptera: Aphididae) (Pappas et al, 2008b). On the other hand, the high survival rate of *C. externa* fed with *A. kuehniella* eggs was comparable to rearing with *A. argillaceae*, suggesting that this prey has satisfactory nutritional properties (Souza et al, 2008). Eggs of *A. kuehniella* can be stored frozen, which reduces costs compared with fresh prey, but the freezing period reduces nutritive quality (Mohaghegh and Amir-Maafi, 2007). High larval mortality of *D. prasina* lacewings was also reported after they were fed with *Aphis nerii* (Boyer de Fonscolombe, 1841) (Hemiptera: Aphididae) (Pappas et al, 2007) or *E. kuehniella* (Pappas et al, 2008a, b). Food quality also affected the development and survival of Coccinellidae larva and Pentatomidae nymphs (Isikber and Copland, 2002; Kalushkov and Hodek, 2001, 2004).

A shorter pupal stage, described here in *C. externa* fed with newly hatched *S. frugiperda* larvae, is important for biological control programs, as earlier maturation can lead to a more rapid population increase (Aquad et al, 2003; Pappas et al, 2007, 2008a). The duration of larval and pupal stages of *C. externa* were longer with ad libitum provision of food and reduced transfer between tubes (Aquad et al, 2003). This suggests that limitation of prey availability and handling may reduce the duration of the larval and pupal stages of this predator.

Prey limitation and increased handling has been previously found to reduce the viability of *C. externa* pupae (Aquad et al, 2003). This suggests that the ad libitum availability of *S. frugiperda* in the same tube

could improve the developmental success of this predator. The lower survival of third instar *C. externa* larvae and duration of larval and pupal stages fed with newly hatched *S. frugiperda* larvae provided with limited prey and daily tube changes suggests that limited prey and handling are not optimal condition for this predator (Aquad et al, 2003).

Although first instar *C. externa* larvae perforated the egg-shell of *S. frugiperda*, this prey was not suitable for development. This has been observed previously for this predator fed with *S. frugiperda* (Aquad et al, 2003). Similarly, development was compromised in the ladybeetle predator *Stethorus punctillum* (Weize, 1891) (Coleoptera: Coccinellidae) fed with *Tetranychus urticae* (Koch, 1836) (Acari: Tetranychidae) (Ragkou et al, 2004), and the lacewing *D. prasina* fed with *A. nerii* (Pappas et al, 2007). These results suggest that the survival of *C. externa* may be lower in the corn crop at the beginning of the infestation by *S. frugiperda* (Castillejos et al, 2000; Hoballah et al, 2004), a period during which eggs and adults of this pest are present (Matos Neto et al, 2004, 2005). However, studies involving the association of these insects, especially in the field, should be conducted.

The higher consumption (number of eggs or larvae) of *A. kuehniella* eggs by *C. externa* during the early and final instars can be attributed to the growth of this Chrysopidae and its increasing food necessity. This was also observed for *C. externa* fed with *B. tabaci* biotype B, *M. persicae*, *A. kuehniella*, or *S. frugiperda* (Silva et al, 2004; Barbosa et al, 2008). Prey density can also affect consumption, as was reported for *Podisus nigrispinus* (Dallas, 1851) (Heteroptera: Pentatomidae) feed with *S. frugiperda* larvae (Zanuncio et al, 2008). Higher consumption of prey provides females with great body mass, which correlates with higher fecundity (Zanuncio et al, 2002, 2005; Lemos et al, 2009). An example is seen for *C. sanguinea*, which is heavier and has higher fecundity when fed with *T. citricida* compared with a diet of *Aphis spiraeicola* (Patch, 1914) (Hemiptera: Aphididae) (Michaud, 2000). This is important because body weight indicates the amount of nutrients stored, which can affect mating, dispersion, flight and fecundity in insects (Omkar et al, 2006).

Table 4 - Weight of prey consumed (mean \pm standard error of mean) of the developmental stages (DS) (first instar – FI, second instar – SI, third instar – TI, and larval – L) of *Chrysoperla externa* (Neuroptera: Chrysopidae) fed with newly laid (T1) or one-day-old *Spodoptera frugiperda* (Lepidoptera: Noctuidae) eggs (T2); newly hatched (T3), one- (T4) or two-day-old (T5) *S. frugiperda* larvae; or one-day-old *Anagasta kuehniella* (Lepidoptera: Pyralidae) eggs (T6).

DS	T1	T2	T3	T4	T5	T6
	Consumption (weight of prey consumed)					
FI	1.93 \pm 0.09 ^b	2.07 \pm 0.22 ^b	3.44 \pm 0.23 ^b	10.84 \pm 0.73 ^a	6.15 \pm 0.74 ^a	1.74 \pm 0.03 ^b
SI	2.20 \pm 0.24 ^b	3.15 \pm 0.89 ^b	5.78 \pm 1.38 ^a	-	-	5.06 \pm 0.10 ^a
TI	8.16 \pm 1.09 ^b	11.03 \pm 0.63 ^a	8.50 \pm 1.48 ^b	-	-	12.90 \pm 0.11 ^a
L	12.30 \pm 1.43 ^b	16.26 \pm 1.03 ^a	17.74 \pm 3.10 ^a	-	-	19.71 \pm 0.24 ^a
CV	4.81%					

means followed by the same letter per line do not differ by the test of Tukey ($P < 0.05$); CV= Coefficient of variation

The higher consumption (number of eggs or larvae) of *A. kuehniella* eggs by *C. externa* may be due to its smaller size and weight and, consequently, the need of Chrysopidae to eat large numbers. The weight of 40 *A. kuehniella* eggs (1.56 mg) was less than 40 newly laid (2.03 mg) or 40 one-day-old (2.60 mg) eggs, 40 newly hatched (1.93 mg) or 40 one- (5.06 mg), or 40 two-day-old (5.70 mg) *S. frugiperda* larvae. This was observed for *C. externa* and *C. cubana* fed with Pyralidae eggs (Souza et al, 2008). However, the higher consumption (number of eggs or larvae) of *A. kuehniella* eggs by *D. prasina* than of *Toxoptera* sp. (Hemiptera: Aphididae) eggs and/or *Pinnaspis* sp. suggests that this prey is suitable for this predator (Pappas et al, 2008b). Moreover, the similar prey consumption (number of eggs or larvae) of *Diatraea saccharalis* (F., 1794) (Lepidoptera: Pyralidae), *Sitotroga cerealella* (Olivier, 1819) (Lepidoptera: Gelechiidae) and *A. kuehniella* by *Ceraeochrysa cincta* (Schneider, 1851) (Neuroptera: Chrysopidae) larvae (Pappas et al, 2007) suggests that species of Chrysopidae may have different food requirements.

In conclusion, newly laid eggs, one-day-old eggs or newly hatched larvae of *S. frugiperda* can be used as prey for *C. externa*, but the development of this predator was better with *A. kuehniella* eggs, suggesting that the latter may be better suited for mass production.

Acknowledgements

To “Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)”, “Fundação de Amparo a Pesquisa do Estado de Minas Gerais (FAPEMIG)” and “Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES)” for financial support. To “Paper Check” and “Asia Science” for correction and editing this manuscript.

References

Alcantra E, Carvalho CF, Santos TM, Souza B, Santa-Cecília LVC, 2008. Biological aspects and predatory capacity of *Ceraeochrysa cubana* (Hagen, 1861) (Neuroptera: Chrysopidae) fed on *Aphis*

gossypii Glover, 1877 (Hemiptera: Aphididae) under different temperatures. Ciênc Agrotec 32: 1047-1054

Auad AM, Carvalho CF, Souza B, Barbosa LR, 2003. Biological aspects of *Chrysoperla externa* (Hagen, 1861) (Neuroptera: Chrysopidae) fed on eggs and larvae of *Spodoptera frugiperda* (JE Smith, 1797) (Lepidoptera: Noctuidae). Rev Bras Milho Sorgo 2: 106-111

Barbosa LR, Carvalho CF, Souza B, Auad AM, 2008. Efficiency of *Chrysoperla externa* (Hagen, 1861) (Neuroptera: Chrysopidae) in the *Myzus persicae* (Sulzer, 1776) (Hemiptera: Aphididae) population reduction in sweet pepper (*Capsicum annum* L). Ciênc Agrotec 32: 1113-1119

Beserra EB, Parra JRP, 2004. Biology and parasitism of *Trichogramma atopovirilia* Oatman & Platner and *Trichogramma pretiosum* Riley (Hymenoptera, Trichogrammatidae) on eggs of *Spodoptera frugiperda* (JE Smith) (Lepidoptera, Noctuidae). Rev Bras Entomol 48: 119-126

Brekvens N, Bonte J, Berkvens D, Deforce K, Tirry L, De Clercq P, 2008. Pollen as an alternative food for *Harmonia axyridis*. Biol Control 53: 201-210

Castillejos V, García L, Cisneros J, Goulson D, Cave RD, Caballero P, Williams T, 2001. The potential of *Chrysoperla rufilabris* and *Doru taeniatum* as agents for dispersal of *Spodoptera frugiperda* nucleopolyhedrovirus in maize. Entomol Exp Appl 98: 353-359

Cole PG, Home PA, 2006. The impact of aphicide drenches on *Micromus tasmaniae* Walker (Neuroptera: Hemerobiidae) and the implications for pest control in lettuce crops. Aust J Entomol 45: 244-248

Figueiredo MLC, Dias AMPM, Cruz I, 2006. Relationship between fall armyworm and their natural biological control agents in the maize crop. Pesqui Agropecu Bras 41: 1693-1698

Gao F, Liu XH, Ge F, 2007. Energy budgets of the Chinese green lacewing (Neuroptera: Chrysopidae) and its potential for biological control of the cotton aphid (Homoptera: Aphididae). Insect Sci 14: 497-502

- Hagerty AM, Kilpatrick AL, Turnipseed SG, Sullivan MJ, Bridges WC, 2005. Predaceous arthropods and lepidopteran pests on conventional, Bollgard, and Bollgard II cotton under untreated and disrupted conditions. *Environ Entomol* 34: 105-114
- Hoballah ME, Degen T, Bergvinson D, Savidan A, Tamò C, Turlings TCJ, 2004. Occurrence and direct control potential of parasitoids and predators of the fall armyworm (Lepidoptera: Noctuidae) on maize in the subtropical lowlands of Mexico. *Agr Forest Entomol* 6: 83-88
- Hogervorst PAM, Wäckers FL, Carette AC, Romeis J, 2008. The importance of honeydew as food for larvae of *Chrysoperla carnea* in the presence of aphids. *J Appl Entomol* 132: 18-25
- Isikber AA, Copland MJW, 2002. Effects of various aphid foods on *Cycloneda sanguinea*. *Entomol Exp Appl* 102: 93-97
- Kalushkov AA, Hodek I, 2001. New essential aphid prey for *Anatis ocellata* and *Calvia quatuordecimgottata* (Coleoptera: Coccinellidae). *Biocontrol Sci Techn* 11: 35-39
- Kalushkov AA, Hodek I, 2004. The effects of thirteen species of aphids on some life history parameters of the ladybird *Coccinella septempunctata*. *Bio-Control* 49: 1984-1992
- Kovanci OB, Kovanci B, Gencer NS, 2007. Species composition, seasonal dynamics and numerical responses of arthropod predators in organic strawberry fields. *Biocontrol Sci Techn* 17: 457-472
- Kunkel BA, Cottrell TE, 2007. Oviposition response of green lacewings (Neuroptera: Chrysopidae) to aphids (Hemiptera: Aphididae) and potential attractants on pecan. *Environ Entomol* 36: 577-583
- Lawo NC, Romeis J, 2008. Assessing the utilization of a carbohydrate food source and the impact of insecticidal proteins on larvae of the green lacewing, *Chrysoperla carnea*. *Biol Control* 44: 389-398
- Lemos WP, Serrão JE, Zanuncio JC, Lacerda MC, Zanuncio TV, Ribeiro RC, 2009. Body weight and protein content in the haemolymph of females of the zoophytophagous predator *Brontocoris tabidus* (Heteroptera: Pentatomidae) with different diets and ages. *J Plant Dis Protect* 116: 218-222
- López-Arroyo JI, Tauber CA, Tauber M, 2000. Storage of lacewing eggs: post-storage hatching and quality of subsequent larvae and adults. *Biol Control* 18: 165-171
- Matos Neto FC, Cruz I, Zanuncio JC, Silva CHO, Picanço MC, 2004. Parasitism by *Campoletis flavicincta* on *Spodoptera frugiperda* in corn. *Pesqui Agropecu Bras* 39: 1077-1081
- Matos Neto FC, Zanuncio JC, Cruz I, Guedes RNC, Picanço MC, 2005. Progeny production and parasitism by *Campoletis flavicincta* (Hym. Ichneumonidae) as affected by female ageing. *Biol Agric Hort* 22: 369-378
- Michaud JP, 2000. Development and reproduction of ladybeetles (Coleoptera: Coccinellidae) on the citrus aphids *Aphis spiraecola* Patch and *Toxoptera citricida* (Kirkald) (Homoptera: Aphididae). *Biol Control* 18: 287-297
- Michaud JP, 2005. On the assessment of prey suitability in aphidophagous Coccinellidae. *Eur J Entomol* 102: 385-390
- Mohaghegh J, Amir-Maafi M, 2007. Reproduction of the predatory stinkbug *Andrallus spinidens* (F.) (Heteroptera: Pentatomidae) on live and frozen prey. *Appl Entomol Zool* 42: 15-20
- Nakahira K, Nakahara R, Arakawa R, 2005. Effect of temperature on development, survival, and adult body size of two green lacewings, *Mallada desjardinsi* and *Chrysoperla nipponensis* (Neuroptera: Chrysopidae). *Appl Entomol Zool* 40: 615-620
- Omkar MG, Singh SK, 2006. Optimal number of matings in two aphidophagous ladybirds. *Ecol Entomol* 31: 1-4
- Pappas ML, Broufas GD, Koveos DS, 2007. Effects of various prey species on development, survival and reproduction of the predatory lacewing *Dichochrysa prasina* (Neuroptera: Chrysopidae). *Biol Control* 43: 163-170
- Pappas ML, Broufas GD, Koveos DS, 2008a. Effect of temperature on survival, development and reproduction of the predatory lacewing *Dichochrysa prasina* (Neuroptera: Chrysopidae) reared on *Ephestia kuehniella* eggs (Lepidoptera: Pyralidae). *Biol Control* 45: 396-403
- Pappas ML, Broufas GD, Koveos DS, 2008b. Effect of relative humidity on development, survival and reproduction of the predatory lacewing *Dichochrysa prasina* (Neuroptera: Chrysopidae). *Biol Control* 46: 234-241
- Ragkou VS, Athanassiou CG, Kavallieratos NG, Tomanovic Z, 2004. Daily consumption and predation rate of different *Stethorus punctillum* instars feeding on *Tetranychus urticae*. *Phytoparasitica* 32: 154-159
- Russel DF, 1989. MSTAT-C Statistical Package Program ver. 2.1. Michigan State University, East Lansing
- Sattar M, Fatima B, Ahmed N, Abro GH, 2007. Development of larval artificial diet of *Chrysoperla carnea* (Stephens) (Neuroptera: Chrysopidae). *Pak J Zool* 39: 103-107
- Sattar M, Abro GH, 2009. Comparative effect of natural and artificial larval diets on biology of *Chrysoperla carnea* (Stephens) (Neuroptera: Chrysopidae). *Pak J Zool* 41: 335-339
- Silva CG, Souza B, Auad AM, Bonani JP, Torres LC, Carvalho CF, Ecole CC, 2004. Development of immature stages of *Chrysoperla externa* fed on nymphs of *Bemisia tabaci* biotype B reared on three hosts. *Pesqui Agropecu Bras* 39: 1065-1070
- Silva RB, Zanuncio JC, Serrão JE, Lima ER, Figueiredo MLC, Cruz I, 2009. Suitability of different arti-

- ficial diets for development and survival of stages of predaceous ladybird beetle *Eriopis connexa* (Coleoptera: Coccinellidae). *Phytoparasitica* 37: 115-123
- Souza B, Costa RIF, Tanque RL, Oliveira OS, Santos FA, 2008. Predation among *Chrysoperla externa* (Hagen, 1861) and *Ceraeochrysa cubana* (Hagen, 1861) (Neuroptera: Chrysopidae) larvae under laboratory conditions. *Ciênc Agrotec* 32: 712-716
- Tavares WS, Cruz I, Petacci F, Assis Júnior SL, Freitas SS, Zanuncio JC, Serrão JE, 2009. Potential use of Asteraceae extracts to control *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and selectivity to their parasitoids *Trichogramma pretiosum* (Hymenoptera: Trichogrammatidae) and *Telenomus remus* (Hymenoptera: Scelionidae). *Ind Crop Prod* 30: 384-388
- Tavares WS, Cruz I, Fonseca FG, Gouveia NL, Serrão JE, Zanuncio JC, 2010a. Deleterious activity of natural products on postures of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and *Diatraea saccharalis* (Lepidoptera: Pyralidae). *Z Naturforsch C* 65: 412-418
- Tavares WS, Costa MA, Cruz I, Silveira RD, Serrão JE, Zanuncio JC, 2010b. Selective effects of natural and synthetic insecticides on mortality of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) and its predator *Eriopis connexa* (Coleoptera: Coccinellidae). *J Environ Sci Heal B* 45: 557-561
- Vandekerckhove B, Van Baal E, Bolckmans K, De Clercq P, 2006. Effect of diet and mating status on ovarian development and oviposition in the polyphagous predator *Macrolophus caliginosus* (Heteroptera: Miridae). *Biol Control* 39: 532-538
- Wade MR, Zalucki MP, Wratten SD, Robinson KA, 2008. Conservation biological control of arthropods using artificial food sprays: current status and future challenges. *Biol Control* 45: 185-199
- Zaki FN, Gesraha MA, 2001. Production of the green lacewing *Chrysoperla caranea* (Steph) (Neuropt: Chrysopidae) reared on semi-artificial diet based on the algae, *Chlorella vulgaris*. *J Appl Entomol* 125: 97-98
- Zanuncio JC, Molina-Rugama AJ, Santos GP, Ramalho FS, 2002. Effect of body weight on fecundity and longevity of the stinkbug predator *Podisus rostralis*. *Pesqui Agropecu Bras* 37: 1225-1230
- Zanuncio JC, Beserra EB, Molina-Rugama AJ, Zanuncio TV, Pinon TBM, Maffia VP, 2005. Reproduction and longevity of *Supputius cincticeps* (Het.: Pentatomidae) fed with larvae of *Zophobas confusa*, *Tenebrio molitor* (Col: Tenebrionidae) or *Musca domestica* (Dip: Muscidae). *Braz Arch Biol Techn.*48: 771-777
- Zanuncio JC, Silva CA, Lima ER, Pereira FF, Ramalho FD, Serrão JE, 2008. Predation rate of *Spodoptera frugiperda* (Lepidoptera: Noctuidae) larvae with and without defense by *Podisus nigrispinus* (Heteroptera: Pentatomidae). *Braz Arch Biol Techn* 51: 121-125

