

Available online at <http://jddtonline.info>

REVIEW ARTICLE

PHYTOCONSTITUENTS AND THERAPEUTIC POTENTIALS OF *DATURA STRAMONIUM* LINN

*Das Sanjita, Puneet kumar, Basu S.P.

Noida Institute of Engineering and Technology, 19, Knowledge Park-2, Institutional area, Phase-2, Greater Noida, U.P., INDIA-201306

*Corresponding Author's Email: Sanjita8@yahoo.co.in, Phone: 9818223359

Received 01 April 2012; Revised 09 April 2012; Accepted 01 May 2012, Available online 15 May 2012

ABSTRACT

Datura stramonium Linn is one of the widely well known folklore medicinal herb. It is a wild growing flowering plant and was investigated as a local source for tropane alkaloids which contain a methylated nitrogen atom (N-CH₃) and include the anti-cholinergic drugs atropine, and scopolamine. This plant has contributed various pharmacological actions in the scientific field of Indian systems of medicines like analgesic, antiasthmatic activities. The present paper will give exclusive information on the chemical constituents and mainly psychopharmacological activities of this plant. *Datura stramonium* has been scientifically proven to have alkaloids, tannins, carbohydrates, proteins. This study highlights the pharmacological activities of *Datura stramonium*, which may be due to the presence of its scientifically proven chemical constituents.

Keywords: *Datura stramonium*, phytochemistry, ethnopharmacology, pharmacology.

INTRODUCTION

From ancient civilization it was traditionally used for religious visionary purposes throughout the world and used for witchcraft in medieval Europe. The god Lord Shiva was known to smoke Cannabis and *Datura*. People still provide the small thorn apple during festivals and special days as offerings in Shiva icons at temples. An extract made from the leaves is taken orally for asthma and sinus infections and stripped bark are applied externally to swellings, burns and ulcers¹. The incidence of *D. stramonium* poisoning is sporadic with a cluster of poisoning cases in the 1990s and 2000s, the United States media contained stories occurring mostly among adolescents and young adults dying or becoming seriously ill from ingesting. Some medicinal uses of the plant are its anti-inflammatory property of all part of the plants², stimulation of the central nervous system (CNS)³, respiratory decongestion⁴, treatment of dental and skin infections, alopecia^{5,6} and also in the treatment of toothache⁷. It is a hallucinogenic plant that causes serious poisoning. Consumption of any part of the plant may result in a severe anticholinergic reaction that may lead to toxicity and occasionally cause diagnostic difficulties⁸. Cases of poisoning have been reported after eating the berries; death may occur from heart failure after ingesting 125 seeds⁹, because the seeds contain the highest concentration and has a rapid onset of action¹⁰ and thus may be potentially useful as an alternative to atropine for the treatment of the muscarinic symptoms of organophosphate toxicity and some of central anticholinergic effects¹¹. The wide distribution, the strong toxicity and the potential for occurrence in foodstuffs are responsible for the numerous incidents in humans¹²⁻¹⁷.

Vernacular Names:

Sanskrit	: Dhastura
Bengali	: Dhattura
English	: Angel's Trumpet
Gujrati	: Unmatta
Hindi	: Dattura
Marathi	: Kanaka
Malayalam	: Marau mmam
Tamil	: Ummatta

Taxonomical Classification:

Kingdom	: Plantae
Subkingdom	: Tracheobionta
Superdivision	: Spermatophyta
Division	: Magnoliophyta
Class	: Magnoliopsida
Subclass	: Asteridae
Order	: Solanales
Family	: Solanaceae
Genus	: <i>Datura L</i>
Species	: <i>Stramonium</i>

Botanical description

Jimson weed (*Datura stramonium*) is known to contain highly toxic tropane alkaloids, including the pharmacologically active compounds atropine and scopolamine. *D. stramonium* is a bushy, smooth, foetid annual on rich soil it may attain a height of even 6 feet. The branching stems are spreading and leafy, stout, erect, smooth, a pale yellowish green in color, branching repeatedly in a

cardiovascular collapse²⁴. Its anticholinergic compounds are likely to produce delirium and stupor but rarely cause deep coma. The ethanolic extracts obtained from both leaf and seed in the thorn apple were investigated for acaricidal, repellent and oviposition deterrent properties against adult two-spotted spider mites under laboratory conditions. Leaf & seed extracts, which were applied in 167,250 and 145,750 mg/l concentrations, respectively caused 98% and 25 % mortality among spider mite adults after 48 h²⁵. Exposure of the foetus to this plant when a mother uses it for asthma, will cause a continuous release of Ach, resulting in the desensitizing of nicotinic receptors, this could ultimately result in permanent damage to the foetus²⁶. The main effects of jimsonweed seeds were: decreased body weight gain, serum alkaline phosphatase and blood urea nitrogen. Female rats showed more marked responses to jimson weed seed than did males. In addition to the effects seen in both sexes, the females developed decreased serum total protein and cholesterol, and increase serum glutamic, pyruvic transaminase and chloride, red blood cell count, haemoglobin concentration and packed red cell volume²⁷. An extract prepared from the seeds of the DS possess activity typical of a protein haemagglutinin or lectin. The extract is capable of agglutinating erythrocytes from several species, and is non-specific with regard to human ABO blood groups. Laboratory monitoring of changes in some blood parameters in horses intoxicated with jimsonweed was carried out. It was established that the intoxication was accompanied by hyperchromaemic, erythrocytosis, leukocytosis, neutrophilia and regenerative shift, lymphocytopenia, aneosinophilia, increased haematocrit values and low erythrocyte sedimentation rat²⁸. All parts of the plant are toxic but the highest amount of the alkaloids is contained in ripe seeds²⁹. They act as competitive antagonist of acetylcholine at peripheral and central muscarinic receptor sites. Poisoning results in widespread paralysis of parasympathetic innervated

organs³⁰. Datura aqueous leaf extract induces cytotoxicity & oxidative stress in human cancer cell lines. Severe toxicity has been associated with coma and seizures, although death is rare³¹.

Anticholinergic activity of *D. stramonium*: The alkaloids found in *D. stramonium*, are organic esters used clinically as anticholinergic agents. Jimson weed (also called Jamestown weed, thornapple and stinkweed) has been reported as a drug of abuse³²⁻³³ and has been involved in the accidental poisoning of humans and animals³⁴⁻³⁷. Symptoms of acute jimson weed poisoning included dryness of the mouth and extreme thirst, dryness of the skin, pupil dilation and impaired vision, urinary retention, rapid heartbeat, confusion, restlessness, hallucinations, and loss of consciousness. Chronic and subchronic toxic effects of concern could be more subtle than those observed in acute poisonings of man, which have come to the attention of the medical profession, or of domestic animals that have consumed the plant as a source of food.

CONCLUSION

Present study gives a broad information about the bioactive constituents, ethnopharmacology along with the scientifically claimed medicinal uses of *D. stramonium*. Several alkaloids, carbohydrates, fat, proteins and tannins have been reported to be present in different parts of *D. stramonium*. Plant shows various types of activities such as analgesic and antiasthmatic activity which may be due to the presence of the investigated active chemical constituents. In future to extend this study, elaborated psychopharmacological studies will be done as the plant contains alkaloids. Furthermore other parts of the plants are to be exploited for further pharmacological investigations along with the establishment of mechanism and chemical constituents responsible for the respective therapeutic potential.

REFERENCES:

- Asprey GF and Thornton P, Medicinal Plants of Jamaica, W. I. Med. J. 1953, 3,17-41.
- Spring MA, Ethnopharmacologic analysis of medicinal plants used by Laotian Hmong refugees in Minnesota, J. Ethnopharmacol. 1989, 26, 65-91.
- Guharov SR, Barajas M, Intense stimulant effect: atropine intoxication from the ingestion and smoking of Jimson weed (*Datura stramonium*), Vet. Toxicol. 1991, 33, 588-589.
- Zagari A, Medicinal plants, Vol. 3, 5th ed. Tehran University Publication, Tehran, Iran. 1992, pp. 889.
- De Foe V and Senatore F, Medicinal plants and phytotherapy in the Amal Fitan Cost, Salerno province Campania, Southern Italy, J.Ethnopharm, 1993, 39, 39-51
- John D, One hundred useful drugs of the Kani tribes of Trivandrum forest divisions, Kerala, India, Inter. J. crude drug Res. 1984, 22, 17-39.
- Abebe W, A survey of prescriptions used in traditional medicine in Gondar region, North-West Ethiopia: general pharmaceutical practice, J. Ethnopharm, 1986, 18, 147-165.
- Diker D, Markovitz D, Rothman M. and Sendovski U, Coma as a presenting sign of *Datura stramonium* seed tea poisoning, European J.of Internal Med. 2007, 34(2), 336-338.
- Nehaul, BBG, Datura poisoning in British Guiana. Report of four cases, W. I. Med. J. 1955, 4, 57-59.
- Chang SS, Wu ML, Deng JF, Lee CC, Chin TF and Liao SJ, Poisoning by Datura leaves used as edible wild vegetables, Vet. Human Toxicol. 1999, 41, 242-245.
- Theodore CB, Jasan C, Dallas B, Melanie O, Jimson Weed Extract as a Protective Agent in Severe Organophosphate Toxicity. Acad Emerg Med. 2004, 11(4), 335-338.
- Roblot F, Montaz L, Delcoustal M, Gaboriau E, Chavagnat JJ, Morichaud G, Pourrat O, Scepti M and Patte D, *Datura stramonium* poisoning: The diagnosis is clinical, treatment is symptomatic, Revue de Med. Inteme. 1995, 16, 187-190.
- Arouko H, Matray D, Braganca C, Mpaka P, Chinello L, Castaing F, Bartou C and Poisot D, Voluntary poisoning by ingestion of *Datura stramonium*. Another cause of hospitalisation in youth seeking strong sensations, Annual de Med. Inteme. 2003, 154, 46-50.
- Boumba A, Mitselou A and Vougiouklakis T, Fatal poisoning from ingestion of *Datura stramonium* seeds, Vet. Human Toxicol. 2005, 46, 81-82.
- Ertekin V, Selimoglu MA and Altinkaynak S, A combination of unusual presentations of *Datura stramonium* intoxication in a child: Rhabdomyolysis and fulminant hepatitis, J. of Emergency Med. 2005, 28, 227- 228.
- Oberndorfer S, Grisold W, Hinterholzer G and Rosner M, Coma with focal neurological signs caused by *Datura stramonium* intoxication in a young man, J. N. Neuro. Psy. 2002, 73, 458- 459.

17. Shaikh. AL and Sablay, Z M, Hallucinogenic plant poisoning in children, *Saudi Med.J.* 2005, 26, 118- 121.
18. Romeike A, Beiträge zur chemischen Physiologie der mydriatisch wirkenden Solanaceen- Alkaloide, *Pharmazie.* 1953, 8, 668-674 and 729-747.
19. Phillipson JD and Handa SS, N-oxides of hyosciamine and hyoscyne in the Solanaceae, *Phytochem.* 1975, 14: 2683-2690.
20. Bucher E and Meszaros L, Stechapfelsamen (*Datura spec.*) und deren Alkaloide in Futtermitteln. *Kraftfutter* 1989, 3(89), 76-82.
21. Bucher E, Tischler J and Ludsteck M, HPLC-Bestimmung der Alkaloide (Scopolamin und Hyoscyamin) von Stechapfelsamen (*Datura spec.*) in Futtermittel., *Landwirtschaftliche Forschung.* 1989, 42, 293-309.
22. Berkov S, Zayed R and Doncheva T, Alkaloid patterns in some varieties of *Datura stramonium*, *Fitoterapia.* 2006, 77 (3), 179-182.
23. Oseni OA, Olarinoye CO and Amoo IA, Studies on chemical compositions and functional properties of thorn apple (*Datura stramonium* L) Solanaceae, *African J. Food Sc.* 2011, 5 (2), 40 – 44.
24. Alberto K, Claudia S, Ludmilla K and Arnon B, Toxic Delirium due to *Datura stramonium*, *Israel Med. Asso. J.* 2001, 3, 538-539.
25. Kurnal NA, Yalcin SCC, Acaricidal, repellent and oviposition deterrent activities of *Datura stramonium* L. against adult *Tetranychus urticae* (Koch), *JPest Sci.* 2009, 14, 54-57.
26. Pretorius E and Marx J, *Datura stramonium* in asthma treatment and possible effects on prenatal development *Environ, Toxicol. and Pham.* 2006, 21(3), 331-337.
27. Dugan GM, Gumbmann MR and Friedman M, Toxicological evaluation of jimson weed (*Datura stramonium*) seed, *Food and Chem. Toxicol.* 1989, 27, 501-510.
28. Binev R, Valchev I. and Nikolov J, Haematological studies on Jimson weed (*Datura stramonium*) intoxication in horses, *Trakia J. of Sci.* 2006, 4(1), 43-48.
29. DeFrates LJ, Hoehns JD, Sakornbut EL, Glascock DG and Tew AR, Antimuscarinic intoxication resulting from the ingestion of moonflower seeds, *The Annals of Pham.* 2001, 39, 173- 176.
30. Friedman M and Levin E, Composition of jimson weed (*Datura stramonium*) seeds, *J. of Agri. and Food Chem.* 1989 37, 998-1005.
31. Dewitt MS, Swain R, Gibson LB, The dangers of jimson weed and its abuse by teenagers in the Kanawha Valley of West Virginia, *WV Med J,* 1997, 93(4), 182-185.
32. Klein-schwartz W and Oderda GM, Jimsonweed intoxication in adolescents and young adults, *Am. J. Dis. Child.* 1984, 138, 737-739.
33. Ogrady TC, Brown J and Jacamo J, Outbreak of jimson weed abuse among Marine crops personnel at Camp Pendleton, *Milit. Med.* 1983, 148, 732-734.
34. Clark WL, Jemigan RH and Erwin JW, Epidemiological report jimson weed poisoning--Tennes-see. *Morbidity and Mortality Weekly Report,* US Department of Health, Education and Welfare, 1964, pp. 126-127.
35. Taha SA and Mahdi AW, *Datura* intoxication in Riyadh, *Trans. R. Soc. trop Med. Hgy.* 1984, 78, 134-135.
36. Urich RW, Bowerman DL, Levisky JA and Pflug JL, *Datura stramonium*: a fatal poisoning, *J. Forens Sci.* 1982, 27, 948-954.
37. Weintraub S, Stramonium poisoning, *Postgrad. Med.* 1960, 28: 364-367.
38. Feenaughty DA, Atropine poisoning: jimsonweed, *J. emerg. Nurs.* 1982, 8, 139-141.
39. Lampe KF and McCann MA, *AMA Handbook of poisonous and Injurious Plants,* American medical Association, Chicago, 1985, pp. 70-72.
40. Thabet H, Brahm N, Amamou M, BenSalah N, Hedhill and Yacoub M, *Datura stramonium* poisonings in humans, *Vet. Human Toxicol.* 1999, 41, 320-321.