

Available online at <http://jddtonline.info>

## REVIEW ARTICLE

MEDICO ETHNOBOTANICAL PERSPECTIVES OF JYOTISMATI (*CELASTRUS PANICULATUS WILLD*): A HERBAL TRANQUILIZERDebnath Monojit\*<sup>1</sup>, Pushpan Reshmi<sup>2</sup>, Kumari Harshita<sup>2</sup>, Nishteswar K<sup>3</sup><sup>1</sup>Scholar M.Pharm(Ayur),IPGT & RA, Gujarat Ayurved University, Jamnagar,361008.<sup>2</sup>Scholar PhD,IPGT & RA, Gujarat Ayurved University, Jamnagar,361008.<sup>3</sup>Professor & HOD,Dravyaguna,IPGT & RA, Gujarat Ayurved University, Jamnagar,361008.\*Corresponding Author's Email: [monodebnath@gmail.com](mailto:monodebnath@gmail.com), Phone:+91 8460656811

Received 14 March 2012; Revised 23 April 2012; Accepted 01 May 2012, Available online 15 May 2012

## ABSTRACT:

The primary objectives of the present study is to find out and documentation of various medico ethnobotanical claims of jyotismati from different tribal population throughout the India and to make a valuable discussion on the modern evaluation of the drug. Jyotismati (*Celastrus paniculatus* Willd.) is a woody climber of vedic lore. In several studies seed oil was screened for its sedative and tranquillizing properties. But the tribal claims of other parts of jyotismati are yet to be studied. The plant jyotismati is used throughout the tribal population of India for wound healing, cough, insomnia, opium poisoning. The details of medico ethnobotanical aspect of the plant jyotismati and the recent researches carrying out on the plant clearly indicate that jyotismati plays a key role in the healthcare system of India. Modern researches are revalidating the tribal uses of jyotismati. The indiscriminate use of seed and fruits may lead for its inclusion to the endangered list. So in search of its substitution the leaf and bark should be thoroughly studied experimentally and clinically. Now research regarding the other parts of this plant (stem bark, root, leaf ) is required to establish the useful therapeutic profile of the whole plant and to prove the ethnobotanical claims. Although a vivid scientific enquiry and clinical studies have to be encouraged and the adverse reactions are to be recorded.

**Keywords:** Jyotismati, Classical use, Ethnobotanical use, Modern evaluation.

## INTRODUCTION:

About 7500-8000 species of plants are estimated to be used for human and veterinary health care in the century. About 3000 plants species are reported to be used in the codified Indian system like Ayurveda(900 species), Siddha(800 species) , Unani(700 species), and Amchi (300 species).

Tribals make use of surrounding flora and fauna in the management of various ailments. Acharyas of Ayurveda also advocated to obtain the knowledge of herb from forest dwellers, cowherds etc. A thorough scan of medico ethnobotanical information indicates that in tribal medicine classical drugs are being used for a different condition along with non classical drugs in their tribal medical practises. Jyotismati is one of such classical drugs which is being used by tribals for different conditions not indicated by ayurvedic classics. Bark is reported to have abortifacient, wound healing, sedative and bronchodialator activities. Root is used as an antidote for snake bite poisoning. Leaves are proved to be emmenagogue and the leaf sap is employed as a good antidote for opium poisoning. Root and Leaf paste are used in headache.<sup>1</sup>

## BOTANICAL DESCRIPTIONS OF JYOTISMATI:

It is a large, woody scrambling or climbing Polygamodioecious type of shrub, which with assistance from a nearby tree climbs up to over 10 m. It occurs almost throughout India ascending to an altitude of 1800 m in the subtropical Himalayas. Main stem and thick branches have

yellowish bark. Leaves acuminate or acute and glabrous. Flowers are small, unisexual, yellowish green, fragrant, borne interterminal , pendulous panicles. Capsules are three celled, globose, 6-10 mm across,wrinkled and greenish brown in colour, when mature. Capsule contains 3-6 ovoid or ellipsoid, reddish brown in colour, encased in complete arillus of scarlet colours. Plant flowers and fruits at various times throughout the year.

## CLASSICAL USES OF JYOTISMATI:

1) **Udara roga**(Abdominal disease)- Intake of jyotismati seed oil with sarja kshara,hingu and milk will inhibit udara roga (Susruta). Intake of seed oil with milk for 8 days inhibits sannipata udara roga (Vangasena).

2) **Nastartava** or anartava(Amenorrhoea)- Leaf of jyotismati should be triturated with kanji(Sour gruel) to regularize the menstruation (Chakradutta).

3) **Sidhma**- Jyotismati oil processed with water of apamarga kshara on external application alleviates sidhma (Ashtangahridaya).

4) **Lutha Visha**(Spider Poisoning)- Jyotismati patra and Karkati patra svarasa is mixed and administered in spider poisoning (Vrinda Madhava).

5) **Urdhwajatrugata vikaras**- Siroroga, Apasmara and pinasa(Charaka).

## ETHNOMEDICAL USES OF JYOTISMATI:

Table: 1: Tribal claims of jyotis mati from different regions of India.

ACTION	PART USED	SOURCE(Tribal Pockets)
Abortifacient <sup>2,3</sup>	Bark	Sikkim, Bengal, Bihar, Orissa
Anaemia <sup>4</sup>	part not mentioned	Sikkim, Bengal, Bihar, Orissa
Antidote to snake bite <sup>5</sup>	Root	Uttarpradesh
Backache <sup>6</sup>	Seed	Jammu, Kashmir, Himachal Pradesh
Bodyache <sup>7,8</sup>	seed oil	Punjab, Hariyana, Rajasthan, Gujarat; Sikkim, Bengal, Bihar, Orissa
Bone fracture <sup>9</sup>	Bark	Punjab, Hariyana, Rajasthan, Gujarat
Bronchities <sup>10</sup>	Bark	Not specified
Carbuncle <sup>4</sup>	part not mentioned	Sikkim, Bengal, Bihar, Orissa
Cold <sup>11,12</sup>	Seed	Uttarpradesh
Colic <sup>4</sup>	part not mentioned	Sikkim, Bengal, Bihar, Orissa
Cough <sup>11,12</sup>	Seed	Uttarpradesh
Diarrhoea <sup>11,12</sup>	part not mentioned	Uttarpradesh
Digestive complaints <sup>13</sup>	Seed	Punjab, Hariyana, Rajasthan, Gujarat
Dysentery <sup>11,12</sup>	Fruit	Uttarpradesh
Eczema <sup>14</sup>	seed oil	Madhyapradesh, Maharashtra, Andhra Pradesh
Fever <sup>11,15,12</sup>	Seed	Uttarpradesh
Gastric complaints <sup>16</sup>	Bark	Sikkim, Bengal, Bihar, Orissa
Gout <sup>16,17</sup>	Seed	Sikkim, Bengal, Bihar, Orissa; Madhyapradesh, Maharashtra, Andhra Pradesh
Hair tonic <sup>18</sup>	Seed	Uttarpradesh
Headache <sup>19</sup>	leaf+root	Andaman and Nicobar group of Island
Leprosy <sup>15</sup>	Seed	Uttarpradesh
Memory <sup>17</sup>	Seed	Punjab, Hariyana, Rajasthan, Gujarat
Paralysis <sup>17</sup>	Seed	Punjab, Hariyana, Rajasthan, Gujarat
Rheumatism <sup>13,20</sup>	Seed	Punjab, Hariyana, Rajasthan, Gujarat; & aman and Nicobar group of Island
	seed oil	Sikkim, Bengal, Bihar, Orissa
Scabies <sup>21,8</sup>	Fruit	Sikkim, Bengal, Bihar, Orissa
Sex disease <sup>22</sup>	Root	Sikkim, Bengal, Bihar, Orissa
Skin disease <sup>21,23</sup>	Fruit	Sikkim, Bengal, Bihar, Orissa
	Seed	Andaman and Nicobar group of Island
Snake bite <sup>5</sup>	Root	Uttarpradesh
Sores <sup>4</sup>	part not specified	Sikkim, Bengal, Bihar, Orissa
Stimulant <sup>13,24</sup>	Seed	Punjab, hariyana, Rajasthan, Gujarat
Stomachache <sup>25</sup>	seed oil locally	Sikkim, Bengal, Bihar, Orissa
Swollen veins <sup>26,25</sup>	bark	Sikkim, Bengal, Bihar, Orissa
	seed oil	Sikkim, Bengal, Bihar, Orissa
Syphillis <sup>4</sup>	part not specified	Sikkim, Bengal, Bihar, Orissa
Tonic <sup>17</sup>	Seed	Punjab, Hariyana, Rajasthan, Gujarat
Wounds <sup>27</sup>	Bark	Andaman and Nicobar group of Island
	Seed <sup>8</sup>	Sikkim, Bengal, Bihar, Orissa
	seed+ leaf <sup>28</sup>	Uttarpradesh
Edible <sup>28</sup>	Fruit	Sikkim, Bengal, Bihar, Orissa
Vegetable <sup>29,30</sup>	flower, fruit	Andaman & Nicobar group of Island; Punjab, Hariyana, Rajasthan, Gujarat
Luminant <sup>31,32</sup>	seed oil	Sikkim, Bengal, Bihar, Orissa
Edible oil, Cooking medium, control epileptic bouts <sup>33</sup>	Seed oil	

## MODERN RESEARCH STUDIES ON *CELASTRUS PANICULATUS*:

- 1) Jyotismati as a Medhya rasayana have very good effect in case of mental depression.<sup>34</sup>
- 2) Jyotismati is having very potent activity against amvata.<sup>35</sup>
- 3) Tranquilizing activity of Malkangni oil was reported.<sup>36</sup>
- 4) Seed extract shows Anti-spermatogenic effect in rat.<sup>37,38</sup>
- 5) Root bark Chloroform extract showed significant antimalarial activity against *Plasmodium falciperum* in vitro.<sup>39</sup>
- 6) It showed significant CNS depressant effect and a clear synergism with pentobarbital.<sup>40</sup>
- 7) The effect of seed oil on learning and memory was found to be significant<sup>41</sup>. In another set of experiments the seed oil (Chronically administered) was demonstrated to selectively reverse the scopolamine –induced task performance deficit (Central muscarinic receptor blockade) in a navigational memory task in young adult rats<sup>42</sup>. Aqueous extract of the seeds showed cognition enhancing activity in the male wister rats on shuttle- box.<sup>43</sup>
- 8) Alcoholic (70%) extract of whole plant showed tranquilizing action in rats<sup>44</sup>. Alcoholic extract also displayed anti-inflammatory activity against carrageenan induced hind paw oedema and cotton palate induced granuloma in albino rats<sup>44</sup>. Both the alcoholic extract<sup>44</sup> as well as a pure sesquiterpinoid polyol ester (ex- seed oil)<sup>45</sup> showed antipyretic analgesic activities in standard animal studies. The drug clearly shows its protective effect in stress<sup>46</sup>.
- 9) Aqueous alcoholic (50 %) extract of seeds exhibited hypolipidaemic effect. Extract also showed atherosclerotic effect.<sup>47</sup>
- 10) Seed has been shown to have antimalarial activity<sup>48</sup>. Methanolic extract shows anti oxidant activity<sup>49</sup>.
- 11) A Clinical trial was taken on 30 patients of residual schizophrenia to assess the effect of Smritisagara rasa, an ayurvedic herbo-mineral preparation consisting *Celastrus paniculatus* as one of the ingredients. Administered in doses of 250 mg TDS with honey for three months. Clinical studies revealed that 11 out of 30 patients showed

significant improvement and approximately similar number showed moderate improvement.<sup>50</sup>

## CHEMICAL CONSTITUENTS:

The leaves contains alkaloids, a glycoside and colouring matter, whereas the oil extracted from seeds contain sterols, alkaloids and a bright colouring matter, Celapanin, Celapanigin, Celapagin, Celastrine and paniculatin are the some important alkaloids present in the seeds. The oil also contains sesquiterpene like dipalmitoyl glycerol and alkaloids also.<sup>51</sup>

## DISCUSSION:

Jyotismati is a plant being used from Vedic lore. Ethnobotanical studies claim the significant use of its root as antidote in snake bite and the use of its bark in bone fracture, bronchitis, abortion, gastric complaints and swollen veins. Leaf is used for sedative and wound healing activity in many regions. Modern research work evaluates the seed oil activity in favour of its ethnomedical claims.

The bark is abortifacient, depurative and a brain tonic. The leaves are emmenagogue and the leaf sap is a good antidote for opium poisoning. The seeds are acrid, bitter, thermogenic, emollient, stimulant, intellect promoting, digestive, laxative, emetic, expectorant, appetizer, aphrodisiac, cardiogenic, anti-inflammatory, diuretic, diaphoretic, febrifuge and tonic, abdominal disorders, leprosy, skin diseases, paralysis, asthma, leucoderma, cardiac debility, inflammation, nephropathy, amenorrhoea, dysmenorrhoea. Those modern evaluations prove the efficacy of this medicine in tribal life. So, a vivid scientific enquiry and clinical studies have to be encouraged and the adverse reactions are to be recorded.

## CONCLUSION:

Jyotismati is being used now a day mainly for its tranquilizing activity but several other uses are practised in the tribal populations. Modern researches are revalidating the tribal uses of jyotismati. The indiscriminate use of seed and fruits may lead for its inclusion to the endangered list. So in search of its substitution the leaf and bark should be thoroughly studied experimentally and clinically. Now research regarding the other parts of this plant (stem bark, root, leaf) is required to establish the useful therapeutic profile of the whole plant and to prove the ethnobotanical claims.

## REFERENCES:

- 1) Sudha Parimala GH, Shashidhar, Ch. Sridevi, V. Jyothi and R. Suthakaran, Anti-inflammatory activity of *Celastrus paniculatus* seeds, International Journal of PharmTech Research, 2009, 1(4), 1326-1329.
- 2) Tarafder CR. Ethnogaecology in relation to plants-II. Plants used in abortion. J. Econ. Tax. Bot. 1983, 4: 507-516.
- 3) Brothakur, S. K. Certain plants in the folklore and folkline of the karbis (Mikirs) of Assam. In S. K. Jain (Ed) Glimpses of Indian Ethnobotany. 1981, 170-181.
- 4) Jain SK, Tarafder CR. Medicinal plant lore of the Santals. A revival of P.O. Boddington's work. Econ. Bot. 1970, 24, 241-278.
- 5) Maheswari JK, Singh KK, Saha S. Ethnobotany of tribals of Mirzapur district, U.P. Nation. Bot. Res. Inst., Lucknow, 1986.
- 6) Vir Jee, Dar GH, Kachroo P. Bhat GM. Taxo-ethnobotanical studies of the rural areas in district Rajouri (Jammu). J. Econ. Tax. 1984, Bot. 5, 831-838.
- 7) Bedi SJ. Ethnobotany of the Ratan Mahal Hills, Gujarat, India. Econ Bot. 32, 1978, 278-284.
- 8) Pal DC, Srivastava JN. Preliminary notes of ethnobotany of Singbhum district, Bihar. Bull. Bot Surv. Nodia, 1976, 18:247-250.
- 9) Jain SK. Plants in Indian Medicine and Folklore associated with healing of bones. Indian J. Orthopaed. 1987, 1: 95-104.
- 10) Joshi MC, Patel MB, Mehta PJ. Some folk medicines of Dangs, Gujarat State. Bull. Medico-ethnobot. Res. 1: 1980, 301-317.

- 11) Singh KK, Maheswari JK. Traditional Therapy amongst the tribals of Varanasi District, Uttarpradesh. J. Econ. Tax. Bot. Bot. 4: 1983, 829-838.
- 12) Singh K. K. & Maheswari J.K. Forest in the life and Economy of the Tribals of Varanasi district, U.P. J. Econ. Tax. Bot. 6: 1985,109-116.
- 13) Jain, S.P. Ethnobotany of Morni and Kalesar (Anbala-Haryana). J. Econ. Tax. Bot. 5: 1984, 809-813.
- 14) Maheswari. J. K. , Kalakoto B. S. & Brijlal. Ethnomedicine of Bhil Tribe of Jhabua district, Madhya Pradesh. Anc. Sci. Life 5: 1986, 255-261.
- 15) Sharma, P.K., Dhyani S. K. & Shanker V. Some useful and medicinal plants of the district Dehradun and Siwalik . J. Sci. Res. Pl. Med. 1: 1979, 17-43.
- 16) Goel,A.K. Sahoo & Mudgal V. A contribution to the ethnobotany of Santal Pargana, Bihar, 1984, Bot. Surv. India, Howrah.
- 17) Sebedtian, M.K. & Bhandari M.M. Medico ethnobotany of Mount Abu, Rajasthan, India. J. Ethnopharmacol. 12: 1984, 223-230.
- 18) Maheswari, J.K., Singh K. K. & Saha S. Ethnomedical uses of plants by Tharus in Kheri district, U.P. Bull. Medico-ethnobot. Res. 1: 1980,318-337.
- 19) Jain, S. K. Medicinal Plantlore of the tribals of Bastar. Econ. Bot. 19: 1965,236-250.
- 20) Shah, G.L. Some economically important plants of Salsette island near Bombay. J. Econ. Tax. Bot. 5: 1984. 753-765. Tribedi, G. N., R. N. Kayal & H. N. Rai Chaudhuri. Some medicinal plant in Mayur bhanj (Orissa). Bull. Bot. Surv. India 24: 1982.,117-120.
- 21) Saxena, H. O., Brahman M. & Dutta P. K.. Ethnobotanical studies in Orissa, 1381,232-244, In S. K. Jain (Ed.) Glimpses of Indian Ethnobotany.
- 22) Upadhaye, A., Kumbhojkar M. S. & Vartak V. D. Observation on wild plants used in folk medicine in the rural areas of the Kolhapur district. Anc. Sci. Life 6: 1986, 119-121.
- 23) Shah, G. L., Menon A. R. & Gopal G. V. An account of ethnobotany of Saurashtra n Gujaratstate (India). J. Econ. Tax. Bot. 2: 1981, 173-182.
- 24) Srivastava, D. K. & Varma S. K., An ethnobotanical study of Santhal pargana, Bihar. Indian Forester 107: 1981, 30-41.
- 25) Chaudhuri, Rai H. N., Molla H. A., D. C. Pal & B. Roy , Plants used in traditional medicine by some tribals of Jalpaiguri district, West Bengal. Bull. Bot. Surv. India 24: 1982,87-90.
- 26) Sharma BD , Malhotra S. K. A contribution to the ethnobotany of tribal areas in Maharastra, J. Econ. Tax. Bot. 5: 1984, 533-537.
- 27) Gaur, R.D., Sharma M.P & Semwal J. K. Ethnotoxic plants of Garhwal hills in India. East. Anthropol. 33: 1980,159-163.
- 28) Maji, S. & Sikdar J.K. A taxonomic survey and systematic census on the edible wild plants of Midnapore district, West Bengal. J. Econ. Tax. Bot. 3: 1982, 717-737.
- 29) Jain, S. K. 1963. Studies in Indian Ethnobotany –Less known uses of 50 commonplants from tribal areas of Madhya Pradesh . Bull Bot. Srv. India. 5:223-226.
- 30) Jain, S.k. Wild plant foods of the tribals of Bastar (Madhya Pradesh) . Proc Nat. Inst. Sci. India. 30 B: 1964, 56-80.
- 31) Jain, S.K. Plant resources in tribal areas of Bastar, Vanya jati 12: 1964, 147-173.
- 32) Jain S. K. & De J. N. Observations on ethnobotany of Purulia district , West Bengal . Bull. Bot. Surv. India 8: 1966, 237-251.
- 33) Dr. Koppula Hemadri ,“A Treatise on tribal medicine” ; Dr. Koppula Hemadri’s House of Tribal Medicine, Vijayawada-520 008,2011.
- 34) Doshi T.R ,The study of Medhya Rasayana drug (Vacha & Jyotismati) w.s.r. to their effect on depression , IPGT & RA ,Gujarat Ayurved University ,Jamnagar; 1991.
- 35) Pawar P.S. ,A pharmaco clinical study on jyotismati in the management of amvata, IPGT & RA , Gujarat Ayurved University ,Jamnagar;2001.
- 36) Gaitande,B.B. et al.,Pharmacological activity of Malkanguni an indigenous tranquilizing drug (Preliminary reports), Curr. Med. Pract., Vol.1, 1957,PP-169.
- 37) Bidwai,P.P.; Wangoo. D. & Bhullar, N. Antispermatic action of *Celastrus paniculatus* Seed extract in Rat with reversible changes in the liver, J. Ethno. Vol.28(3) 1990,PP, 293-303.
- 38) Wangoo, D., Antispermatic effects of *Celastrus paniculatus* seed extract on the tests of albino rats .Eitoterapia, 1988,Vol59(5),PP- 377-382.
- 39) Pavanand .K. et al Schizontocidal activity of *Celastrus paniculatus* Willd. Agaist Plasmodium falciperum in vitro, Phytotherapy Res, 1989, Vol 3,PP 136-139.
- 40) Dandia , East Pharm, 1990, 33(396) , 39, Ahumada et al. Phytoter Res, 1991,5, 29.
- 41) Nalini K., Karanth K. S., Rao A, Aroor A.R., Journal of Ethnopharmacology.,1995, 47, 101.
- 42) Gattu M., Boss K.L., Terry A. V. , J. J. Buccafusco, Pharmacol. Biochem. Behav.,1997, 57, 793.
- 43) Kumar M. H.,Gupta Y. K., Phytomedicine,2002, 9 ,302.
- 44) Singh N., Chand N., Kohli R.P.,J. Res. Indian Medicine,1974, 9, 1.
- 45) Joglekar G. V., Balwani J.H., J. Res. Indian Medicine, 1967, 2, 190.
- 46) Kakrani H. K., Nair G.V., Kalyani G. A., Satyanarayana D., Fitoterapia, 1985,56, 293.
- 47) Mathur N. T., Verma M., Dixit V. P, Indian Drugs, 1993, 30, 76.
- 48) Pavanand K. et al., Phytoter. Res.,1989, 3, 136.
- 49) Russo A.,Izzo A.,Cardile V.,Borrelli F.,vanilla A., phytomedicine,2001, 8, 125.
- 50) P.C. Sharma, M.B. Yelne, T. J. Dennis, Database on Medicinal Plant Used in Ayurveda by CCRAS, New Delhi, 2005.
- 51) M. Bhanumathy, S. B. Chandrasekar, Uma Chandur, T. Somasundaram, Phyto-pharmacology of *Celastrus paniculatus*: An Overview, International Journal of Pharmaceutical Sciences and Drug Research 2010; 2(3): 176-181.